

# LIFETIME INTRA-CONSTELLATION CALIBRATION OF THE RAPIDEYE SENSORS OVER PICS

Cody Anderson, Michael Thiele, Dr. Andreas Brunn

Calcon 2013



RapidEye | www.rapideye.com September 2, 2013

### OUTLINE



- Background
- Procedure
  - Individual Tile
  - Cumulative Site
  - Complete Satellite-Band
- Full Constellation Satellite-Band Comparison
- Conclusions




# **BACKGROUND**

### THE RAPIDEYE CONSTELLATION

RapidEye
Delivering the World


- 5 optical satellites.
- 5 spectral bands.
- 6.5 m GSD, 77km swathwidth.
- >5 million sq km/day imaging capacity.
- Daily revisit of any target possible!!



# RAPIDEYE'S TILE CONCEPT




- RapidEye uses a 25km x25km tile reference system.
- RapidEye's Pseudo-Invariant Calibration Sites (PICS) are defined using these tiles.
- An image take may not cover an entire cal-site.



# RAPIDEYE'S 26 PICS








# **PROCEDURE**


Individual Tile
(Preformed for each satellite and band on an individual basis)

### INDIVIDUAL TILE FILTERING



# Samples per year (6 minimum)







# INDIVIDUAL TILE FILTERING (CONT.)



Simple mean + stand. dev. threshold.







# INDIVIDUAL TILE FILTERING (CONT.)



# Median/Mean Filter ± Shrinking R<sup>2</sup>.






### INDIVIDUAL TILE LINEAR FIT



Linear Least Squares Fit of filtered tile means.






# **PROCEDURE**

Cumulative Site (Preformed for each satellite and band on an individual basis)

### FILTER ALL TILES IN A SITE





This Cal-Site Includes 28 Tiles

### LINEAR FIT ALL TILES IN A SITE

100 2009


2010 2011

Date

2012

2011





2011

Date

27 Tiles Pass the Individual Filter Stage for this Cal-Site

### FILTER TILE LINEAR FITS

110 Weg 110

2011 2012

Date

100 L 2009

2010 2011 2012


Date

100 <del>---</del> 2009

2011

Date





Linear fits are filtered by mean of the slope ± std. dev. <u>17 Tiles Pass this stage</u>.

### NORMALIZE TILE LINEAR FITS

100 <del>---</del> 2009

2011

Date

110 Weg 110

2011 2012


Date

100 L 2009

2010 2011 2012

Date






Divide individual tile means by the linear fit value at a certain calibration zero point

### CAL-SITE LINEAR FIT





2010

RapidEye | www.rapideye.com September 2, 2013 Slide 17


2011

Date

2012

# CAL-SITE LINEAR FIT (CONT.)







# **PROCEDURE**

Complete Satellite-Band (Preformed for each satellite and band on an individual basis)

### **CAL-SITE FILTERING**

2009


2010

2011

Date

2012







The Individual Cal-Sites with a linear fit RMSE > the mean RMSE + the std. dev are filtered.

### TOTAL COMBINED LINEAR FIT








# TOTAL COMBINED LINEAR FIT (CONT.)








# FULL CONSTELLATION SATELLITE-BAND COMPARISON


### **BLUE BAND**






### **GREEN BAND**





### **RED BAND**






### **RED-EDGE BAND**








### **NIR BAND**







# **CONCLUSIONS**

### CONCLUSIONS



- There is some seasonal variation in the cal-sites.
  - Analyzing over an entire year helps minimize this effect.
- The RapidEye fleet has been quite stable since starting operations.
  - Maximum drop for one satellite-band is RE 4 Band 4 (3.90%)
- All Satellite Bands agree well with each other.
  - Maximum difference between Satellites is in Band 4 (3.24%).
- All degradations are accounted for in RapidEye products.



Cody Anderson: <a href="mailto:anderson@rapideye.com">anderson@rapideye.com</a>

Michael Thiele: thiele@rapideye.com

Dr. Andreas Brunn: <u>brunn@rapideye.com</u>

RapidEye :: Delivering the World

Germany | USA | Canada | Luxembourg

info@rapideye.com www.rapideye.com



RapidEye | www.rapideye.com September 2, 2013