



## Common Vitamin and Mineral Deficiencies in Utah

*Jeffery O Hall and D.R. ZoBell*

Department of Animal, Dairy, and Veterinary Sciences

### Introduction

Many minerals and vitamins have been proven in research studies to be essential for optimal growth, physiologic function, and productivity in animals. Data from the analytical section of the Utah Veterinary Diagnostic Laboratory would indicate significant incidences of vitamin and mineral deficiencies in Utah. There has been an increase in recent years that appears to be associated with producers decreasing or completely stopping the practice of vitamin-mineral supplementation. Based on interaction with producers they attribute some of this to high fuel, hay, and other production factors in which has resulted in searches for ways to cut input costs. A common finding with many of the diagnosed deficiencies is a lack of vitamin-mineral supplementation either long term or due to cost cutting.

Increasing incidence of adverse neonatal health effects, due to vitamin or mineral deficiencies, were observed in 2009, but further increases are occurring in 2010. This continued increase is likely due to some herds which stopped supplementation in the fall of 2008 having body reserves that allowed for healthy calves in the spring of 2009. But continued lack of supplementation has resulted in depletion of the body reserves and poor calf health in the spring of 2010. Even though this has just started to come to light, many producers may be using the wrong or a marginal mineral/vitamin supplementation program, which needs to be addressed and corrected.

This paper is directed at the health effect of common vitamin and mineral deficiencies and provides a summarization of the most commonly analyzed tissues and fluids that are used for diagnosing specific deficiencies. The paper also touches on immune system effects and appropriate supplementation.

### Deficiency Diagnoses

Historically, testing for deficiencies has been performed on diets and/or dietary components to ensure “adequate” concentrations in the diet. However, general mineral analysis does not identify the chemical forms of these minerals, which can dramatically alter their bioavailability and utilization. In addition, certain dietary factors can alter bioavailability of certain vitamins.

Although not possible for some of the minerals, the most specific means of diagnosing a mineral deficiency is by testing animals for unique functional deficits or deficiencies of specific mineral containing proteins or enzymes. This type of testing is often impractical from a field perspective, due to individual test costs or rigorous sample handling requirements. But, when possible, this type of testing eliminates the need to know the specific molecular characteristics of a dietary mineral and the potential for competitive interactions of antagonistic minerals for absorption/utilization. For minerals that do not have identified physiologic indices for which testing can be performed, direct quantification from animal tissues or serum may provide a reliable indication of the overall mineral status of the animal or group.

Testing of adequacy of fat soluble vitamins is commonly achieved by testing serum or liver tissue. It is essential that serum be separated from the red blood cells soon after collection. In addition, serum should be maintained frozen and protected from sun light while being shipped to the testing laboratory.

Vitamin and mineral deficiencies can be suggestively diagnosed by the development of clinical disease or by post-mortem identification of tissue lesions. But, proof of deficiencies often requires analytical verification since most do not have very unique clinical signs or lesions. In some instances, circumstantial proof of a

deficiency can be provided by positive response to supplementation of a suspected deficient vitamins or minerals. But, positive response may have nothing to do with the supplementation and may be just a time responsive correction of some other clinical condition.

An individual vitamin or mineral may have multiple means of measurement for identification of deficiencies, but most have one that is more specific than the others. For example, dietary concentrations may or may not be reflective of the amount that is bioavailable. Or, an individual tissue concentration may or may not reflect functional availability at the target or functional site.

The age of the animal being tested also is important for proper interpretation of status. For example, feti accumulate some minerals at different rates during gestation, necessitating adequate aging of the fetus for interpretation. In addition, some minerals, for which little is provided in milk, accumulate at higher concentrations during gestation in order to provide neonates with adequate body reserves for survival until they begin foraging. This is especially prevalent with copper, iron, selenium, and zinc. Thus, the “normal range” for these minerals in body storage tissues would be higher in early neonates than in an adult animal. One must be careful to make sure that the testing laboratory is interpreting the results based on the age of the animals tested, as some laboratories try to interpret all samples as if they were from adult animals.

When individual animals are tested, the prior health status must be considered in interpreting vitamin and mineral concentration of tissues. Disease states can shift mineral from tissues to serum or serum to tissues. For example, diarrhea can result in significant loss of sodium, potassium, and calcium from the body. Or, acidosis will cause electrolyte shifts between tissues and circulating blood. It is known that infectious disease, stress, fever, endocrine dysfunction, and trauma can alter both tissue and circulating serum/blood concentrations of certain minerals and electrolytes. Thus, evaluation of multiple animals is much more reflective of mineral status within a group than testing individual animals that are ill or have died from other disease states.

## **Live Animal Sampling**

A variety of samples are available from live animals that can be analyzed for vitamin-mineral content. The most common samples from live animals are serum and whole blood. These samples are adequate for measurement of several minerals, but it must be recognized that some disease states, as well as feeding times, can result in altered or fluctuating serum concentrations. Other samples from live animals that are occasionally used for analyses include liver biopsies, urine, and milk. But,

since milk mineral content can vary through lactation, vary across lactations, and be affected by disease it is not typically used to evaluate whole animal mineral status. Furthermore, hydration status significantly affects urinary mineral concentrations, rendering it a poor sample for evaluation of mineral status. For Vitamin A and E, serum is the best sample.

Serum should be separated from the red/white blood cell clot within the 1 to 2 hours of collection. If the serum sets on the clot for long periods of time, minerals that have higher intracellular content than serum can leach into the serum and falsely increase the serum content. Minerals for which this commonly occurs include potassium and zinc. In addition, hemolysis from both natural disease and due to collection technique can result in increase serum concentrations of iron, manganese, potassium, selenium, and zinc. Vitamin A and E can begin breaking down in serum if not separated from the red blood cells and frozen within 1-2 hours of collection. Serum for vitamin A and E analysis should be stored to prevent breakdown from sunlight exposure.

The best type of collection tube for serum or whole blood is royal blue-top vacutainer tubes, as they are trace-metal free. Typical red-top clot tubes will give abnormally increased results for zinc content as a zinc containing lubricant is commonly used on the rubber stoppers. For minerals other than zinc or vitamins A and E, serum samples from the typical red-top clot tubes are adequate. Similarly, serum separator tubes are typically adequate for vitamin-mineral analyses, except for zinc.

Samples should be appropriately stored for adequate sample preservation. Liver biopsies, urine, and serum can be stored frozen long term or refrigerated if mineral analysis is to be completed within a few days. Whole blood and milk should be refrigerated but not frozen, as cell lysis or coagulation of solids, respectively, will result.

## **Post-Mortem Animal Sampling**

A variety of post-mortem animal samples are available that can be analyzed for vitamin-mineral content. The most common tissue analyzed for mineral content is liver, as it is the primary storage organ for many of the essential minerals. In addition, bone is used as the primary storage organ for calcium, phosphorous, and magnesium. For Vitamin A and E, liver is the tissue of choice for analysis, but it needs to be relatively fresh. Tissue degradation will correspondingly decrease the vitamin A and E present.

Post mortem samples should be stored frozen until analyzed to prevent tissue degradation. If samples are to be analyzed within 1-2 days, they can be stored under refrigerated conditions.

## Copper Deficiency

Copper deficiency is one of the most commonly encountered nutritional problems in ruminants particularly in the intermountain west, but copper excess is also commonly encountered, especially in sheep. In contrast, copper deficiency is rare in non-ruminants. Clinical signs of deficiency can present as a large array of adverse effects. Reduced growth rates, decreased feed conversion, abomasal ulcers, lameness, poor immune function, sudden death, achromotrichia, and impaired reproductive function are commonly encountered with copper deficiency.

Cows will do all they can to ensure adequate copper is in calves when they are born. They will actually deplete their own body reserves to ensure neonatal adequacy. As such, neonates diagnosed with copper deficiencies are proof of maternal deficiencies. With copper being an essential component of the immune function, this maternal deficiency likely results in poor colostrums quality and inadequate neonatal protection even in calves that get adequate volumes of colostrums.

The best method for diagnosing copper status is via analysis of liver tissue, although much testing is performed on serum. Deficiency within a herd will result in some animals that have low serum copper concentrations, but serum content does not fall until liver copper is significantly depleted. In herds that have had livers tested and found a high incidence of deficiency, it is not uncommon for a high percentage of the animals to have “normal” serum concentrations. At the Utah Veterinary Diagnostic Laboratory, it is commonly recommended that 10% of a herd or a minimum of 5-10 animals be tested in order to have a higher probability of diagnosing a copper deficiency via serum quantification. Even with herd deficiency, low serum copper concentrations may only be seen in 10% or more of the individuals. Herds that may be classified as marginally deficient based on liver testing may have predominantly “normal” serum copper concentrations. Thus, serum copper analysis should be viewed as a screening method only. Another factor that can influence diagnosis of copper deficiency in serum is the presence of high serum molybdenum. As the copper-sulfur-molybdenum complex that forms is not physiologically available for tissue use, “normal” serum copper content in the presence of high serum molybdenum should always be considered suspect. In addition, the form of selenium supplementation can alter the normal range for interpretation of serum copper status, with selenite supplemented cows having a lowered normal range for serum copper.

Excessive supplementation of copper in dairy cattle is a relatively common finding at the Utah Veterinary

Diagnostic Laboratory. Liver copper concentrations greater than 200 ppm are routinely identified. But, in recent months, some cases of deficiencies have been identified, due to cessation of mineral supplementation programs.

The recommended adequate liver copper concentration range in adult cattle is 25 to 100 ppm. In comparison, late term fetal or neonatal liver should have 65 to 150 ppm copper to be considered normal.

## Manganese Deficiency

Manganese deficiency in ruminants is associated with impaired reproductive function, skeletal abnormalities, and less than optimal productivity. Cystic ovaries, silent heat, reduced conception rates, and abortions are reported reproductive effects. Neonates that are manganese deficient can be weak, small, and develop enlarged joints or limb deformities. Manganese deficiencies in beef cattle are most commonly seen in areas of highly alkaline soils, due to much poorer plant uptake of manganese.

Manganese deficiency, although not reported often, is identified routinely in dairy cattle when tested. Of interest is the fact that most testing of beef cattle (greater than 95%) finds normal manganese concentrations in liver, blood, and serum, but in these same matrices greater than 50%, 75%, and 95%, respectively, of dairy cattle tested are below recommended normal concentrations (unpublished data, Utah Veterinary Diagnostic Laboratory). This may, in part, be due to high calcium and phosphorous content of dairy rations, which can be antagonistic to the bioavailability of manganese.

Of the samples available, liver is the most indicative of whole body status, followed by whole blood and then serum. As red blood cells have higher manganese content than serum, hemolysis can result in increased serum content. Since the normal serum concentration of manganese is quite low, many laboratories do not offer this analysis because of inadequate sensitivity. Overall, response to supplementation has frequently been used as a means of verifying manganese deficiency, but it is critical that a bioavailable form be utilized.

## Selenium Deficiency

As an essential mineral, selenium is commonly identified as deficient in ruminants, but infrequently in dairy cattle. Selenium deficiency is also identified in many non-ruminant species. Selenium deficiency is associated with reduced growth rates, poor feed efficiency, poor immune function, impaired reproductive performance, and damage to muscle tissues. “White muscle disease,” a necrosis and scarring of cardiac and/or skeletal muscle, is linked to severe selenium deficiency;

although, it can be caused by vitamin E deficiency as well.

Cows will do all they can to ensure adequate selenium is in calves when they are born. They will actually deplete their own body reserves to ensure neonatal adequacy. As such, neonates diagnosed with selenium deficiencies are proof of maternal deficiencies. With selenium being an essential component of the immune function, this maternal deficiency likely results in poor colostrums quality and inadequate neonatal protection even in calves that get adequate volumes of colostrums.

Diagnosis of a deficiency can be made by analysis of liver, whole blood, or serum for selenium content or by analysis of whole blood for glutathione peroxidase, a selenium dependent enzyme, activity. The most specific analysis is that of whole blood glutathione peroxidase, as it verifies true functional selenium status. Liver is the optimal tissue to analyze for selenium content as it is a primary storage tissue. With serum and whole blood, the former better reflects recent intake, while the latter better reflects long term intake status. Since seleno-proteins are incorporated into the red blood cells when they are made and the cells have a long half-life, selenium content of whole blood is a better reflection of intake over the previous months than serum.

In order to adequately diagnose selenium deficiency, the dietary form of the selenium intake by the animals is important. Natural selenium, predominantly in the form of selenomethionine is metabolized and incorporated into selenium dependent proteins, but can also be incorporated into non-specific proteins in place of methionine. Inorganic selenium is metabolized and predominantly incorporated into selenium dependent proteins. Thus, "normal" concentrations in serum and whole blood differ depending on whether the dietary selenium is a natural organic form or an inorganic supplement.

The recommended adequate liver selenium concentration range in adult cattle is 0.25 to 0.50 ppm. In comparison, late term fetal or neonatal liver should have 0.35 to 0.65 ppm selenium to be considered normal.

## **Zinc Deficiency**

Zinc is an essential mineral that is required by all cells in animals. Zinc plays a role in numerous enzymatic reactions. Deficiencies of zinc are associated with reduced growth, poor immune function, diminished reproductive performance, and poor offspring viability, as well as skin lesions in severe cases.

Tissue zinc concentrations do not reflect body status well. Of the common samples tested, liver and serum are the best indicators of zinc status. But, serum and liver

zinc can be altered by age, infectious diseases, trauma, fever, and stress. Response to zinc supplementation has shown that some animals having low-end normal liver or serum zinc can still show improvement in some clinical conditions. Thus, liver and serum only verify deficiency when these samples have very low zinc content.

## **Vitamin A Deficiency**

Vitamin A is an essential fat soluble vitamin in ruminants. It is essential for all cell replications and is especially important in epithelial integrity. It plays an important role in tight junctions between cells, as well as being an important antioxidant in the body and in mucosal secretions. Vitamin A deficiency is associated with poor growth rates, poor feed intake, poor immune function, poor reproductive performance, and high incidences of diarrhea in calves. Loss of efficient tight junctions in the epithelial cell lining of the digestive tract allows opportunistic pathogens to invade and cause disease.

Vitamin A is provided in the diet via green growing vegetation or supplementation. Dead, brown forages have relatively no Vitamin A content. Thus, for grazing livestock, they must accumulate enough body reserves of vitamin A to carry them through the winter and have enough left to provide adequate vitamin A to their offspring. Therefore, it is more common to see vitamin A deficiencies in the springs after significant drought years, due to decreased time for body reserve accumulation. Unlike minerals, much of the vitamin A provided to the neonate is via the colostrums and in milk fats. Also, early calving has increased the incidence of neonatal vitamin A deficiencies due to lack of green forage at the time of parturition.

Vitamin A analysis can be efficiently performed on serum or liver tissue. It is important that samples be stored frozen and protected from light to prevent degradation of the vitamin A.

## **Vitamin E Deficiency**

Vitamin E is an essential fat soluble vitamin in ruminants. It is essential for all cells as an important antioxidant in the body in conjunction with selenium. Vitamin E deficiency is associated with poor growth rates, poor immune function, poor reproductive performance, poor muscle function, poor cardiovascular function, and "white muscle disease."

Vitamin E is provided in the diet via green growing vegetation or supplementation. Dead, brown forages have relatively no Vitamin E content. Thus, for grazing livestock, they must accumulate enough body reserves of vitamin E to carry them through the winter and have enough left to provide adequate vitamin E to their

offspring. Therefore, it is more common to see vitamin E deficiencies in the springs after significant drought years, due to decreased time for body reserve accumulation. Much of the vitamin E provided to the neonate is via the colostrums and in milk fats, although it is also transferred across the placenta. Also, early calving has increased the incidence of neonatal vitamin E deficiencies due to lack of green forage at the time of parturition.

Vitamin E analysis can be efficiently performed on serum or liver tissue. It is important that samples be stored frozen and protected from light to prevent degradation of the vitamin E.

## Effects on Immune Status

Deficiencies in vitamins and minerals have a two part impact on immune function in neonates. Firstly, since neonates are still developing their immune capabilities, these deficiencies have a direct negative impact on that development. And, indirect immune compromise is via the mother's poor immune function. At the time in which it is essential that the mothers be immune competent in order to produce antibodies for the colostrums, they often are deficient due to depletion from movement to the fetus. Additionally, poor immune function at the time of vaccination can result in very poor vaccine response, which in turn results in poor immune memory and antibody production. Thus, herd deficiencies would be expected to result in poor colostrums quality. This poor quality equates to a higher incidence of disease in the offspring due to poor maternal protection. Often this is seen as high incidence of neonatal diarrheas and/or high incidence of neonatal/juvenile pneumonias.

## Optimization of Supplementation

There are three basic time periods in which it is critical that vitamin-mineral status is optimal. Firstly, since the majority of minerals are transferred to the fetus during the last trimester of gestation, the 3 months prior to parturition are essential for offspring to be born with adequate body reserves and leave the mothers with enough to have good immune status for colostrum production. Secondly, as these vitamin-mineral deficiencies play a significant role in reproductive health, the time period from parturition to breeding is critical to ensure that the mother's system is replenished for optimal breed-back efficiency. And, thirdly, any time period in which animals are to be vaccinated, one must ensure adequate vitamin-mineral status in order to maximize response to vaccines. The authors routinely suggest animals be on a well balanced vitamin-mineral

supplementation plan for a minimum of 30 days prior to any vaccination.

## Summary

A variety of samples can be tested for vitamin-mineral content, but may not provide any indication of the overall mineral status of the animal. Appropriate diagnosis of mineral status involves thorough evaluation of groups of animals. The evaluation should include a thorough health history, feeding history, supplementation history, and analysis of several animals for their mineral status.

Dietary mineral evaluation should only be used to augment the mineral evaluation of animal groups. If minerals are deemed to be adequate in the diet, but the animals are found to be deficient, antagonistic interactive effects of other minerals need to be investigated. As an example, high sulfur or iron can cause deficiencies in copper and selenium even when there are adequate concentrations in the diet.

Overall, common vitamin-mineral deficiencies are significant hindrances to profitability in the livestock industry. Poor reproductive performance results in increased incidence of culling for open cows. Poorer than optimal feed efficiency and weight gain impact the bottom line in terms of pounds of cattle sales. And, poor calf health results in deaths and disease. The resultant increased disease incidence results in lost income in terms of treatment costs and poorer overall growth rates and gains in affected animals.

Any procedures mentioned in this article should be carried out by your veterinarian. A qualified nutritionist should be employed to determine mineral or vitamin requirements for your individual situation.

Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran's status. USU's policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions.

Utah State University employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran's status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USU-sponsored events and activities.

This publication is issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Noelle E. Cockett, Vice President for Extension and Agriculture, Utah State University.