

1970

Attendance Report: Small Mammal Workshop

Clive D. Jorgensen

Follow this and additional works at: http://digitalcommons.usu.edu/dbiome_memo

 Part of the [Earth Sciences Commons](#), [Environmental Sciences Commons](#), and the [Life Sciences Commons](#)

Recommended Citation

Jorgensen, Clive D., "Attendance Report: Small Mammal Workshop" (1970). *Memorandum*. Paper 97.
http://digitalcommons.usu.edu/dbiome_memo/97

This Article is brought to you for free and open access by the US/IBP Desert Biome Digital Collection at DigitalCommons@USU. It has been accepted for inclusion in Memorandum by an authorized administrator of DigitalCommons@USU. For more information, please contact dylan.burns@usu.edu.



IBP
DESERT BIOME

ATTENDANCE REPORT

Small Mammal Workshop

Helsinki, 1970

by

Clive D. Jorgensen
Center for Environmental Studies
Brigham Young University
Provo, Utah

April 1971

The material contained herein does not constitute publication. It is subject to revision and reinterpretation. The author requests that it not be cited without his expressed permission.

ATTENDANCE REPORT

Small Mammal Workshop: Helsinki, 1970

Clive D. Jorgensen, Brigham Young University, Provo, Utah

This meeting was the third since 1966, and included papers concerning almost all aspects of small mammal ecology. Most of the speakers addressed themselves directly to the problems of IBP, and although the species were different from those found in the desert, the problems and discussions concerned with them were rather helpful. Much of this report is included in a summary papers presented by Frank Golley and Lech Ryszkowski which also attempts to point out the direction of future work that should be undertaken by the international working group on small mammals. The proceedings of this meeting will be published as the third in a series, beginning with the 1966 proceedings, "Secondary Productivity in Terrestrial Ecosystems," and including the 1968 proceedings, "Energy Flow through Small Mammal Populations."

When the working group was first organized in 1966, two research programs were outlined for further work: (1) a standard or minimum program to provide information on the impact of small mammal populations on primary production, and (2) an advanced program to develop methods to investigate special problems associated with measuring energy flow in small mammal populations. In partitioning the effort, density determinations, sex ratios, body weight distributions, age distributions, and sexual activity were assigned to the first program; methods to determine densities, food habits, metabolism, and energy assimilation were specified for the second program. The efforts of this research have produced many helpful findings. A brief review of the most important of these (as far as the desert IBP work is concerned) may assist those who are looking for specific kinds of information.

Standing crop measurements have been reported from Bulgaria -- *Picea* -- *Fagus* mountain forest for *Clethrionomys glareolus* and *Apodemus flavicollis* (Markov et al., 1968); Czechoslovakia -- mixed deciduous forest for *C. glareolus* and *A. flavicollis* (Pelikan, 1968; Turcek, 1968); Finland -- meadow for *Microtus agrestis* (Myllymaki, 1969a), Poland -- cultivated fields for *Microtus arvalis* (Trojan 1969), peat bog for *Microtus oeconomus* (Bulchalczyk and Pucek, 1968), mountain meadow for *Pitymys subterraneus*, *M. agrestis*, and *A. flavicollis* (Grodzinski et al., 1966), Cladino -- Pinetum community for *C. glareolus* and *A. flavicollis* (Ryszkowski, 1969a, 1969b), Fagetum community for *C. glareolus* and *A. flavicollis* (Bobek, 1969), Tileo -- carpinetum community for *C. glareolus* and *A. flavicollis* (Aulak, 1967), and Cirecio -- Alnetum community for *C. glareolus* and *A. flavicollis* (Aulak, 1967); Rumania -- cultivated fields for *M. arvalis* (Hamar and Tuta, 1969); Sweden -- meadow for *M. agrestis* and a spruce plantation for *C. glareolus* (Hansson, 1969a); United States -- mixed deciduous forests for *Peromyscus gossypinus*, *Blarina berevicauda*, and *Ocotomys nutalli* (Gentry et al., 1968); and USSR -- several communities (Svarc et al., 1969).

A standard minimum method was proposed by Grodzinski et al. (1969) to obtain density estimates for small mammals. This method consists of pre-baiting for five days and then removal trapping of the resident population for six days on a 5.06 ha grid, which contains 256 points with two snap traps at each location. This method has been variously used, tested, and modified by Chelkowska and Ryszkowski (1967), Gentry et al. (1968), Hansson (1969b), Janion et al. (1968), Pelikan (1969), Ryszkowski (1969a), Ryszkowski and Petruszewicz (1967), Smirnov (1969), and Smith et al. (1969). Work has also been done to estimate the boundary effects of the grid using stained bait (Adamczyk and Ryszkowski 1968, Halisova 1969) and bait impregnated with radioisotopes (Myllymaki 1969b). Perhaps the most promising method of assessing border effect of kill trapping was proposed at the meeting by Michael Smith who used assessment lines, thus allowing for estimates of variance in the size of the grid. His method is also more flexible for varied communities and has been tested in the desert at the Rock Valley site in southern Nevada.

Data on natality, survival, and longevity are not numerous but have been reported for a number of species (Bujalska et al., 1968; Petruszewicz, 1968, 1969; Petruszewicz et al., 1969; Pucek et al., 1969; Ryszkowski, 1967; and Walkowa, 1969). These and other population parameters received considerably more attention at the Helsinki meeting, but still, very little data are yet available.

Progress has been made in the determination of energy parameters for certain species. Metabolic characteristics under laboratory conditions (Grodzinski and Gorecki, 1967; Grodzinski and Klekowski, 1968; Trojan and Wojciechowska, 1969), digestibility of food (Drozd, 1967; Petrides and Stewart, 1967), caloric values and body composition of different tissues (Golley, 1967; Gorecki, 1967; Evans, 1969), and growth (Petruszewicz et al., 1968) have been determined for a variety of species. These topics were also discussed during the Helsinki meeting but considerably more time was spent in discussing methods and rates of transfer within the ecosystem.

Mineral cycling, interactions of small mammals with other components of the ecosystem, and distribution characteristics have not been studied and a request for more of this type of work was made. It was felt that it was especially important that studies be expanded to include the deserts, the tundra, and the tropics.

The following statement of future plans is taken directly from the summary submitted by Frank Golley and Lech Ryszkowski.

It is appropriate at this stage in the development of the working group that we examine our progress in terms of the overall program of the IBP. At least one of the initial objectives of the IBP/PT program was to investigate whole ecological systems, with specialists contributing to the solution of whole system problems by investigating those groups of organisms in which they had special competence. The small mammal working group was one of these special groups and one of the final objectives of our investigations was to contribute to the understanding of the production dynamics of terrestrial ecosystems. Over time the ecosystem view has tended to be lost in our enthusiasm for special aspects of our studies and there has been little whole-system work accomplished, except in a few national programs. Since our working group has been highly successful and our knowledge of the impact of small mammal populations on production has advanced greatly, it may be useful for us to restate the basic underlying concept of the ecosystem IBP studies. Such a statement is especially appropriate today because it is widely understood that the environment of the planet Earth is deteriorating under the stress of expanding human populations and many people are looking to biologists and ecologists for solutions to environmental problems.

Man depends upon the ecosystems of the planet for sustenance, air, water, shelter, and for psychological needs. Because of the growing populations of human beings and more intensive use of ecosystems throughout the planet, it is essential that we develop sound plans for the use of the resources of the biosphere. Planning requires knowledge of ecosystem structure, function, and relationships. It is especially important that we have knowledge about how the most abundant and significant ecosystems influence the hydrologic cycle, and interact with the geosphere in the maintenance of soil fertility. We must also know the rates of loss of material from systems, the rates and form of production of ecosystems, the rates of energy flux through the system, and the resiliency of ecosystems to stress from the environment or man.

It is well known that ecological systems are composed of producers, which convert solar energy to chemical energy and are the initial step in the cycling of chemical materials in the system, decomposers, which break down organic material, and consumers, which probably function mainly in control processes on producers and decomposers. Small mammals are consumers and we should consider their broader role in control of ecosystem processes, as well as to continue our studies on the impact of populations on primary production. For example, a larger proportion of primary production channeled through the small mammal populations is expended as heat energy than that channeled through insect populations. In this respect, small mammals may function as a "steam engine" regulator in an ecosystem. Similarly, the formation of burrows by small mammals can influence the location and movement of organic matter in the soil. We should be concerned with such questions as how do mammals change the pathways or the rates of flow of energy and materials? How do mammals interact with other groups of consumers to influence these processes?

A focus of the small mammal working group such as we suggest has further value. Mammals are of economic concern as competitors with man for food and fiber and as disease vectors. History is replete with evidence that increase in density of man or mismanagement of wastes results in an increase in mammalian and other pest species. While control measures can be instituted, they are seldom successful over long periods of time. In contrast, an ecosystem approach, because it considers all aspects of the system, may provide the basis for successful management of pests, as well as other components of the system. In our opinion, effective and sustained management of our environment demands this total approach, rather than exclusive concentration on immediate, local solutions to specific problems.

If it is agreed that our working group should accept the ecosystem or biogeocenosis as the ultimate focus of our research, then an expanded program should be organized which takes this large concern into consideration as well as to further research those specific problems which we have investigated over the past five or ten years. First, we should support the Man and Biosphere program and become officially associated with it, if that is possible. Second, we should expand our research to other major ecosystems. Third, we should include studies on cultivated and urban-village ecosystems in our program. Fourth, we should expand our study of the interaction of mammal populations with other components of the system.

To this end, we suggest that the following reviews be prepared for presentation at a meeting of the working group in 1972. These papers would not only summarize the state of knowledge of small mammals but would also provide the information to design an expanded ecosystem-oriented program for the working group. We propose that the following papers comprise the main part of the meeting, with shorter comments and supporting or critical statements from other investigators included as comments in the published proceedings:

(1) density estimation and density in different habitats, (2) age indices and growth curves, (3) natality and survival rates in the natural environment, (4) bioenergetic parameters and foods, (5) energy flow and mineral cycling in different habitats, (6) the role of small mammals in food chains, in conditioning the environment, and in energy flow and mineral cycling of the whole system, (7) mammals in cities and in stores of food and fiber, and (8) mammals as disease vectors.

Environmental planning without consideration of the role of small mammals in ecosystems will be incomplete. We have an obligation to identify the kinds of information required for effective management of the earth and to supply that information appropriate to our specialty. A change in the emphasis of our working group does not necessarily change the kind of research carried out by the scientist in the field but, rather, it enlarges the view of the group and makes our studies more meaningful in a larger context.

Generally, the meeting was the most relevant I have attended and it helped my perspective greatly. Some of my general reactions are itemized below:

1. The small mammal working group does not seem to be coordinating well with ongoing Biome programs. I feel that the presence of Norman French and myself was helpful in pointing out the advantages of a closer association.
2. Modeling efforts were obvious by their absence. The general lack of equipment may account for much of this lack, but they have not, as yet, made a clear effort to coordinate with ongoing programs. This would be of considerable help. An invitation to them to contact the Desert Biome for assistance was extended by both Norman French and myself.
3. The group will meet again to concentrate on sampling methods. It is hoped that the basis for the proposed meeting will be real data and programs that can be submitted to a simulated population. The simulator will likely come from the efforts already begun in the Desert Biome at Brigham Young University.
4. Members of this group had made no effort to sample small mammals with live traps. I was the only person present with a proposal for live trapping. This proposal was well received and, if it had been in sufficiently final form, could have been included in the proceedings even though it was not included on the agenda. The group expressed a desire to see it as soon as possible so as to test it under European conditions.
5. The role of predators was discussed considerably. I decided that, generally, in the forests and meadows of Europe predators transfer an amazing quantity of energy and must be considered. We might also consider them in the desert.
6. I feel we should retain a working relationship with this group. The Desert Biome has much to offer, and we can learn such from an association. Also, I would like to request support for my own participation in future meetings.
7. Thank you for the opportunity; it has been of significant help.

LITERATURE CITED

- Adamczyk, K. and L. Ryszkowski. 1968. Estimation of the density of a rodent population using stained bait. *Acta Theriol.* 13: 295-311.
- Aulak, W. 1967. Estimation of small mammal density in three forest biotopes. *Ekol. Pol. A.* 15: 755-788.
- Bobek, B. 1969. Survival, turnover, and production of small rodents in a beech forest. *Acta Theriol.* 14: 191-210.
- Buchalczyk, T. and Z. Pucek. 1968. Estimation of the numbers of *Microtus alconomus* using the standard minimum method. *Acta Theriol.* 13: 461-482.
- Bujalska, G., R. Andrzejewski, and K. Petruszewicz. 1968. Productivity investigations of an island population of *Clethrionomys glareolus* (Schreber, 1780). II-Natality. *Acta Theriol.* 13: 415-425.
- Chelkowska, H. and L. Ryszkowski. 1967. Causes of higher abundance estimates of small rodents at the edges of sampling areas in forest ecosystems. *Ekol. Pol. A.* 15: 737-746.
- Drozd, A. 1967. Food preference, food digestibility and the natural food supply of small rodents. In secondary productivity of terrestrial ecosystems (ed. Petruszewicz) Warszawa - Krakow. p. 325-330.

- Evans, D.M. 1969. The effects of changes in nutrition on populations of *M. agrestis* L. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa. p. 157-158.
- Gentry, J.B., F.B. Golley, and M.H. Smith. 1968. An evaluation of the proposed International Biological Program census method for estimating small mammal populations. *Acta Theriol.* 13: 313-327.
- Golley, F.B. 1967. Methods of measuring secondary productivity in terrestrial vertebrate populations. In secondary productivity of terrestrial ecosystems. (ed. Petruszewicz) Warszawa. p. 99-124.
- Gorecki, A. 1967. Caloric values of the body in small mammals. In secondary productivity of terrestrial ecosystems. (ed. Petruszewicz) Warszawa-Krakow. p. 315-321.
- Grodzinski, W. and A. Gorecki. 1967. Daily energy budgets of small rodents. In secondary productivity of terrestrial ecosystems. (ed. Petruszewicz) Warszawa-Krakow. p. 295-314.
- Grodzinski, W., A. Gorecki, K. Janas, and P. Migula. 1966. Effects of rodents on the primary productivity of alpine meadows in Bieszczady Mountains. *Acta Theriol.* 11: 419-431.
- Grodzinski, W. and R. Z. Klekowski. 1968. Methods of ecological bioenergetics. Warszawa-Krakow. 252 pp.
- Halisova, V. 1969. A new method of marking small mammals with special coloured bait in IBP investigations. (unpublished manuscript).
- Hamar, M. and A. Tuta. 1969. Estimation of the density of *Microtus arvalis* (Pall) in clover fields by means of buried metal cylinders. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa. P. 81-88.
- Hansson, L. 1969a. The standard minimum method in grassland habitats. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa. p. 31-38.
- _____. 1969b. Home range, population structure and density estimates at removal caught with edge effect. *Acta Theriol.* 14: 153-160.
- Janion, M., L. Ryszkowski, and J. Wierzbowska. 1968. Estimate of number of rodents with variable probability of capture. *Acta Theriol.* 13: 285-294.
- Markov, G., L. Christov, and S. Gerasimov. 1968. Rodent trapping with use of different kinds of bait. *Small mammal newsletter* 2: 41-44.
- Myllymaki, A. 1969a. Productivity of a free-living population of the field vole, *Microtus agrestis*. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa. p. 255-265.
- _____. 1969b. Trapping experiments on the water vole *Arvicola terrestris* with the aid of the isotope technique. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa. p. 39-55.
- Pelikan, J. 1969. Testing and elimination of the edge effect in trapping small mammals. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa. p. 57-61.
- _____. 1968. IBP rodent programme in Czechoslovakia. *Small Mammal Newsletter* 2: 3-9.
- Petrides, G.A. and P.G. Stewart. 1969. Determination of energy flow in small mammals using chromium. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa p. 25-30.
- Petruszewicz, K. 1968. Calculation of the number of individuals born by a population. *Bull. Acad. Pol. Sci Cl. II* 16: 546-553.
- _____. 1969. Estimation of numbers of new-born animals. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa. O. 181-185.
- Petruszewicz, K., R. Andrzejewski, G. Bujalska, and J. Gliwicz. 1968. Productivity investigation of an island population of *Clethrionomys glareolus* Schreber. 1780. IV-Production. *Acta Theriol.* 13: 435-445.
- _____. 1969. The role of spring, summer and autumn generations in the productivity of a free-living population of *Clethrionomys glareolus* (Schreber, 1780). In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa p. 235-254.
- Ryszkowski, L. 1967. Short cut methods for the estimation of mean length of life in small mammal populations. In secondary productivity of terrestrial ecosystems. (ed. Petruszewicz) Warszawa-Krakow. p. 283-294.

- _____. 1969a. Operation of the standard minimum method. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa. 0. 13-24.
- _____. 1969b. Estimates of consumption of rodents populations in different pine forest ecosystems. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa p. 281-289.
- Ryszkowski, L. and K. Petruszewicz. 1967. Estimation of energy flow through small rodent populations. In secondary productivity of terrestrial ecosystems. (ed. Petruszewicz) Warszawa-Krakow. p. 125-146.
- Smirnov, V.S. 1969. Improvement of methods for estimating population size and survival of rodents. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa. p. 89-98.
- Smith M.H., J.B. Gentry, and F.B. Golley. 1969. A preliminary report on the examination of small mammal census method. In energy flow through small mammal census method. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa p. 25-30.
- Svarc, S.S., V.N. Bolsakov, V.G. Olnier, and O.A. Pjostolova. 1969. Population dynamics of rodents from northern and mountainous geographical zones. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa. p. 205-220.
- Trojan, P. 1969. Energy flow through a population of *Microtus arvalis* (Poll.) in an agrocenosis during a period of mass occurrence. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa. p. 267-279.
- Trojan, P. and B. Wejciechowska. 1969. Ecological model and tables of the daily costs of maintenance (DEB) of *Microtus arvalis* (Poll.) *Ekol. Pol. A.* 17: 314-342.
- Turcek, F.J. 1968. Small mammal secondary production study on mat-grasslands within the IBP in Czechoslovakia. *Small Mammal Newsletter* 2: 10-13
- Walkawa, W. 1969. Operation of compensation mechanisms in exploited populations of white mice. In energy flow through small mammal populations. (ed. Petruszewicz and Ryszkowski) Warszawa p. 247-253.