Natural Resources and Environmental Issues

Volume 15 Saline Lakes Around the World: Unique Systems with Unique Values

Article 22

1-1-2009

Planktonic bloom-forming Nodularia in the saline Lake Alchichica, Mexico

Maria Guadalupe Oliva

Proyecto de Investigacion en Limnologia Tropical, Universidad Nacional Autonoma de Mexico, Los Reyes

Alfonso Lugo

Proyecto de Investigacion en Limnologia Tropical, Universidad Nacional Autonoma de Mexico, Los Reyes

Javier Alcocer

Proyecto de Investigacion en Limnologia Tropical, Universidad Nacional Autonoma de Mexico, Los Reyes

Laura Peralata

Proyecto de Investigacion en Limnologia Tropical, Universidad Nacional Autonoma de Mexico, Los Reyes

Luis Alberto Oseguera

Proyecto de Investigacion en Limnologia Tropical, Universidad Nacional Autonoma de Mexico, Los Reyes

Follow this and additional works at: http://digitalcommons.usu.edu/nrei

Recommended Citation

Oliva, Maria Guadalupe; Lugo, Alfonso; Alcocer, Javier; Peralata, Laura; and Oseguera, Luis Alberto (2009) "Planktonic bloomforming Nodularia in the saline Lake Alchichica, Mexico," *Natural Resources and Environmental Issues*: Vol. 15, Article 22. Available at: http://digitalcommons.usu.edu/nrei/vol15/iss1/22

This Article is brought to you for free and open access by the Quinney Natural Resources Research Library, S.J. and Jessie E. at DigitalCommons@USU. It has been accepted for inclusion in Natural Resources and Environmental Issues by an authorized administrator of DigitalCommons@USU. For more information, please contact becky.thoms@usu.edu.



Planktonic Bloom-Forming Nodularia in the Saline Lake Alchichica, Mexico

María Guadalupe Oliva¹, Alfonso Lugo¹, Javier Alcocer¹, Laura Peralta¹ & Luis Alberto Oseguera^{1,2}

¹Proyecto de Investigación en Limnología Tropical, FES Iztacala, Universidad Nacional Autónoma de México, Av. De Los Barrios No. 1, Los Reyes Iztacala, 54090 Tlalnepantla, Edo. De México, México; ²Programa de Posgrado en Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México

Corresponding author:

María Guadalupe Oliva, Proyecto de Investigación en Limnología Tropical, FES Iztacala, Universidad Nacional Autónoma de México, Av. De Los Barrios No. 1, Los Reyes Iztacala, 54090 Tlalnepantla, Edo. De México, México E-mail: oliva@servidor.unam.mx

ABSTRACT

Lake Alchichica displays a characteristic cyanobacterial bloom associated with the onset of the stratification period (April-June). It mostly consists of the nitrogen-fixing cyanobacterium Nodularia spumigena. Filaments of Nodularia were straight, usually with an evident colorless, transparent sheath, with gas vesicles and without akinetes. The bloom developed in waters with a conductivity (K_{25}) of 13.47 to 14.14 mS cm⁻¹ (total dissolved solids around 8.7-9.2 g l⁻¹). Water column temperature was between 14.1 and 21.2°C, the pH fluctuated between 8.8 and 10.0, and dissolved oxygen between 0 and 9.3 mg 1⁻¹. The annual mean concentration of N-NO₃ was highest in 2001 (1.0 μM) and lowest in 2000 (0.4 µM). For P-PO₄ the highest mean value was measured in 2002 (0.8 μM) with the lowest value in 1999 (0.4 µM). We monitored the bloom throughout four annual cycles (1999-2002). The intensity of growth and extent of the presence of N. spumigena in Lake Alchichica differed among the years. The highest mean density was observed in 2001 (1.36 x 10¹¹ cells m⁻²), while the lowest values were found in 2000 (1.54 x 1010 cells m-2). Mean values for 1999 and 2002 were 2.08 x 1010 cells m-2 and 2.29 x 10¹⁰ cells m⁻², respectively. N. spumigena is found regularly during a three-month period, but peak concentrations are reached only for a few weeks. In calm weather and following solar heating, the Nodularia filaments floated to the lake surface, causing the bloom to become apparent. The intensity of growth was correlated with the annual concentration of N-NO₃, indicating the role of the cyanobacteria as a source of new nitrogen to the lake.

INTRODUCTION

Blooms of the nitrogen fixing cyanobacteria *Nodularia* spumigena Mertens have been reported around the world. They are common in the Baltic Sea (Komárek et al. 1993; Musial & Pliński 2003), in the North Sea and in brackish Elesmere (Nehring 1993), and in numerous estuaries and bays along the southern and western coast of Australia and New Zealand (Musial & Pliński 2003). Blooms have also been reported for North and South America: in Great Salt Lake (Felix & Rushforth 1979), in brackish coastal waters

and inland lakes and ponds in British Columbia, Canada (Nordin & Stein 1980), in Pyramid and Walker lakes in Nevada, USA (Cooper & Koch 1984; Galat et al. 1990), in Brazil (Werner & Rosa 1992), and in a shallow coastal lagoon in Uruguay (Pérez et al. 1999). Cyanobacterial blooms cannot be explained simply by the biological features of the species. The phenomenon is induced by many environmental factors which play an important limiting and controlling role. Among these factors Musial & Pliński (2003) and Mazur Marzec et al. (2006) mention phosphorus and nitrogen supply, as well as the N:P ratio, surface water temperature, light availability, calm weather, thermal stratification, and salinity.

The occurrence of Nodularia blooms in the hyposaline Lake Alchichica, Mexico, has previously been observed for several years, associated to the onset of the stratification period (April-June). Arredondo Figueroa et al. (1983) and Alcocer et al. (2000) reported Nodularia blooms in Lake Alchichica in the 1980s and 1990s (Oliva et al. 2001). Similar Nodularia blooms also develop in Lake Atexcac (Macek et al. 1994; Tavera & Komárek 1996), a nearby lake that shares ecological and morphological features with Lake Alchichica. According to personal observations and recorded data, the intensity and extent of the presence of Nodularia appeared to differ among the years. In the present paper we discuss the spatial and temporal dynamics of Nodularia during four annual cycles (1999-2002) in Lake Alchichica. The main physical and chemical variables are also presented.

AREA OF STUDY

Lake Alchichica is a deep crater lake located in the state of Puebla (19° 24' N and 97° 24' W), Central Mexico (Figure 1). The annual air temperature in the area fluctuates between 5.5 and 30°C with a mean value of 14.4°C. Having an arid climate, with annual precipitation of less than 500 mm and an annual evaporation rate of 1590 mm (Adame et al. 2008), this high altitude plateau (≥ 2300 m above sea level) named Los Llanos de San Juan can be described as a "cool desert".



Figure 1-Lake Alchichica, Puebla, Mexico.

The surface area of Lake Alchichica is 2.3 km² (diameter 1733 m). It has a volume of 9.42 x 10⁷ m³, a maximum depth of 62 m and a mean depth of 40.9 m (Filonov et al. 2006). The lake is warm monomictic (Alcocer et al. 2000). Mixing takes place from the end of December or the beginning of January until the onset of the stratification period by the end of April or beginning of May. A well-developed thermocline is present from June-July until October-November. After November, the thermocline deepens and becomes weaker until its breakup in late December or early January. Besides the spring bloom of *Nodularia*, Lake Alchichica displays a winter diatom bloom coinciding with the mixing period.

Lake Alchichica is a unique hyposaline $(8.5 \pm 0.2 \text{ g I}^{-1};$ Na-Mg and Cl-HCO₃ dominated) and alkaline (pH = 9.0 \pm 0.1) aquatic system (Vilaclara et al. 1993), characterized by several endemic species and unique features such as tufa towers. The endemic biota described include the atherinid fish *Poblana alchichica* (De Buen 1945), the ambystomatid salamander *Ambystoma taylori* (Brandon et al. 1981), the isopod *Caecidotea williamsi* (Escobar Briones & Alcocer 2002), and more recently the centric diatom *Cyclotella alchichicana* (Oliva et al. 2006).

MATERIALS AND METHODS

Sampling took place monthly at the central and deepest part of the lake during 1999–2002. Mid-day *in situ* profiles of temperature, dissolved oxygen, pH and conductivity (K₂₅) were obtained with a calibrated Hydrolab® DS3/SVR3 multiparameter water-quality data logger and logging system (discrete readings every meter). Ten water samples (depth 2, 5, 10, 15, 20, 25, 30, 40, 50 and 60 m) for phytoplankton analysis were obtained with a 51 Niskin-type water sampler. Two 500 ml sub-samples from each sampling depth were fixed, one with 4% formaldehyde and

the other with Lugol's solution (1%). Cyanobacteria were counted in a 50 ml settling chamber with a Zeiss inverted microscope following the Utermöhl method (APHA 1985; Wetzel & Likens 2000). The length and width of *Nodularia* filaments, vegetative cells and heterocytes were measured with an eyepiece micrometer. A hundred independent measurements were made for each cell type. The length of the filament measured was then divided by the length of the cells. This procedure was repeated several times with different filaments and an arithmetic mean was established (APHA et al. 1985). Integrated values for the water column were obtained using linear interpolation, and the total abundance of *Nodularia* was expressed per unit area (cells m⁻²) rather than per volume units since the phytoplankton rarely homogeneously distributed (Payne 1986).

Another set of samples was collected from the same depths for nutrient analyses. These samples were maintained in the cold (4°C) and in darkness until analysis. Nitrate and phosphate (soluble reactive phosphorus) analysis followed standard methods (Strickland & Parsons 1972), adapted by Kirkwood (1994) for a Skalar Sanplus segmented flow autoanalyzer system.



Figure 2–Nodularia spumigena straight filaments showing heterocytes. Scale bar = $10 \mu m$.

RESULTS

Nodularia in Lake Alchichica showed the following morphological traits: filaments were straight and composed of discoid vegetative cells 7-12 μm wide (mean 8.1 μm) and 2.7-3.6 μm long (mean 3.4 μm), with gas vesicles. Filaments usually had an evident colorless, transparent sheath. Heterocytes were 6.1-10.9 μm wide (mean 10 μm) and 4.5-6.4 μm long (mean 5.6 μm). They were present after every 12 to 16 vegetative cells (mean 13 cells) (Figure 2). Akinetes were never observed. According to its

morphological features and comparison with other descriptions (Nordin & Stein 1980; Pérez et al. 1999), our species corresponds to *Nodularia spumigena*. It is important to point out the existence of several morphotypes associated with local environmental conditions (Musial & Plinski 2003; Mazur Marzec et al. 2006). A definitive identification of *N. spumigena* requires studies that include ultrastructural and molecular biology considerations (Albertano et al. 1996; Hayes & Barker 1997). Such studies are still pending on the *Nodularia* of Lake Alchichica.

The K_{25} of Lake Alchichica varied from 13.47 to 14.14 mS cm⁻¹ (around 8.7-9.2 g l⁻¹) during the study period, confirming its hyposaline nature. The water column temperature ranged between 14.1 and 21.2°C, the pH fluctuated between 8.8 and 10.0, and dissolved oxygen between 0 and 9.3 mg l⁻¹ during the four years. Anoxic conditions were present in the deeper layers during the stratification period. Table 1 presents the N-NO₃ annual mean concentration, being highest in 2001 (1.0 μ M) and lowest in 2000 (0.4 μ M). For P-PO₄ the highest mean value was measured in 2002 (0.8 μ M) and the lowest value in 1999 (0.4 μ M). *N. spumigena* blooms in Lake Alchichica became apparent when calm weather and solar heating were present and filaments floated to the lake surface (Figure 3).

Table 1–Annual range and mean concentration values of N-NO₃ and P-PO₄³ in Lake Alchichica, 1999–2002.

Year	1999	2000	2001	2002
N-NO ₃ (μM) Range Annual mean value	0.04-4.2 0.7	0.01-1.9 0.4	0.06-5.2 1.0	0.01-4.8 0.5
P-PO ₄ (μM) Range Annual mean value	0.01-3.2 0.4	0.05-3.6 0.6	0.2-4.0 0.7	0.12-4.2 0.8

N. spumigena was the dominant species of the phytoplankton assemblage within the cyanobacteria bloom in all four years. The intensity of growth and extent of the presence of *N. spumigena* in Lake Alchichica differed among years (Figure 4). In 2001 the highest mean integrated column density was found (1.36 x 10¹¹ cells m⁻²), while 2000 had the lowest values (1.54 x 10¹⁰ cells m⁻²). Mean values for 1999 and 2002 were 2.02 x 10¹⁰ and 2.29 x 10¹⁰ cells m⁻², respectively. The maximum single-depth density was 1.05 x 10⁵ cells ml⁻¹ (surface water, May

30th 2001). The species was found along a four-month period (April to July) with peak concentrations lasting a few weeks only. In 1999 *Nodularia* was present for longer (7 months), including the mixing phase of the lake, when a diatom bloom is usually present. In 2002 the presence of *Nodularia* began in May and lasted until September.

We tried to establish a criterion to assess the presence of a bloom. The average value (4.64 x 10¹⁰ cells m⁻²), using all data when *Nodularia* was present, was used as a criterion for bloom conditions. Values above this average were considered as bloom. According to this criterion, blooms were present in June 1999, in May and June 2001 and in June 2002, but not in 2000 (Figure 4).



Figure 3–Accumulation of filaments during the *N. spumigena* bloom.

DISCUSSION

N. spumigena trichomes exist in different forms. In the Baltic Sea, N. spumigena displays three different forms: straight, coiled and spirally coiled (Komárek et al. 1993). In contrast, the population from Lake Alchichica is always straight, similar to other North and South American N. spumigena populations such as those in brackish coastal waters and inland lakes of British Columbia, Canada (Nordin & Stein 1980), in Walker and Pyramid Lakes, Nevada (Cooper & Koch 1984; Galat et al. 1990), in Great Salt Lake, Utah (Stephens 1990), in Rio Grande do Sul, Brasil (Werner & Rosa 1992), and in Castillos Lagoon, Uruguay (Pérez et al. 1999).




Figure 4–Interannual variation of densities of *N. spumigena* integrated over the water column in Lake Alchichica, 1999–2002. X = annual mean value.

Analyses of the 16S rRNA nucleotide sequence by Lehtimäki et al. (2000) indicated a great similarity between genotypes of the genus Nodularia. Although this gene is conserved and cannot therefore be used as a tool in the taxonomic identification of *Nodularia* species, there was a notable difference in the gene sequence between toxic and non-toxic Nodularia species. Laamanen et al. (2001) and Lyra et al. (2005) concluded that in the Baltic Sea there are only three species of *Nodularia*: one planktonic and toxin producing, with gas vesicles, which fits the description of N. spumigena, and two benthic, non-nodularin-producing species, without gas vesicles, namely N. sphaerocarpa and N. harveyana. The Lake Alchichica population is planktonic and contains gas vesicles; however, there is currently no evidence that this species produces toxins like N. spumigena from the Baltic Sea. Levels of nodularin will be tested in the future.

Some blooms of *N. spumigena* have been related to eutrophication processes (Hallegraeff 1993), although Pérez et al. (1999) mentioned that cyanobacterial blooms may be natural events and are not necessarily a consequence of eutrophication. The presence of *Nodularia* blooms in Lake Alchichica agrees with the last point of view, since this is an oligotrophic lake.

There are several factors that influence the growth and the extent of development of N. spumigena blooms in water bodies, the most important being salinity, temperature and light availability (Hamel & Huber 1985). N. spumigena requires salt concentrations between 5 and 20 g Γ^1 to be present. Maximum growth is achieved at salt concentrations from 5 to 10 g Γ^1 , especially when sulfate and sodium are dominant (Nordin & Stein 1980). Salinity in Lake Alchichica is in the range between 8.5 and 9 g Γ^1 , and its water contains high concentrations of sulfate and sodium (Vilaclara et al. 1993); both characteristics appear adequate

Nodularia growth requirements. High temperatures promote Nodularia blooms in saline and brackish waters (Lehtimäki 2000). According to Nordin & Stein (1980), best growth is achieved between 25 and 30°C. The highest temperatures in Lake Alchichica surface waters range between 18 and 21°C, below the optimum range. Nordin & Stein (1980) indicate pH values above 10.0 to be the best for Nodularia growth; Lake Alchichica's pH is close to this optimum value. Light availability is also an important factor for Nodularia to develop. From April to June, while Nodularia usually reaches its maximum densities, solar radiation showed the highest values (up to 1200 W m⁻²). In addition, Lake Alchichica is a transparent lake with a photic zone depth between 13 and 38 m (Adame et al. 2008). A short period that combines high solar radiation and calm weather appears to favor accumulation of the cyanobacteria at the lake surface, making presence of Nodularia clearly visible. Cloudy and windy weather prevents the cyanobacteria from accumulating at the water surface (Lehtimäki 2000).

Nitrogen and phosphorus concentrations and the N/P ratio are also important factors. Scarcity of nitrogen promotes development of nitrogen fixing cyanobacteria such as *Nodularia* (Paerl 1996). Due to its nitrogen-fixation capacity, *Nodularia* has a competitive advantage when N is the limiting nutrient for phytoplankton growth. The low dissolved inorganic nitrogen: soluble reactive phosphorus ratios (1.2-12.9, mean 6.71) found in Lake Alchichica by Adame et al. (2008) clearly suggests nitrogen to be the nutrient that most likely limits phytoplankton growth.

Several investigations have shown that low inorganic N concentrations precede blooms of nitrogen fixing cyanobacteria, while a decrease in phosphate levels has been observed in conjunction with blooms of *N. spumigena* (Cooper & Koch 1984). High concentrations of ammonium,

nitrate, and organic nutrients are released into the water during cyanobacterial decay (Engström-Öst et al. 2002). In Lake Alchichica, the highest N-NO₃ concentrations were coupled with the highest *N. spumigena* growth (2001), and the lowest NO₃. concentrations with the lowest densities (2000).

It is difficult to compare N. spumigena densities from different lakes, since values are given in different units (filaments ml⁻¹ or cells ml⁻¹) and are not calculated on an area basis (i.e., cells m⁻²). Maximum densities are commonly found at or close to the surface (around a depth of one meter) but in our case N. spumigena was usually found in the upper 10 m and some times throughout the entire water column (60 m). Lake Alchichica peak concentrations (1.05 x 10⁵ cells ml⁻¹) are much higher than those reported from Walker Lake (3.4 x 10⁴ cells ml⁻¹) by Cooper & Koch (1984) and from Pyramid Lake (3200 cells ml⁻¹) by Galat et al. (1981). Felix & Rushforth (1979) mentioned maximum densities of one thousand filaments per milliliter in Great Salt Lake. According to Paerl (1988), densities above 10⁴ to 10⁶ cells per milliliter of a single species could indicate the presence of a bloom. Following this criterion, N. spumigena bloomed in Lake Alchichica only in May 2001 with values between 4.1 x 10⁴ and 1.1×10^5 cells ml⁻¹ were observed in the top five meters. The rest of the time, densities were always below 10⁴ cells per milliliter.

N. spumigena appears to be an important source of nitrogen for Lake Alchichica. Falcón et al. (2002) measured acetylene reduction rates of up to $78.9 \pm 12.1 \mu \text{mol m}^{-2} \text{ h}^{-1}$ during the 2001 bloom. As mentioned previously, N-NO₃ concentrations after the 2001 bloom were the highest of the whole period, while the minimum N-NO₃ concentrations coincided with the lowest N. spumigena densities in 2000. This observation provides indirect support for the relevance of this species as a source of new nitrogen to the lake, as has been observed in other saline lakes (Hamilton Galat & Galat 1983). Nitrogen appears to be incorporated in the lake throughout the Nodularia bloom, and the nitrogen is released as ammonia during its mineralization by heterotrophic bacteria. Other modes of incorporation such as foraging are considered to be of low importance due to the fact that this species is poorly used as food for zooplankton (Hamilton Galat & Galat 1983).

In conclusion, while the water temperature is below its optimum range, the salinity, pH and light conditions explain the presence of *N. spumigena* in Lake Alchichica. The concentrations of nitrogen and phosphorus and their ratio determine the peak growth of *N. spumigena* during the onset of the early stratification. The intensity of *N. spumigena* growth appears to be related with the postpeak N-NO₃ concentration, indicating the role of the cyanobacteria as a source of new nitrogen to the lake through N₂ fixation and subsequent mineralization.

ACKNOWLEDGEMENTS

This research was supported by two grants from the Consejo Nacional de Ciencia y Tecnología (CONACyT): Limnoecología tropical: Interacciones cuenca de drenajelag. (CONACYT-41667) and Acoplamiento Física Lacustre-Procesos Biogeoquímicos en un Lago Tropical Profundo (CONACYT-34893-T). We thank Sergio Castillo (ICMyL, UNAM) for nutrient analyses.

REFERENCES

- Adame, M.F., J. Alcocer & E. Escobar. 2008. Size-fractionated phytoplankton biomass and its implications for the dynamics of an oligotrophic tropical lake. Freshwater Biology 53: 22–31.
- Albertano, P., D. Di Somma, D. Leonardi, A. Canini & M. Grilli Caiola. 1996. Cell structure of planktic cyanobacteria in Baltic Sea. Algological Studies 83: 29–54.
- Alcocer, J., A. Lugo, G. Vilaclara, M.R. Sánchez & E. Escobar. 2000. Water column stratification and its implications in a tropical, warm monomictic, saline lake Alchichica, Puebla, Mexico. Verhandlungen der Internationale Vereinigung für Limnologie 27: 3166–3169.
- American Public Health Association Washington (APHA). 1985. Standard Methods for the Examination of Water and Wastewater. American Public Health Association, Washington, DC, 1269 pp.
- Arredondo Figueroa, J.L., L.E. Borrego Enríquez, R.M. Castillo Domínguez & M.A. Valladolid Laredo. 1983. Batimetría y morfometría de los lagos maars de Cuenca de Oriental, Puebla, México. Biotica 8 (1): 37–47.
- Brandon, R.A., E.J. Maruska & W.T. Rumph. 1981. A new species of neotenic *Ambystoma* (Amphibia, Caudata) endemic to Laguna Alchichica, Puebla, Mexico. Bulletin of the Southern California Academy of Sciences 80: 112–125.
- Cooper, J.J. & D.L. Koch. 1984. Limnology of the desertic terminal lake, Walker Lake, Nevada, USA Hydrobiologia 118: 275–292.
- De Buén, F. 1945. Investigaciones sobre ictiología Mexicana. Anales del Instituto de Biología Universidad Nacional Autónoma de México 16: 475–532.
- Engström-Öst, J., M. Koski, K. Schmidt, M. Viitasalo, S.H. Jónasdóttir, M. Kokkonen, S. Repka & K. Sivonen. 2002. Effects of toxic cyanobacteria on a plankton assemblage: community development during decay of *Nodularia spumigena*. Marine Ecology Progress Series 232: 1–4.
- Escobar Briones, E. & J. Alcocer. 2002. *Caecidotea williamsi* (Crustacaea: Asellidae), a new species from a saline crate-lake in the eastern Mexican Plateau. Hydrobiologia 477: 93–105.
- Falcón, L.I., E. Escobar Briones & D. Romero. 2002. Nitrogen fixation patterns displayed by cyanobacterial consortia in Alchichica crater-lake, Mexico. Hydrobiologia 467: 71–78.
- Felix, E.A. & S.R. Rushforth. 1979. The algal flora of the Great Salt Lake, Utah. Nova Hedwigia 31: 163–195.

- Filonov, A., I. Tereshchenko & J. Alcocer. 2006. Dynamic response to mountain breeze circulation in Alchichica, a crater lake in Mexico. Geophysical Research Letters 33: L07404, DOI:10.1029/2006GL025901.
- Galat, D.L., E.L. Lider, S. Vigg & S.R. Robertson. 1981. Limnology of a large, deep, North American terminal lake, Pyramid Lake, Nevada, USA Hydrobiologia 82: 281–317.
- Galat, D.L., J.P. Verdin & L.L. Sims. 1990. Large-scale patterns of *Nodularia spumigena* blooms in Pyramid Lake, Nevada, determined from Landsat imagery: 1972–1986. Hydrobiologia 197: 147–164.
- Hallegraeff, G.M. 1993. A review of harmful algal blooms and their apparent global increase. Phycologia 32(2): 79–99.
- Hamel, K.S. & A.L. Huber. 1985. Relationship of cellular phosphorus in the cyanobacterium *Nodularia* to phosphorus availability in the Peel-Harvey estuarine system. Hydrobiologia 124: 57–63
- Hamilton Galat, K. & D.L. Galat. 1983. Seasonal variation of nutrients, organic carbon, ATP, and microbial standing crops in a vertical profile of Pyramid Lake, Nevada. Hydrobiologia 105: 27–43
- Hayes, P.K. & G.L.A. Barker. 1997. Genetic diversity within Baltic Sea populations of *Nodularia* (Cyanobacteria). Journal of Phycology 33: 919–923.
- Kirkwood, D.S. 1994. Sanplus Segmented flow analyzer and Its applications. Seawater Analysis. Skalar Co., Germany.
- Komárek, J., M. Hübel, H. Hübel & J. Šmarda. 1993. The *Nodularia* studies. 2. Taxonomy. Algological Studies 68: 1–25.
- Laamanen, M.J., M.F. Gugger, J. Lehtimäki, K. Haukka & K. Sivonen. 2001. Diversity of toxic and nontoxic *Nodularia* isolates (cyanobacteria) and filaments from the Baltic Sea. Applied and Environmental Microbiology 67: 4638–4647.
- Lehtimäki, J. 2000. Characterisation of cyanobacterial strains originating from the Baltic Sea with emphasis on *Nodularia* and its toxin, nodularin. Department of Applied Chemistry and Microbiology University of Helsinki, Finland. pdf-version, http://ethesis.helsinki.fi. Accessed 10 December 2007.
- Lehtimäki, J., C. Lyra, S. Suomalainen, P. Sundman, L. Rouhiainen, L. Paulin, M. Salkinoja-Salonen & K. Sivonen. 2000. Characterization of *Nodularia* strains, cyanobacteria from brackish waters, by genotypic and phenotypic methods. International Journal of Systematic and Evolutionary Microbiology 50: 1043–1053.
- Lyra, C., M.J. Laamanen, J.M. Lehtimäki, A. Surakka & K. Sivonen. 2005. Benthic cyanobacteria of the genus *Nodularia* are non-toxic, without vacuoles, able to glide and genetically more diverse than planktonic *Nodularia*. International Journal of Systematic and Evolutionary Microbiology 55: 555–569.
- Macek, M., G. Vilaclara & A. Lugo. 1994. Changes in protozoan assemblages structure and activity in a stratified tropical lake. Marine Microbial Food Webs 8: 235–249.

- Macek, M., G. Vilaclara, A. Lugo & J. Alcocer. 2007. Lago de Atexcac. In: De la Lanza, E.G. & P.S. Hernández. (eds), Las Aguas Interiores de México: Conceptos y Casos. AGT, México: 199–212.
- Mazur Marzec, H., A. Krężel, J. Kobos & M. Plínski. 2006. Toxic *Nodularia spumigena* blooms in the coastal waters of the Gulf of Gdańsk: a ten-year survey. Oceanologia 48: 255–273.
- Musial, A. & M. Plinski. 2003. Influence of salinity of the growth of *Nodularia spumigena* Mertens. Oceanological and Hydrobiological Studies 32: 45–52.
- Nehring, S. 1993. Mortality of dogs associated with a mass development of *Nodularia spumigena* (Cyanophyceae) in a brackish lake at the German North Sea coast. Journal of Plankton Research 15: 867–872.
- Nordin, R.N. & J. Stein. 1980. Taxonomic revision of *Nodularia* (Cyanophyceae/Cyanobacteria). Canadian Journal of Botany 58: 1211–1224.
- Oliva, M.G., A. Lugo, J. Alcocer, L. Peralta & M.R. Sánchez. 2001. Phytoplankton dynamics in a deep, tropical, hyposaline lake. Hydrobiologia 466: 299–306.
- Oliva, M.G., A. Lugo, J. Alcocer & A.E. Cantoral Uriza. 2006. *Cyclotella alchichicana* sp. nov. from a saline Mexican lake. Diatom Research 21: 81–89.
- Paerl, H.W. 1988. Nuisance phytoplankton blooms in coastal, estuarine and inland waters. Limnology and Oceanography 33: 823–847.
- Paerl, H.W. 1996. A comparison of cyanobacterial bloom dynamics in freshwater, estuarine and marine environments. Phycologia 35(6): 25–35.
- Payne, A.I. 1986. The Ecology of Tropical Lakes and Rivers. Wiley, New York.
- Pérez, M.C., S. Bonilla, L. De León, J. Šmarda & J. Komárek. 1999. A bloom of *Nodularia baltica-spumigena* group (Cyanobacteria) in a shallow coastal lagoon of Uruguay, South America. Algological Studies 93: 91–101.
- Stephens, D.W. 1990. Changes in lake levels, salinity, and the biological community of Great Salt Lake (Utah, USA) 1847–1987. Hydrobiologia 197: 139–146.
- Strickland, J.D.H & T.R. Parsons. 1972. A practical hand book of seawater analysis. Fisheries Research Board of Canada.
- Tavera, R. & J. Komárek. 1996. Cyanoprokaryotes in the volcanic lake Alchichica, Puebla State, Mexico. Algological Studies 83: 511–538.
- Vilaclara, G., M. Chávez, A. Lugo, H. González & M. Gaytán. 1993. Comparative description of crater-lakes basic chemistry in Puebla State, Mexico. Verhandlungen der Internationale Vereinigung für Limnologie 25: 435–440.
- Werner, V.R. & Z.M. Rosa. 1992. Cyanophyceae da estação ecológica do Taim, Rio Grande do Sul. Revista Brasileira de Biología 52: 481–502.
- Wetzel, R.G. & G.R. Likens. 2000. Limnological Analyses. Springer-Verlag, New York.