

2000

Western lentiform Leafminer

Diane G. Alston
Utah State University Extension

Michael E. Reding
Utah State University Extension

Follow this and additional works at: http://digitalcommons.usu.edu/extension_histall

 Part of the [Horticulture Commons](#)

Warning: The information in this series may be obsolete. It is presented here for historical purposes only. For the most up to date information please visit [The Utah State University Cooperative Extension Office](#)

Recommended Citation

Alston, Diane G. and Reding, Michael E., "Western lentiform Leafminer" (2000). *All Archived Publications*. Paper 423.
http://digitalcommons.usu.edu/extension_histall/423

This Factsheet is brought to you for free and open access by the Archived USU Extension Publications at DigitalCommons@USU. It has been accepted for inclusion in All Archived Publications by an authorized administrator of DigitalCommons@USU. For more information, please contact dylan.burns@usu.edu.



Orchard IPM Series HG/Orchard/04

Western Tentiform Leafminer *Phyllonorycter elmaella*

by Diane G. Alston and Michael E. Reding



Michigan State University

Underside of apple leaf showing mines of the western tentiform leafminer.

Do You Know?

- ◆ Western tentiform leafminer is an indirect pest that mines the leaves of apple and cherry.
- ◆ It can diminish the photosynthetic capability of trees and reduce fruit size and quality.
- ◆ Leafminer populations can fluctuate dramatically within and between years.
- ◆ Biological control by numerous, naturally occurring wasps is common.
- ◆ Successful management requires a good sampling program and knowledge of treatment thresholds:
 - 1 mine per leaf in early June
 - 2 mines per leaf in mid-June to July
 - 5 mines per leaf in August
- ◆ Numbers of mines on leaves should be sampled from June through August.
- ◆ Leafminers are resistant to many common insecticides.
- ◆ Insecticides have a detrimental effect on their natural biological control agents—parasitic wasps.

Western tentiform leafminer populations can vary tremendously between years or even between generations within a single year. Increasing resistance to organophosphate insecticides and variable effectiveness of at least six parasitic wasp species that attack the leafminer can influence the population size.

In Utah, the leafminer typically has three generations per year but may have as many as four in some years. The immature larval stages—sap feeders and tissue feeders—feed on apple and cherry leaves. When densities are high enough, feeding can cause a reduction in fruit size and quality.



Michigan State University

Sap-feeding larva inside mine.




Michigan State University

Adult moths are tiny ($1/8$ - $1/5$ inch long), and can be found resting on leaves and flying in the orchard.

Hosts

apple
cherry
pear
prune

Western Tentiform Leafminer Life History



The two arrows indicate when sap-feeding larvae should be monitored during the second and third generations, because if control treatments are necessary they should target this stage.

Life History

Pupa—Overwintering Stage

- ◆ **Size:** 1/8 to 1/5 inch long
- ◆ **Color:** yellow-brown changing to dark brown just before the adult emerges
- ◆ **Shape:** cylindrical tapering sharply at the rear end
- ◆ **Where:** overwintering pupae are inside mines in fallen leaves; summer pupae are inside mines in leaves on trees
- ◆ Summer pupation lasts 7–10 days
- ◆ Just before emerging, pupae cut through mine and protrude out the lower leaf surface
- ◆ Pupal skins remain attached to the mine after adults emerge and are called “emergence tubes”

Adult

- ◆ **Size:** 1/8 to 1/5 inch long, small, slender moth
- ◆ **Color:** forewings with golden brown and white bands edged with black; hindwings are gray with long hairs on the edges
- ◆ **Shape:** wings are held roof-like over the body when at rest
- ◆ **When:** first generation moths emerge from overwintering pupae in the early spring often around the green tip stage of apple
- ◆ May be up to three additional flights in the summer and fall; adult moths emerge continuously from June through September
- ◆ Life stages begin to overlap after the first generation
- ◆ Females lay eggs shortly after emerging

Egg

- ◆ **Size:** extremely small, about 1/100 to 1/80 inch in diameter and difficult to see without magnification
- ◆ **Color and shape:** transparent when first laid turning yellowish green; round and flat
- ◆ **Where:** on the undersides of leaves
- ◆ Hatches in 1–2 weeks

Larva—Damaging Stage

- ◆ Newly hatched larva mines down through egg case into leaf tissue
- ◆ Passes through five instars

Sap Feeder

- ◆ Instars 1–3
- ◆ **Size and color:** 1/12 inch long by the third instar, cream in color
- ◆ **Shape:** flat, wedge-shaped, and legless
- ◆ **Where:** feeds on sap from the spongy mesophyll layer of leaves with sucking mouthparts
- ◆ Separates outer layer of the leaf undersurface from the tissue above to form a mine, which often follows leaf veins
- ◆ Mines appear as a blotch and are only visible from the undersides of leaves at this stage

Tissue Feeder

- ◆ Instars 4–5
- ◆ **Size and color:** 1/8 to 1/6 inch long, cream turning yellow by the fifth instar
- ◆ **Shape:** cylindrical with legs, round head, and chewing mouthparts
- ◆ **Where:** feeds on leaf tissue just up to, but not through, the epidermis
- ◆ Attaches silken threads to both sides of the mine pulling them together to form a tent-like structure in the leaf
- ◆ Mines can be seen from the upper leaf surfaces and have a characteristic mottled appearance with white spots on top where green tissue has been removed
- ◆ Fifth instar larvae spin a silken chamber for pupating inside the mine

Host Injury

Larvae feeding on leaves can:

- ◆ Reduce photosynthetic capability of leaves
- ◆ Disrupt the growth regulating and ripening processes governed by hormones produced in the leaves
- ◆ Affect fruit quality by slowing sugar development
- ◆ Cause leaf drop, premature ripening, and fruit drop in severe infestations
- ◆ Increase susceptibility to chemical burn in heavily infested leaves
- ◆ Increase fruit sunburn if leaves surrounding fruit are curled or have fallen prematurely

Timing Control

Sampling should focus on the number of mines in leaves. The best time to sample for western tentiform leafminer larvae is at the beginning of the second generation, which usually occurs from mid-June to early July. Sampling earlier in the season for first generation eggs and sap feeder larvae is possible, but the eggs are difficult to locate and identify, and there is a poor relationship between the number of first generation eggs and the resulting mines.

Quick Evaluation Method

How to Sample

- ◆ Begin sampling in mid-June.
- ◆ Walk through the orchard and quickly scan tree terminals (leaf clusters at the ends of shoots) to estimate the number of mines per leaf.
- ◆ Sap feeders are the predominant stage at this time, so look on the undersides of leaves for mines.

Sampling Threshold

Second Generation: Mid-June To Early July

If terminals have:

< 1/2 mine per leaf

Return in 5–7 days to do another quick evaluation.

1/2–1 mine per leaf

Take a complete sample (see below). Sampling should continue through August.

Complete Sampling Method

Scouting for western tentiform leafminer can be done at the same time as mite sampling (see “Web Spinning Spider Mite” bulletin). If you have a large acreage of apples and cherries, you may want to select several representative blocks where you will conduct the scouting program. Randomly select 10 trees scattered throughout a 2- to 5-acre block to sample. If the orchard block is larger than 5 acres, it may be necessary to sample more trees.

How to Sample

- ◆ If the quick evaluation method indicates the need for a complete sample, sample for second generation leafminers during mid-June and early July.
- ◆ Scout designated orchard blocks every 1–2 weeks.
- ◆ On each sampling date, collect 10 leaves from each of the 10 trees (100 leaves total).
- ◆ Count the number of mines that you find and determine the average number of mines per leaf for the orchard block. (Divide the total number of mines by 100. See Western Tentiform Leafminer Sampling Form.)
- ◆ These same leaves can be used to monitor for mites.
- ◆ Third generation leafminers should be sampled in early to mid-August using the quick evaluation or complete method as indicated by the density of the mines present.
- ◆ If the second generation was treated or if sap feeders do not appear until late August or September, then it is not necessary to sample the third generation.

Management

Insecticides targeted for leafminer control are most effective against the sap feeder stage although parasitism is not usually evident until the tissue feeder stage. Therefore, parasite levels from earlier generations should be considered along with current sapfeeder mine densities during second and third generations to determine if treatment thresholds have been reached.

Treatment Thresholds

Leafminer Generation	Treatment Threshold
Second	
Early June	1 or more mines per leaf and parasitism of larvae is less than 30% ^a
Mid-June to July	2 or more mines per leaf and parasitism of larvae is less than 30% ^a
Third	
August	5 or more active mines per leaf ^b

^aOpen up a subsample of mines to look for parasites (see description below).

^bTo determine if mine no longer contains an active larva, look for presence of adult emergence tube attached to mine on lower leaf surface.

Insecticides

Chemical control is most effective against sapfeeding larvae in mines and adult moths flying in orchards. Insecticide sprays should only be applied when leafminer treatment thresholds are exceeded during the second or third generation (see above). Pre-bloom applications targeting first generation sapfeeders are only recommended if large numbers of adult moths are observed in the orchard during April or May, or egg counts exceed three per leaf during tight cluster to pink stage of apple.

The recommended insecticide is:

- ◆ oxamyl (Vydate)—controls adults and sapfeeders

Insecticide Resistance

Western tentiform leafminer has developed resistance to numerous organophosphate and organochlorine insecticides commonly used in orchards over the past 40 years. Oxamyl (Vydate) still appears to be an effective chemical but should only be used when necessary because it disrupts biological control of leafminers and other orchard pests.

Biological Control

There are at least six species of exoparasitic wasps that attack leafminers in the Northwest, and their presence in Utah orchards is common. Leafminer parasitism can be as high as 90% and can eliminate the need for chemical control of even a formerly high population. The wasps lay their eggs inside the mines on the outside of leafminer larvae. Wasp larvae hatch from the eggs and feed on developing leafminer larvae from the outside.

Parasitized Leafminer

To determine if a leafminer larva has been parasitized, open up a mine and look for one of the following:

- ◆ A wasp larva feeding on a deflated leafminer larva
- ◆ The pupal stage of the wasp along with a shriveled leafminer larva
- ◆ An empty mine with the remains of a leafminer larva and a wasp pupal case and no emergence tube formed by the adult leafminer moth

In the last case, the adult wasp has already emerged from its pupa and exited through a smaller hole, which is difficult to observe.

Insecticides to Avoid

Insecticides that are toxic to the parasites should be avoided during mid- to late summer if possible. In Washington studies, the following insecticides were shown to have moderate to high toxicity to *Pnigalio flavipes*, the main parasite of western tentiform leafminer in the Northwest:

- ◆ encapsulated methyl parathion (PennCap)
- ◆ chlorpyrifos (Lorsban)
- ◆ esfenvalerate (Asana)
- ◆ endosulfan (Thiodan)
- ◆ oxamyl (Vydate)

Conservation of parasites is important to a good biological control program and can reduce the need for insecticide control.

Utah State University Extension, an equal opportunity employer, provides programs and services to all persons regardless of race, age, gender, color, religion, national origin or disability. Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Robert L. Gilliland, Vice President and Director, Cooperative Extension Service, Utah State University.

Western Tentiform Leafminer Sampling Form—Apple and Cherry

Complete Sampling Method for Second and Third Generations

(Mid-June to August)

Orchard Block _____ Scout _____ Date _____

Number of Mines Per Leaf

	Leaf 1	Leaf 2	Leaf 3	Leaf 4	Leaf 5	Leaf 6	Leaf 7	Leaf 8	Leaf 9	Leaf 10	Tree Total
Tree 1											
Tree 2											
Tree 3											
Tree 4											
Tree 5											
Tree 6											
Tree 7											
Tree 8											
Tree 9											
Tree 10											
Total number of mines for 10 trees											
Average number of mines per leaf for 10 trees (divide above sum by 100)											per leaf

Instructions

1. Every 1-2 weeks, choose one or several representative orchard blocks for sampling.
2. Randomly select 10 trees scattered throughout a 2- to 5-acre block. If orchard is larger than 5 acres, it may be necessary to sample more trees.
3. On each sampling date, collect 10 leaves from each of the 10 trees (100 leaves total). These same leaves can be used to monitor for mites (refer to mite sampling form).
4. Count the number of mines and determine the average number of mines per leaf for each orchard block (to find the average, divide the total number of mines by 100).
5. If leaf mine counts are high, record counts from every leaf. Otherwise, record only total mine counts from all 10 leaves for each tree.

Treatment Thresholds*

Second Generation

Early June—one or more mines per leaf and parasitism less than 30%

Mid-June to July—two or more mines per leaf and parasitism less than 30%

Third Generation

August—five or more active mines per leaf

*See manual for more complete information