

1993

The Russian Wheat Aphid in Utah

Jay B. Karren
Utah State University

Follow this and additional works at: http://digitalcommons.usu.edu/extension_histall

 Part of the [Horticulture Commons](#)

Warning: The information in this series may be obsolete. It is presented here for historical purposes only. For the most up to date information please visit [The Utah State University Cooperative Extension Office](#)

Recommended Citation

Karren, Jay B., "The Russian Wheat Aphid in Utah" (1993). *All Archived Publications*. Paper 755.
http://digitalcommons.usu.edu/extension_histall/755

This Report is brought to you for free and open access by the Archived USU Extension Publications at DigitalCommons@USU. It has been accepted for inclusion in All Archived Publications by an authorized administrator of DigitalCommons@USU. For more information, please contact dylan.burns@usu.edu.





Utah State University Extension
Fact Sheet No. 80
February 1993

THE RUSSIAN WHEAT APHID IN UTAH

Introduction

Since arriving in Utah in 1987, the Russian wheat aphid, *Diuraphis noxia* (Kurdjumov), has spread to all grain growing areas of the state. It is very unpredictable in that at times it becomes an economic pest and at other times it is just present. In some areas it has caused losses in wheat and barley of up to 50 percent or more. It can be a problem in fall or spring planted grains.

Biology

Russian wheat aphids infest wheat, barley, and triticale, as well as several wild and cultivated grasses. Broadleaf plants such as alfalfa, clover, potatoes, and sunflowers are not hosts. Volunteer grain plays a key role in the life cycle of this pest by providing a food source in the interval between grain harvest and the emergence of fall-seeded crops. Many species of grasses act as reservoir hosts during the late-summer dry season; however, grasses such as barnyard grass and foxtail grass that grow on irrigation ditch banks and other wet waste areas are poor hosts. Most wild desert grasses are normally dormant and unsuitable for aphids during this period. In some cases, winged forms may feed on corn during heavy flights, but no colonization occurs.

In the summer, all Russian wheat aphids are females that do not lay eggs but give birth to live young at a rate of four to five per day for up to four weeks. The new young females can mature in as little as 7-10 days. Because of this reproductive ability, large infestations can spread rapidly. Throughout the year, overcrowding and weather conditions may stimulate the production of winged forms which are easily dispersed on wind currents. Russian wheat aphids prefer to live in leaf whorls and tightly rolled leaves that develop shortly after they begin feeding. As the colonies become crowded or the plant declines, wingless aphids move to neighboring plants. Russian wheat aphids survive Utah's cold winters as wingless nymphs or adults. They seem to be more cold-hardy than the other cereal grain species. They have overwintered in Utah every year since they were found, although the heavy snow, severe cold, or both during the 1988-89 and 1990-91 winters drastically reduced populations in most areas.

Identification

Following are the major characteristics by which the Russian wheat aphid can be identified

and separated from the other small grain aphids:

- Light green, elongated, and spindle shaped, not globular.
- Short antennae.
- Wartlike projection above tail (two tails).
- Cornicles very short and not obvious (no tailpipes on abdomen).

Detection and Damage

The characteristic damage of the Russian wheat aphid is the best way to identify its presence in a field. The aphids cause longitudinal leaf rolling and white (warm weather) or purple (cool weather) streaking on the leaves. The damage is caused by injection of a toxin into the plants as it feeds, which prevents the production of chlorophyll and causes the leaves to curl. Heavily infested plants are severely stunted and sometimes flattened. Later infestation and damage causes the flag leaf to curl and prevents the head from completely emerging. This produces a "gooseneck" head that does not allow proper grain maturation.

- 1. Surveying should begin when plants begin to grow in the spring.
- 2. Check at least 10 plants in 4 areas of each field.
- 3. Survey fields at least once, preferably twice, per week.
- 4. Generally each 1% infestation level will result in a 0.5% yield loss at harvest.

| Season | Plant Growth Stage | Threshold |
|--------|--|--|
| Fall | Seedling (1 tiller) | 10% of plants infested. |
| Fall | Larger plants | Treat if plants are stressed or there is danger of winter kill. |
| Spring | Winter grain--green to appearance of 1st node | 5% of plants with reproducing aphid colonies and fresh damage. |
| Spring | Winter grain--appearance of 1st node to head emergence | 10% of tillers infested. |
| Spring | Spring grain--emergence to head emergence | 10% of tillers infested. |
| Spring | Head emergence to soft dough | Treat only if heavy populations (i.e., more than 20 aphids per plant) develop on 10 to 20% of flag leaves or stems or if high populations are discovered in the heads (more than 3 per head at flowering, 8 per head at water stage, and 15 per head at milky ripe stage). |

After the soft dough stage, insecticides will have no benefit (do not spray).

Recent evidence from other states indicates that much of the reduction in yield attributed to fall infestations of Russian wheat aphids occurs because aphid feeding increases the susceptibility of the plant to winter injury. Feeding reduces the plant's cold tolerance and stunts root growth. As a result, fall crops in areas where winter kill is a common problem cannot tolerate as many Russian wheat aphids as those in areas where winter injury is

uncommon. Early detection is important. Listed below are recommendations for surveying and identifying Russian wheat aphid presence and damage in fields.

Barley Yellow Dwarf Virus

Unlike most other cereal aphids, Russian wheat aphids are currently not significant vectors of barley yellow dwarf virus (BYDV) in North America. This disease causes yellowing beginning at the tips of leaves and progresses towards the base. In contrast the Russian wheat aphid damage begins at the base and progresses towards the tip of the leaves in longitudinal white to yellow streaks. Soon after infestation, the leaves begin to curl protecting the aphids inside.

Control

Plant winter wheat late in the fall after trap collections indicate that late-season flights have subsided.

Plant spring wheat as early as possible to minimize crop exposure to aphids. Initial spring suction trap collections reflect the beginning of flight activity, so spring-planted grains should be monitored intensively after the trap catches begin to collect Russian wheat aphids. Contact your local county agent for suction trap information.

Precautionary Statement

All pesticides have both benefits and risks. Benefits can be maximized and risks minimized by reading and following the label. Pay close attention to the directions for use and the precautionary statements. The information on pesticide labels contains both instructions and limitations. Pesticide labels are legal documents, and it is a violation of federal and state laws to use a pesticide inconsistent with its labeling. The pesticide applicator is legally responsible for its proper use. Always read and follow the label.

Jay B Karren
Extension Entomologist