



## Candidatus Liberibacter of pepper

Authors: Claudia Nischwitz, Extension Plant Pathologist • Erin Petrizzo

### What you should know:

- The non-culturable bacterium *Candidatus Liberibacter solanacearum* infects peppers.
- Bacteria are transmitted by potato psyllids, (*Bactericera cockerelli*).
- Bacteria are transmitted throughout the lifetime of the insect and can also infect tomatoes and potatoes.



Fig. 1. Yellow discoloration and stunted growth of leaves.

### INTRODUCTION

The bacterium *Candidatus Liberibacter solanacearum* infects pepper plants. *Candidatus Liberibacter solanacearum* is found in all states west of the Rocky Mountains, Kansas and in Texas. The disease was first reported in the United States in 2004. Yield losses can reach up to 100%.

### SYMPTOMS

The symptoms on peppers include stunted growth, and yellow discoloration of the foliage (Fig. 1, Fig. 2, Fig. 3). The potato psyllids have been present in Utah for a long time but the bacterium was detected for the first time in a in 2013 (potato sample). In summer 2014, pepper samples were confirmed positive for the first time.

### DISEASE CYCLE

The bacterium is transmitted by potato psyllids (*Bactericera cockerelli*) (Fig. 4). Potato psyllids are very small and look like black, winged aphids on the plants. They are native to North America and found in the Southwestern United States.



Fig. 2. Healthy pepper plant (right). Diseased pepper plant (left)



Fig. 3. Yellow discoloration of pepper foliage.

Under a dissecting micro-scope the characteristic white bands on their back can be seen. Eggs (Fig. 5) are a light to dark yellow-orange color and can be found on short stalks near the edge of the leaf. Nymphs (Fig. 6) are only visible on the leaves with a strong hand lens. Potato psyllids can acquire *Candidatus Liberibacter solanacearum* by feeding on an infected plant. The bacterium can also be passed on from mother to offspring. Depending on the number of potato psyllids feeding, a potato plant can be infected within one to six hours. One adult potato psyllid is capable of infecting a healthy potato plant within six hours. Twenty adult potato psyllids are capable of infecting a healthy potato plant within an hour (Munyanenza 2012).



**Fig. 4.** Adult potato psyllid. (*Bactericera cockerelli*)  
Size: 0.12 inches (3 mm)

## MANAGEMENT

It is important to start scouting for potato psyllids using yellow sticky cards early in the season. Leaves should be inspected on the underside for the presence of psyllid nymphs (Fig. 6). The nymphs are very small and can be seen easier with a strong hand lens. If peppers are planted in an area where potato psyllids have been a problem in the past, imidacloprid insecticide can be applied at planting. If follow-up treatments are needed, insecticides containing spinosad or spiromesifen can be used following the label. Once a plant is infected with *Candidatus Liberibacter solanacearum*, there is no cure for it. It should be removed to reduce the amount of inoculum present in the field.



**Fig. 5.** Potato psyllid egg.  
Size: Less than 0.04 inches (1 mm)

## REFERENCES

Munyanenza, J. E. 2012 Zebra Chip Disease of Potato: Biology, Epidemiology and Management. Am. J. Pot. Res. 89:329-350



**Fig. 6.** Potato psyllid Nymphs.  
Size: 0.08 inches (2 mm)

<sup>4</sup>Image courtesy of <http://www.ag.ndsu.edu/archive/entomology/ndsucpr/Years/2010/August/12/ent.htm>

<sup>5</sup>Image courtesy of Kevin Ong, Texas AgriLife Extension Service, Bugwood.org

<sup>6</sup>Image courtesy of Kevin Ong, Texas AgriLife Extension Service, Bugwood.org

**Precautionary Statement:** Utah State University Extension and its employees are not responsible for the use, misuse, or damage caused by application or misapplication of products or information mentioned in this document. All pesticides are labeled with active ingredients, directions for use, and hazards, and not all are registered for edible crops. "Restricted use" pesticides may only be applied by a licensed applicator. The pesticide applicator is legally responsible for proper use. USU makes no endorsement of the products listed herein.

Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran's status. USU's policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions. Utah State University employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran's status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USU-sponsored events and activities. This publication is issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Dept. of Ag., Noelle E. Cockett, Vice President for Extension and Agriculture, Utah State University.