Utah State University DigitalCommons@USU

All Archived Publications

Archived USU Extension Publications

1-1-1996

Important Components of a Successful Pest Management Program

Diane G. Alston *Utah State University*

Warning: The information in this series may be obsolete. It is presented here for historical purposes only. For the most up to date information please visit The Utah State University Cooperative Extension Office

Recommended Citation

Alston, Diane G., "Important Components of a Successful Pest Management Program" (1996). *All Archived Publications*. Paper 767. http://digitalcommons.usu.edu/extension_histall/767

This Report is brought to you for free and open access by the Archived USU Extension Publications at DigitalCommons@USU. It has been accepted for inclusion in All Archived Publications by an authorized administrator of DigitalCommons@USU. For more information, please contact becky.thoms@usu.edu.



Integrated Pest Management Fact Sheet



IMPORTANT COMPONENTS OF A SUCCESSFUL PEST MANAGEMENT PROGRAM

Diane G. Alston, Extension Entomologist
Department of Biology
Utah State University, Logan UT 84322-5305

April 1996 AG/IPM/02

I. IMPORTANT FIRST STEPS

- **1. Pest Identification**—Before deciding to take any pest control action, be sure you have correctly identified the pest (insect, mite disease, weed vertebrate, etc.). Proper pest identification is essential for determining if control is necessary and for selection of the best suited control options.
- **2.** Surveying for Pests and Injury Caused by Pests—It is very important to look for pests and injury symptoms on a regular basis. It is best to use a consistent sample or survey method. Monitoring traps for insects, diseases, or vertebrates can be used when appropriate. Sampling frequency will depend on the pest and situation, but a general guideline for many insects and diseases is to sample every 1–2 weeks during the main growing season. Look in the places where the pests commonly occur— undersides of leaves, under bark, inside fruits, in soil, etc.

3. Control Action Guidelines:

Action Threshold: "The point at which a pest control action must be taken to prevent unacceptable damage to the ecosystem."


The pest manager should develop a set of guidelines for each situation which guides his/her decision-making on acceptable levels of pest injury. A low level of pest injury can perhaps be ignored if the loss will not justify the cost of control. The economic and aesthetic vales of the management situation need to be considered, and will most likely be the primary factors driving pest control decisions. Other considerations may be the environmental and social costs of pest control decisions.

[Refer to IPM Fact Sheet No. 3: "Pest Management Decision-making: The Economic Injury Level Concept" for more information.]

II. IDENTIFYING "WINDOWS OF OPPORTUNITY"

All types of pests have a life cycle or set of developmental events that occur during their life time. The type of life cycle will vary with the pest. However, most pests have certain weak

points or windows of opportunity" during their life cycle when they are the most vulnerable to control. For insects, these windows are often during the immature life stags. Weeds are typically easiest to control during their seedling stage, early in the season when they are just beginning to grow (annuals) or late in the season when they are preparing for dormancy (perennials). Diseases are often easiest to control by using preventive or early intervention tactics before the disease begins developing or becomes established. To optimize management of a pest, control tactics should be targeted for these weak points.



*Indicates weak points in pest life cycles that should be targeted for control; the number of ** indicates the general level of weakness.

III. GENERAL PEST MANAGEMENT TACTICS

The easiest, lowest cost and often most reliable way to avoid many pest problems is to provide a healthy environment that discourages pest activities and/or reduces the host's (plant, animal, or ecosystem) susceptibility to damage. In general, such methods are long-term stable pest suppression. For pests that directly attack essential or yield producing structures of the host (e.g., for plants: fruits, seeds, flowers,

Short-term vs. Long-term Tactics

Short-term: Long-term:
• Chemicals • Cultural

Mechanical -Habitat modifications

Biological
 Biological

stems, trunks), short-term tactics such as pesticides are often the quickest and most reliable pest control option. A combination of both short- and long-term tactics is often the best approach.

Examples of some general pest management tactics:

1. Cultural Controls

- **Land / Water Management:** Maintain the ecosystem in a healthy state to minimize the competitiveness of pests. In some situations water levels can be regulated to reduce pest problems (e.g., mosquitoes, aquatic weeds, etc.). Avoid over- or under-watering plants to minimize stress.
- **-Sanitation:** Remove (prune, gather, burn) plant parts and debris that can serve as protective or overwintering sites for many pests.
- **Habitat Diversification:** In monoculture situations (e.g., agricultural fields, orchards, landscapes, etc.), diversification of vegetation in the habitat may subsequently increase the diversity of animal life in the habitat. Because such diversification may attract beneficial or pest organisms, the specific approach should be tailored to the specific situation.

- Tolerant / Resident Species and Cultivars: Select insect and disease resistant species and cultivars of plants whenever they are available. For perennial plants, select species and cultivars that are cold hardy (for cold climates). A winter damaged plant or tree will be more susceptible to attack by pests.
- Soils and Nutrition: Although you cannot change the soil type in an area, there are some practices you can follow to improve growing conditions for plants. Proper fertilization is important, but over-fertilization can lead to excessive, lush growth that can be attractive to aphids and other foliage pests. Increasing organic matter of soil where it is low can improve the growth and health of plants.

2. Mechanical Controls

- **Hand Removal:** Hand-pulling of weeds or other pests physically removes them from the situation.
- **Mowing:** Mowing down weeds, especially before they produce seeds, can be a very effective management tool.
- **Traps:** Attractive traps can be used to mass trap and remove pests from the environment or to monitor their activity so that control actions can be appropriately timed. Traps are primarily used for insect and vertebrate pests, and for some disease pests. Attractive traps typically use visual cues (color) and/or odors (pheromones, food baits) to attract pests to them. Passive traps may be placed in areas of common pest activity or use wind currents to catch pests.
- **Physical Barriers:** There are many types of physical barriers that can be used to block or disrupt the movement of pests. For example, sticky bands placed around the trunks of trees can prevent some mite and insect pests from crawling up into trees.

3. Biological Controls

Biological Controls: "Any activity of one species that reduces the adverse effect of another."

- **-Predators:** an organism that eats or consumes another (e.g., predatory insects and mites).
- **Parasites:** an organisms that lives in or on another and kills it (parasitoid) while completing its life cycle (e.g., parasitic insects).
- **Pathogens:** microbial agents that attack and invade other organisms (bacteria, viruses, nematodes).
 - Herbivorous insects of weeds: plant-feeding insects that attack weeds.

Biological control agents can be purchased from commercial suppliers and released for supplementary control of pests. Biological control releases can be inundative (flood an area with the agents to reduce the pest in the short-term) or inoculative (release agent over a period of time to help it colonize and spread). However, most biological control occurs without assistance from people. Many predators, parasites and pathogens occur naturally and are continually working to help keep nature in balance. The importance of "natural enemies" is often not appreciated until a broad spectrum pesticide, which kills many beneficials as well as the targeted pest, is applied and a new pest—suddenly released from biological control—become a serious problem.

There are things you can do to encourage the activities of biological control agents already present in the environment:

- -Avoid the use of broad spectrum pesticides unless necessary. Use selective pesticides and target them to pest-problem spots.
- -Provide a habitat that is more favorable for natural enemies (see Habitat Diversification under Cultural Controls) by choosing adjacent plants that supply nectar, pollen, alternate hosts,

and structural protection. Avoid complete removal of habitat for beneficials (field borders, adjacent crops, ground covers, etc.).

- Provide adequate organic matter in soil to ensure good habitat for soil microorganisms that control a variety of pests, particularly soil pathogens and root-feeding nematodes.

[Refer to IPM Fact Sheet No. 4: "General Concepts of Biological Control" for more information on this topic.]

4. Chemical Controls

Pesticide: "Any substance applied to control insects, fungi, bacteria, weeds, vertebrates or other pests."

Since the advent of "synthetic" pesticides in the 1940s, pest managers have become very reliant on their generally simple-to-use, fast-acting, and effective attributes to manage the majority of pest problems. In recent times, resistance of pests to pesticides because of their high use has become a major problem. To reduce the reliance on pesticides as a single-pronged approach to pest management, pest managers should become more familiar with all types of pest control tactics, so that the best one(s) can be selected for each situation. Although pesticides have been perhaps overused in recent decades, they still remain an important tool in the pest management toolbox. Avoidance of overuse will allow them to remain viable pest control tools for many more decades to come.

Many types of chemicals used before the development of synthetic pesticides are becoming popular once again as alternative or "organic" pest controls. In addition, many new chemical products are being developed and made available, such as "biologicals" and "insect growth regulators."

- **Synthetic Pesticides:** human-made in the laboratory; chemically joined compounds or elements (e.g., most herbicides, diazinon, malathion, carbaryl, benomyl, streptomycin, etc.)
- **Organic Pesticides:** Derived from plant, animal, or naturally occurring rock or petroleum oil sources (e.g., rotenone, sabadilla, ryania, nicotine sulfate, pyrethrum, soaps, oils, microbial agents, lime sulfur, copper, etc.)
- **Biological Pesticides:** A subset of organics that specifically refer to products developed from naturally occurring microbial agents such as bacteria, viruses, and fungi (e.g., *Bacillus thuringiensis*, a bacterial pathogen of many insects.)
- Insect Growth Regulators (IGRs): Kill insects by interfering with the normal processes of juvenile development; common IGRs disrupt either the insect's hormonal process or exoskeleton development; a few synthetic replicas of natural IGR chemicals are commercially available.

Utah State University Extension and its employees are not responsible for the use, misuse or damage caused by application or misapplication of the products or information in this publication, and make no endorsement explicitly or implicitly of this publication or information listed herein.

Utah State University Extension is an affirmative action/equal employment opportunity employer and educational organization. We offer our programs to persons regardless of race, color, national origin, sex, religion, age or disability.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Robert L. Gilliland, Vice-President and Director, Cooperative Extension Service, Utah State University, Logan, Utah. (EP/DF/09-97)