

Comparison of Coincident Rayleigh-Scatter and Sodium Resonance Lidar Temperature Measurements from the Mesosphere-Lower-Thermosphere Region



Leda Sox, Vincent B. Wickwar, Tao Yuan and Neal R. Criddle

Department of Physics and Center for Atmospheric and Space Sciences, Utah State University, Logan, Utah



Introduction

Light Detection and Ranging (lidar) is a ground-based remote sensing technique that has been used to study the middle and upper atmosphere for over four decades [1], [2]. Atmospheric lidar systems transmit laser beams into the atmosphere and then use optical and electronic detector systems to measure backscatter resulting from the interaction between the transmitted photons and atmospheric particles. Two of the most widely used lidar techniques for the study the upper atmosphere are Rayleigh-scatter

(RS) Lidar and sodium (Na) resonance lidar. RS lidar systems measure elastic backscatter from O_2 , N_2 and Ar particles in the atmosphere. RS lidar backscatter measurements give relative neutral density profiles, which are then used to calculate absolute temperature profiles. Na lidar measures resonant scatter from sodium atoms which form a layer in the 80-105 km region of the atmosphere where meteors typically ablate. From these measurements, Na density, temperature and zonal and meridional winds can be deduced.

This comparison of RS and Na lidar temperature data is the first study to show results from collocated two lidar systems. Previously, *Argall and Sica [2007]* presented a comparison of these two techniques from sites several hundred kilometers apart. Their results gave a temperature difference between the two types of lidar and called for a new comparison between collocated lidars. In this initial study, we will examine 19 nights of simultaneous measurements spread throughout one year.

1. Lidar Systems' Specifications

The original RS lidar system ran at a midlatitude site (42° N, 112° W), on the campus of Utah State University (USU), from 1993-2004 [4]. During this time, it gathered temperature data in the 45-90 km altitude range. It has since had an instrumentation upgrade (see Fig. 1) and has been used to collect temperature data from the 70-115 km range since summer 2014.

The Na lidar system ran on the campus of Colorado State University (41° N, 105° W) from 1990-2010 [5]. Since 2010, it has been operating at the same USU site as the RS lidar under the configuration [6] also shown in Figure 1.




Figure 1. Lidar system block diagrams for the current versions of the USU Na lidar [from 6] (left) and RS lidar (right).

2. Lidar Temperature Retrievals




Figure 2. Comparison plots of lidar temperatures versus altitude for 7 out of 19 nights. Orange curves give Na lidar temperatures and green curves give RS lidar temperatures.

The two lidars' temperature retrievals are done using completely different methods. RS lidar temperatures were calculated using a modified version of the Chanin-Hauchecorne method based on the backscattered power [1], [4]. Na lidar temperatures were derived using the method described in *Krueger et al., [2015]* which is based on the spectral shape of the returned signal.

3. Temperature Comparison

To better compare the two lidar datasets, the temperatures from each lidar, at a given altitude, were plotted in a time series in Figure 3. They show that at and below 90 km, the RS temperatures were generally colder than the Na temperatures. At 95 km and above, the RS temperatures are on average warmer than the Na temperatures.




Figure 3. RS and Na lidar temperature and temperature difference plots for five altitudes (85, 90, 95, 100 and 105 km) versus date (dates corresponding to date index given in Table 1).

Index	Date YYYYMMDD	Index	Date YYYYMMDD
0	140620	17	141029
1	140702	18	141104
2	140717	19	141106
3	140722	20	141108
4	140723	21	141109
5	140724	36	150328
10	140912	37	150414
11	140913	41	150610
12	140925	42	150618
13	140926		

In Figure 3 (a) and (b) for 85 and 90 km the seasonal variation of the mesopause is observed. There appears to be no seasonal dependence in the temperature differences between the two datasets.




Figure 4. Hourly temperature perturbations for 25 Sept 2014 as measured by the RS lidar (top) and Na lidar (bottom).

Hourly temperature perturbations were calculated from the both lidars' temperature measurements for 25 Sept 2015. To calculate the perturbations, for each lidar, an all-night average was subtracted from the hourly averages. The lidars' simultaneous observations lasted 10 hours, from about 3:30 AM to 12:30 PM UT.

There is good agreement between the two lidars' hourly temperature perturbations. Both lidars capture what appears to be about an 8-hour period wave with an amplitude of roughly 20 K and a downward phase velocity of 2 m/s.

4. Discussion

In Figure 2, we see that over the full measurement year (June 2014-June 2015), the best agreement between the two techniques happens between about 83-90 km. This can also be seen in temperature differences shown in Figure 3 (a) & (b) and in Figure 5 (top) where the correlation between the two datasets is greater than 0.9 in this altitude range.

The best agreement across the full range of altitudes is seen on the nights of the fall and spring equinoxes (Fig. 2 (b) & (e)). A night close to the fall equinox (25 Sept 2014) was chosen to examine the hourly temperature perturbations from both lidars. The perturbation plots (Fig. 4) show good agreement from hour-to-hour and the same 8-hr wave can be seen in both datasets.

The RS lidar temperature are shown to be colder than those of the Na lidar at 85 and 90 km (Fig. 3 (a) & (b)). A similar observation was made in *Argall and Sica, [2007]*. They compared RS and Na lidar climatologies from several different sites over an altitude range of about 80-95 km and found and that on average, the RS temperatures

were 7 K cooler. While our data show the RS temperatures being colder, our difference is not as strong—having an average of only about 2 K. At 95 km and above, our data shows that the RS temperatures are on average increasingly warmer as one goes up in altitude, reaching an average maximum temperature difference of about 16 K at 105 km (Fig. 3 (c)-(e)).

Conclusions and Future Work

We have presented a comparison of simultaneous temperatures acquired by Rayleigh-scatter and sodium resonance lidars collocated at USU. Several conclusions can be reached through this work:

- The two temperature datasets show the best agreement between about 83 and 90 km.
- The best agreement, spanning all altitudes, is seen near the fall and spring equinoxes (Fig. 2 (b) & (d)).
- Below 90 km, RS lidar temperatures are on average slightly cooler than Na lidar temperatures. At 95 km and above the RS temperatures are significantly warmer than the Na temperatures.
- On an hourly scale, temperature perturbations calculated independently for each lidar's dataset, show good agreement between the two techniques.
- The two sets of temperatures show better correlation as a function of altitude, than as a function of date.
- Occasionally, in summer months, the RS lidar observed a lower-in-altitude mesopause, which the Na lidar did not capture (Fig. 2 (g)).

These comparisons need to be continued with additional simultaneous observations. The apparent warmer RS temperatures above 95 km needs to be further investigated.

Acknowledgments

L. Sox and V. B. Wickwar would like to thank engineers: Thomas Amely, Matthew Emerick and Ryan Martineau. They would also like to thank the dedicated student observers and workers: David Barton Jordan Burns, Preston Hooser, David Moser, Luis Navarro, Rebecca Petrick, Patrick Sharp, Joe Slansky, Warren Schweigert and Bryant Ward. Special thanks for financial support goes to the Utah NASA Space Grant, USU, USU Physics Dept and personal contributions. T. Yuan and the Na lidar were supported under NSF AGS grant number 1135882.

References

[1] Hauchecorne, A. and M. L. Chanin (1980), Density and temperature profiles obtained by lidar between 35 and 70 km, *Geophys. Res. Lett.*, 7, 565-568.
 [2] Fricke, K. H. and U. von Zahn (1985), Mesopause temperatures derived from probing the hyperfine structure of the D_2 resonance line of sodium by lidar, *J. Atmos. Terr. Phys.*, 47, 499-512, doi:10.1016/0021-9169(85)90116-3.
 [3] Argall, P. S. and R. J. Sica (2007), A comparison of Rayleigh and sodium lidar temperature climatologies, *Ann. Geophys.*, 25, 27-35, doi:10.5194/angeo-25-27-2007.
 [4] Herron J. P. (2007), Rayleigh-scatter lidar observations at USU's atmospheric lidar observatory (Logan, UT) – Temperature climatology, temperature comparisons with MSIS, and noctilucent clouds. Doctoral Dissertation. Utah State University Library.
 [5] She, C. Y., S. Chen, Z. Hu, J. Sherman, J. D. Vance, V. Vasoli, M. A. White, J. Yu, D. A. Krueger (2000), Eight-year climatology of nocturnal temperature and sodium density in the mesopause region (80 to 105 km) over Fort Collins, Co (41°N, 105°W), *Geophys. Res. Lett.*, 27, 3289-3292, doi:10.1029/2000GL003825.
 [6] Krueger, D. A., C. Y. She, T. Yuan (2015), Retrieving mesopause temperature and line-of-sight wind from full-diurnal-cycle Na lidar observations, *App. Optics*, 54, doi: 10.1364/AO.54.009469