

Utah State University

DigitalCommons@USU

Dwarf Crops

Research

2002

Temperature Sensitivity of Four Tomato Cultivars

Jonathan M. Frantz

Justin Robinson

Bruce Bugbee

Utah State University, bruce.bugbee@usu.edu

Follow this and additional works at: https://digitalcommons.usu.edu/cpl_dwarfcrops



Part of the [Plant Sciences Commons](#)


Recommended Citation

Frantz, Jonathan M.; Robinson, Justin; and Bugbee, Bruce, "Temperature Sensitivity of Four Tomato Cultivars" (2002). *Dwarf Crops*. Paper 4.

https://digitalcommons.usu.edu/cpl_dwarfcrops/4

This Dataset is brought to you for free and open access by the Research at DigitalCommons@USU. It has been accepted for inclusion in Dwarf Crops by an authorized administrator of DigitalCommons@USU. For more information, please contact digitalcommons@usu.edu.





TEMPERATURE SENSITIVITY OF FOUR TOMATO CULTIVARS

Jonathan M. Frantz: slvyt@cc.usu.edu
 Justin Robinson: justrob@cc.usu.edu
 Bruce Bugbee: bugbee@cc.usu.edu



We evaluated 4 tomato cultivars, selected based on height (Red Robin and Micro-Tom) and familiarity (Pixie and Reimann Philipp) for their sensitivity to temperature. Each cultivar was grown in four 12-cm pots from seeding to form canopies in either 30/25 or 25/20 °C day/night temperature.

Yield of all cultivars except Reimann Philipp decreased with warmer temperatures with Red Robin and Pixie showing the most sensitivity. Fresh yield of the two dwarf varieties were lower than the taller cultivars, but due to higher percent dry matter in the fruit of the Micro-Tom and Red

Robin (see Cultivar Screening summary), the dry yield of the dwarf varieties exceeded that of Reimann Philipp. Yield rate was highest in Pixie and similar in the other cultivars.

Harvest index was highest in Micro-Tom. Micro-Tom produced many small fruit while Pixie produced few large fruit. Fruit of Pixie were smaller in warm temperatures, but Reimann Philipp were larger in warm temperatures. Reimann Philipp stands 40-cm tall at maturity, but collapses as the fruit enlarge making harvest difficult.

