

Utah State University

DigitalCommons@USU

Problems

Intermediate Modern Physics

1-8-2018

Physics 3710 – Problem Set #10

David Peak

Utah State University, david.peak@usu.edu

Follow this and additional works at: https://digitalcommons.usu.edu/intermediate_modernphysics_problems



Part of the [Physics Commons](#)

Recommended Citation

Peak, David, "Physics 3710 – Problem Set #10" (2018). *Problems*. Paper 10.
https://digitalcommons.usu.edu/intermediate_modernphysics_problems/10

This Course is brought to you for free and open access by the Intermediate Modern Physics at DigitalCommons@USU. It has been accepted for inclusion in Problems by an authorized administrator of DigitalCommons@USU. For more information, please contact digitalcommons@usu.edu.



Problems 1-3 refer to: The maximum measured z value for a galaxy is 11.1. As on page 1,

$$\text{GR7, } z = \frac{\lambda_d - \lambda_e}{\lambda_e}.$$

1. Suppose z is due to relative motion. Use the Doppler formula $\lambda_d = \left(\frac{1+\beta}{1-\beta}\right)^{1/2} \lambda_e$ to find the speed, β , with which this galaxy is receding from Earth.
2. Given that the universe scale factor is now 1, find a for this galaxy. (Equation (2), GR7, p4.)
3. What would the CMB temperature have been for this galaxy when it emitted the light we detect now? (GR8, p2)
4. Assume the universe contained only matter and vacuum-energy but with $\Omega_m = 0.27$ and $\Omega_v = 0.73$. Calculate the “age” of the universe in years by integrating $\int_0^a \frac{da'}{a' \sqrt{\frac{\Omega_m}{a'^3} + \Omega_v}} = tH_0$. Use the best-measured value of H_0 in the table on Gr9, p6.

(Note: according to Mathematica

$$\int \frac{dx}{\sqrt{(1-A)/x + Ax^2}} = \frac{2}{3\sqrt{A}} \ln \left[2 \frac{\sqrt{Ax^3 + (1-A)} + \sqrt{Ax^3}}{\sqrt{1-A}} \right].)$$