

Utah State University

DigitalCommons@USU

---

Techniques and Instruments

Crop Physiology Lab

---

5-2020

## Accuracy of the Generic Equation to Convert CCI to Chlorophyll Concentration in the Apogee Model MC-100 Chlorophyll Concentration Meter

Shuyang Zhen

Utah State University, shuyang.zhen@usu.edu

Paul Kusuma

Utah State University, paul.kusuma@aggiemail.usu.edu

Follow this and additional works at: [https://digitalcommons.usu.edu/cpl\\_techniquesinstruments](https://digitalcommons.usu.edu/cpl_techniquesinstruments)



Part of the [Physical Sciences and Mathematics Commons](#)

---

### Recommended Citation

Zhen, Shuyang and Kusuma, Paul, "Accuracy of the Generic Equation to Convert CCI to Chlorophyll Concentration in the Apogee Model MC-100 Chlorophyll Concentration Meter" (2020). *Techniques and Instruments*. Paper 17.

[https://digitalcommons.usu.edu/cpl\\_techniquesinstruments/17](https://digitalcommons.usu.edu/cpl_techniquesinstruments/17)

This Report is brought to you for free and open access by the Crop Physiology Lab at DigitalCommons@USU. It has been accepted for inclusion in Techniques and Instruments by an authorized administrator of DigitalCommons@USU. For more information, please contact [digitalcommons@usu.edu](mailto:digitalcommons@usu.edu).



# Accuracy of the generic equation to convert CCI to chlorophyll concentration in the Apogee model MC-100 Chlorophyll Concentration meter

Shuyang Zhen and Paul Kusuma  
Crop Physiology laboratory  
Utah State University

Chlorophyll strongly absorbs photons below 700 nm and transmits above 700 nm. Because of this property, chlorophyll concentration can be estimated by measuring the ratio of transmitted photons through a leaf at specific wavelengths. Chlorophyll concentration is then calculated using empirical relationships.

The fundamental relationship between transmitted photons and chlorophyll is highly non-linear. The Apogee model MC-100 linearizes the relationship using species-specific relationships that correct for differences in position and orientation of chloroplasts among species. This meter also includes a generic equation based on the average of 22 species (Parry et al., 2014), but accuracy of this generic empirical relationship has not been tested for species not in the original study.

## **Materials and Methods**


Measurements from seven plants (cucumber, radish, spinach, and four cultivars of lettuce) were used to evaluate the accuracy of the generic equation for chlorophyll concentration. Chlorophyll was extracted and measured using the procedures described by Parry et al. (2014). Values for chlorophyll concentration ranged from 50 to 700  $\mu\text{mol per m}^2$ .

## **Results**

The generic equation linearized the optical /absolute relationship for all species and values were close to the 1:1 line (Figure 1).

## **Discussion**

When a species-specific relationship is not available, the generic relationship linearized the optical/absolute relationship across species, and significantly improved the estimation of absolute chlorophyll compared to the use of an uncorrected index of transmitted photons.



**Figure 1.** Relationship between the measured and extracted chlorophyll concentration for cucumber (green), radish (red), spinach (purple), and four varieties of lettuce (blue).

**Literature Cited**

Parry, C., Blonquist, J. M., & Bugbee, B. (2014). *In Situ* Measurement of Leaf Chlorophyll Concentration: Analysis of the Optical/Absolute Relationship. *Plant, Cell, and Environment*, 37, 2508 - 2520.