

1-1-1917

X. Notes on some British Guiana Hymenoptera (exclusive of the Formicidae).

G. E. Bodkin

Department of Science and Agriculture, British Guiana

Follow this and additional works at: https://digitalcommons.usu.edu/bee_lab_be



Part of the [Entomology Commons](#)

Recommended Citation

Bodkin, G. E., "X. Notes on some British Guiana Hymenoptera (exclusive of the Formicidae)." (1917). *Be*. Paper 126.

https://digitalcommons.usu.edu/bee_lab_be/126

This Article is brought to you for free and open access by the Bee Lab at DigitalCommons@USU. It has been accepted for inclusion in Be by an authorized administrator of DigitalCommons@USU. For more information, please contact digitalcommons@usu.edu.



Revision of the genus *Tarucus*.

s thespis, L.

Mus. Ind. Ner., p. 318, N. 136
p. 791, N. 236 (1767).

se to *T. bowkeri*, but differs in
It is of a decidedly brighter
pperside, and has quite narrow
rs, the fringes are longer and
; it has not in the secondaries
as obtains in *bowkeri*. On the
t colour is brown, not whitish
The markings are similar, but
rminal row of spots, well marked
thespis. In the primaries the fold
entirely brown, except for quite
e tornus; the tail is little more
ge.

y brown above in both wings
in *bowkeri*, but with a slight
ision, which, in the secondaries,
ginal area; a small white spot
closing the cell and has three
oots beyond it.
y, where it is widely distributed
ern districts, and has also been

ne type as *bowkeri*, the clasp being
rentiated. The tegumen is distinctly
with a very narrow apex, and very
de lobes, whilst the falces are longer
edoeagus originates of moderate size,
the zone, where it is suddenly excised
a point, this front portion being half
t. The vesica is a beautiful object,
mpet-shaped, and being very finely
rgae excitatae are absent, but the
dified form, somewhat as in *bowkeri*.

OF PLATES XIV-XX.

on facing the PLATES.]

X. Notes on some British Guiana Hymenoptera (exclusive
of the Formicidae). By G. E. BODKIN, B.A., Dip.
Agric. (Cantab.), F.Z.S., F.E.S., Government Eco-
nomic Biologist, Department of Science and Agri-
culture, British Guiana.

(Published by permission of the Director of Science and
Agriculture, British Guiana.)

[Read December 6th, 1916.]

PLATES XXI-XXIII AND SKETCH MAP.

UP to the present time very little has been known of the
habits and life-histories of the *Hymenoptera* of British
Guiana. Schomburgk,* in his "Fauna and Flora of British
Guiana," devotes six pages to the *Hymenoptera* and gives
a few observations with regard to their biology as observed
by himself, but unfortunately many of the scientific
names are quite impossible to trace. Peter Cameron has
published in "Timehri"† a comprehensive list of the
Hymenoptera of this country, with descriptions of many
new species, but no biological notes are attached. Scat-
tered references have appeared from time to time in many
scientific publications, but the majority of these are
descriptions of new species.

The present collection in this laboratory was commenced
in 1911 and has been formed chiefly by myself as oppor-
tunities have occurred. All the commoner species have
now been collected, and in many cases observations made
on their life-histories and habits. The accompanying map
indicates in red dots the areas where collections and
observations have been made. These necessarily follow

* Schomburgk, R., "Fauna und Flora von British Guiana."
Leipzig, 1848.

† Peter Cameron, Hymenoptera of the Georgetown Museum,
"Timehri." Journal of the Royal Agricultural and Commercial
Society, 1911-12. Pt. 1. Parasitica, I, pp. 153-186 (1911). Pt. 2.
Parasitica, I, pp. 306-330 (1911). Pt. 3. Marabuntas or Wasps,
II, pp. 207-231 (1912). Pt. 4. Fossores, II, pp. 412-440 (1912).

TRANS. ENT. SOC. LOND. 1917.—PARTS II, III, IV (MAY '18)

the chief communication routes either by water, road, or trail. Vast areas consequently remain untouched.

The collection at present contains 1,600 specimens, of which 161 species are named. All these identifications have been made either by specialists in the British Museum, through the co-operation of the Imperial Bureau of Entomology, or by specialists in the U.S. National Museum.

The present work clearly shows that in scope it can hardly pretend to do more than outline the vast field which awaits entomologists in this part of the world.

There is a distinct difference between the *Hymenoptera* of the flat, cultivated and inhabited coast lands and those of the forest-clad area. Many species of common occurrence on the coast lands are never found in the forests, and *vice versa*. The climate of these two areas also varies, the interior districts having a higher rainfall and a somewhat higher temperature. The trade-winds which sweep the coast lands most of the year are not experienced to any extent inland. No opportunity has yet occurred to investigate the large tracts of savannah lands which exist at the back of the Colony near the Brazilian frontier.

The observations on the habits and life-histories are mostly my own, but a number of interesting notes by the following gentlemen must be acknowledged: Mr. C. B. Williams, Mr. L. D. Cleare, Jnr., Mr. H. W. B. Moore, and Mr. A. A. Abraham.

I have endeavoured to give as full information as possible concerning each species. Where only one specimen of a species has been collected I have given the locality of collection.

The Ants have not been included in these notes, as Mr. W. C. Crawley has already published* an account of the Family.

Family APIDAE.

Subfamily SPHECODINAE.

Genus TEMNOSOMA, Smith.

T. aeruginosum, Smith. Issororo, N.W.D.

* "Ants from British Guiana," W. C. Crawley, B.A. *Annals and Magazine of Natural History*, Ser. 8, vol. xvii, May, 1916.

some British Guiana Hy

Subfamily ANDRENI

Genus AUGOCHLORA,

A. graminea, F. } Taken whil
A. thalia, Smith. } of "wouralli
A. calypso, Sm. } Issororo, N.W.

Subfamily PANURGI

Genus MEGALOPTA,

M. sodalis, Vachal. Issororo, N.
frequently attracted to artificial ligh

Subfamily XYLOCOP

Genus XYLOCOPA

X. fimbriata, F. The commonest British Guiana. The female is bl ochreous yellow with green eyes females is greater than males—abo may frequently be seen collecting of commonly occurring flowers, e large red *Hibiscus* and the flower It possesses a powerful sting, and th by a loud buzzing sound. I hav species in the interior. Any dry de for nesting purposes, such as rotte stumps. The softer kinds of wood wood infested by these bees soon l with their borings, and large quan seen piled up at the foot of the log. tapped a shrill buzzing noise may contained bees. A log of wood 5 diameter on being split open wa imagos and 25-30 larvae and pup three entrance holes, and these le bored at right angles to the grain galleries the cells are formed, usuall never more. Each cell is about about $\frac{3}{4}$ inch in diameter. The c one another by a partition or "wad together and hardened by the bee $\frac{1}{4}$ inch in thickness. The galleries smooth and very neatly executed with a small, fairly solid mass co mixture of honey and pollen. I

Subfamily ANDRENINAE.

Genus AUGOCHLORA, Smith.

A. graminea, F. } Taken while feeding on flowers
A. thalia, Smith. } of "wouralli" (fish-poison plant),
A. calypso, Sm. } Issororo, N.W.D.

Subfamily PANURGINAE.

Genus MEGALOPTA, Smith.

M. sodalis, Vachal. Issororo, N.W.D. This species is frequently attracted to artificial light.

Subfamily XYLOCOPINAE.

Genus XYLOCOPA, L.

X. fimbriata, F. The commonest wood-boring bee in British Guiana. The female is black and the male an ochreous yellow with green eyes. The proportion of females is greater than males—about 4 to 1. Both sexes may frequently be seen collecting honey from a number of commonly occurring flowers, especially those of the large red *Hibiscus* and the flowers of the Pigeon Pea. It possesses a powerful sting, and the flight is accompanied by a loud buzzing sound. I have never observed the species in the interior. Any dry decaying wood is utilised for nesting purposes, such as rotten paling-posts or tree-stumps. The softer kinds of wood are preferred. Logs of wood infested by these bees soon become literally riddled with their borings, and large quantities of frass may be seen piled up at the foot of the log. If such a log is sharply tapped a shrill buzzing noise may be heard caused by the contained bees. A log of wood 5 feet long by 2 feet in diameter on being split open was found to contain 20 imagos and 25-30 larvae and pupae. There were about three entrance holes, and these led directly into galleries bored at right angles to the grain of the wood. In such galleries the cells are formed, usually three or four together, never more. Each cell is about an inch in length and about $\frac{3}{4}$ inch in diameter. The cells are separated from one another by a partition or "wad" of sawdust cemented together and hardened by the bee. These partitions are $\frac{1}{4}$ inch in thickness. The galleries and cells are perfectly smooth and very neatly executed. Each cell is stocked with a small, fairly solid mass consisting apparently of a mixture of honey and pollen. It is a dark yellow in

routes either by water, road, or frequently remain untouched. Present contains 1,600 specimens, of named. All these identifications by specialists in the British operation of the Imperial Bureau specialists in the U.S. National

clearly shows that in scope it can more than outline the vast field which this part of the world.

difference between the *Hymenoptera* and inhabited coast lands and those

Many species of common occurrences are never found in the forests, climate of these two areas also varies, having a higher rainfall and a some- more. The trade-winds which sweep of the year are not experienced to No opportunity has yet occurred ge tracts of savannah lands which of the Colony near the Brazilian

in the habits and life-histories are number of interesting notes by the must be acknowledged: Mr. C. B. reare, Jnr., Mr. H. W. B. Moore, and

to give as full information as possible species. Where only one specimen collected I have given the locality of

been included in these notes, as already published * an account of

family APIDAE.

family SPHECODINAE.

TEMNOSOMA, Smith.

ith. Issororo, N.W.D.

Guiana," W. C. Crawley, B.A. Annals History, Ser. 8, vol. xvii, May, 1916.

colour with a peculiar though not unpleasant odour. These masses of foodstuff weigh about 2.5 grammes, and an equal quantity is deposited in each cell. On this mass the egg is finally deposited and gradually increases in size as development proceeds; later, the segmentation of the future larva may dimly be perceived through the enveloping shell. It is sausage-shaped, slightly curved, and almost transparent in the earlier stages. Length 1.7 cm., diameter 0.4 cm. During development one end becomes somewhat larger than the other. Eventually the extremely thin pellicule strips off and the wrinkled larva is exposed, which starts feeding immediately. The larval excrement is hard, and formed in short rods black in colour. The full-grown larva measures 3.9 cm. in length, and is creamy white in colour. The larval stage lasts about three weeks. Gradually the outlines of the pupa may be seen through the larval skin, which eventually peels off. The period between the full-fed larva and the final stripping of the larval skin is 48 hours. The pupa is at first creamy white, and in form roughly resembles the future adult insect. Length of pupa 2.5 cm., breadth 1.4 cm. Within the first week of pupation a general darkening of colour takes place, the eyes going almost black; hardening of the integument occurs simultaneously. These two processes progress rapidly till the perfect insect is formed in about 3 weeks. It then makes its exit from the cell by gnawing away the wad of hardened sawdust.

X. barbata, F. A fairly common species on the coast lands.

X. brasilianorum, F. A not uncommon species within the forest area. Rockstone, Essequibo River, and H.M. Penal Settlement, Mazaruni River.

X. aurulenta, F. An uncommon species within the forest area. Rockstone, Essequibo River.

Subfamily PROSOPIDINAE.

Genus CAUPOLICANA, Spinola.

C. eximia, Smith. Essequibo River, in vicinity of Rockstone.

Subfamily ANTHOPHORINAE.

Genus EUCERA, Scopoli.

E. festiva, Sm. Vicinity of Georgetown.

some British Guiana

Genus EXOMALOPUS

E. globosa, F. Botanic Garden

Genus EPICHAERUS

E. rustica, Oliv. Rockstone,

Genus MELITOPUS

M. englossoides, Lep. From coast lands and in the interior. This bee has a cream-coloured body, the edges of blades of Para grass embedded in the tissues. They remain quite apparent. They remain quite common about coast lands.

Genus CENTRUS

C. longimana, Lep. A common species on the coast lands and in the interior. It is attracted to smelling substances such as salt, and is often found with this substance on board ships. Frequently seen about the edges of the tidal passages in the big rivers. It is apparently attracted there by sea-birds on these objects. It is common about the interior.

C. lanipes, F. Fairly common in the interior. Has been taken in the act of collecting soft mud from the interior.

C. versicolor, F. Taken on coast lands to artificial light. Appears to be attracted to artificial light.

C. personata, Sm. It has similar habits to *C. personata*, Sm. Taken under similar conditions in the District.

C. labrosa, Friese. An uncommon species in the District. Settlement, Mazaruni River.

Genus THYGATIS

T. rubricata, Sm. Rockstone,

Genus EUGLOSSA

E. dimidiata, F. A fairly common species in the parts of the Colony. Observed collecting mud from a pathway. TRANS. ENT. SOC. LOND. 1918.

though not unpleasant odour. weigh about 2.5 grammes, and are packed in each cell. On this mass the larva gradually increases in size and later, the segmentation of the larva is perceived through the enveloped, slightly curved, and almost cylindrical stages. Length 1.7 cm., diameter 0.5 cm. At one end becomes somewhat flattened. Eventually the extremely thin larva and the wrinkled larva is exposed immediately. The larval excrement consists of rods black in colour. The larva is 3.9 cm. in length, and is creamy white. The larval stage lasts about three weeks. The pupa may be seen through the transparently peels off. The period of the pupa and the final stripping of the pupa is at first creamy white, and resembles the future adult insect. Length 1.4 cm. Within the first few days a darkening of colour takes place, most black; hardening of the cuticle occurs. These two processes are completed. The perfect insect is formed in about 10 days. It exits from the cell by gnawing through the sawdust.

A very common species on the coast

A not uncommon species within the colony, Essequibo River, and H.M. Mazaruni River.

A common species within the forest of Essequibo River.

PROSOPIDINAE.

PROSOPIDINA, Spinola.

Essequibo River, in vicinity of

ANTHOPHORINAE.

ANTHOPHORA, Scopoli.

of Georgetown.

some British Guiana Hymenoptera.

301

Genus EXOMALOPSIS, Spinola.

E. globosa, F. Botanic Gardens, Georgetown.

Genus EPICHARIS, Klug.

E. rustica, Oliv. Rockstone, Essequibo River.

Genus MELITOMA, Latr.

M. euglossoides, Lep. From cultivated Cotton blossoms, Georgetown. This bee has a curious habit of clinging to the edges of blades of Para grass, with its mandibles firmly embedded in the tissues. The reason for this is not apparent. They remain quite motionless in this position, and at times may be seen in considerable numbers. Fairly common about coast lands.

Genus CENTRIS, F.

C. longimana, Lep. A common species both on the coast lands and in the interior. It is attracted to strong-smelling substances such as salt fish, and will follow boats with this substance on board for long distances up the rivers. Frequently seen about buoys moored in the centre of the tidal passages in the big rivers and elsewhere. It is apparently attracted there by the excreta dropped by sea-birds on these objects. It has a swift, noisy flight.

C. lanipes, F. Fairly common on the coast lands and in the interior. Has been taken at flowers and also in the act of collecting soft mud from a pathway.

C. versicolor, F. Taken on one occasion while attracted to artificial light. Appears to occur only in the interior.

C. personata, Sm. It has similar habits to *C. longimana*, Lep. Taken under similar conditions in the North-west District.

C. labrosa, Friese. An uncommon species. H.M. Penal Settlement, Mazaruni River.

Genus THYGATER, Holmb.

T. rubricata, Sm. Rockstone, Essequibo River.

Genus EUGLOSSA, Latr.

E. dimidiata, F. A fairly common species in most parts of the Colony. Observed on several occasions to collect mud from a pathway. A somewhat clumsy insect, TRANS. ENT. SOC. LOND. 1918.—PARTS II, III, IV. (MAY '18) X

easy to capture, but when alarmed assumes a rapid flight. Frequently observed flying about and alighting upon the bark of certain trees, though on closely inspecting the bark no feature which might attract the bee could be observed.

E. cordata, F. The commonest *Euglossa* in British Guiana. Observed in all areas visited. It will construct its somewhat sticky nest in all sorts of curious places, such as the inside of a disused reel of cotton, interior of empty cartridge case, eye-piece of polariscope, keyholes, small cavities in timber, etc. It also takes over the disused cells of *Sceliphron fistulare*, Dahlb. In the forest areas I have observed this bee to construct its nest on the under surface of a leaf. The nest varies in the number of its cells, some only containing two or three, others as many as six. The cells are roughly about 1 cm. in length and about 5 mm. in breadth, with the ends neatly rounded. Each cell is stored with a viscid mixture of honey and pollen, on which the larva feeds. The walls of the cells are thin and soft, being constructed of some dark vegetable substance. The whole exterior is sticky. This insect has an extremely rapid, darting flight, and is particularly fond of the white, sweet-smelling flowers of a certain species of prickly wild Solanaceous plant. While the bee is within the white blossoms it emits an exceedingly shrill buzzing sound, which is often difficult to locate.

E. surinamensis, L. Another very common species of *Euglossa* throughout British Guiana. It is greatly attracted to the flowers of the same Solanaceous plant as attracts *E. cordata*, L. Its loud buzzing and somewhat slow flight makes its presence conspicuous. The nest is constructed sometimes in artificial holes in timber, in cavities in the trunks of trees, and at times on the top of roof-beams. The substance of the nest is thin flakes of bark, or sometimes flakes of white plaster from houses, exceedingly strongly cemented together. Several cells are usually formed united in an irregular mass. Each cell is stored up with a quantity of honey and pollen of moderately hard consistency. The cells are about 1.5 cm. in length, with a diameter of about 1 cm., ovoid in shape, with a perfectly smooth lining.

E. nigrita, Lep. Not a particularly common species. So far only observed on the coast lands. On one occasion the bees were observed to be nesting within a hollow beam in the large dining-hall of one of the largest hotels in

Georgetown. The bees passed to and fro regardless of the proximity of human beings.

E. piliventris, Guér. An uncommon species. Observed on the Essequibo River. Attracted to flowery naceous plant previously mentioned.

E. analis, Lep. An uncommon species. Its nest, consisting of many ovoid cells was found in the soil on the East Coast of Demerara. The cells were dark in colour, and joined together. The bees were taken from this nest.

E. cayennensis, Lep. (= *fasciata*, Lep.) A very common and widely distributed species throughout the country. Nesting habits not observed.

E. decorata, Sm. An uncommon species. Observed on the Essequibo River.

Genus EXAERETE, H.

E. smaragdina, Guér. A common species. Observed of the interior, especially the N.W.D. The bees were collected while flying about plants from which a strong-smelling sap was exuding, in which the bees were feeding. Occasional specimens were taken on the lands and in the Botanic Gardens, but nesting habits unobserved.

Subfamily MEGACHILIDAE

Genus MEGACHILE, L.

M. lobitarsis, Smith. An uncommon species. Distribution not observed.

M. lanata, F. This insect has only been observed and that beneath the Government buildings in town. The mud cells were situated in a 3/4-inch iron pipe. The nest was cylindrical and contained about four cells. The cells were cemented together, making the whole nest very strongly and hard to break. It was found in India.*

Subfamily COELIOXYGIDAE

Genus COELIOXYGUS, L.

C. simillima, Smith. Apparently not a common species. Nesting habits not observed. The bees were found in a wooden post.

* Lefroy, H. Maxwell, "Indian Insects."

n alarmed assumes a rapid flight. flying about and alighting upon the rough on closely inspecting the bark attract the bee could be observed. commonest *Euglossa* in British Guiana areas visited. It will construct nests in all sorts of curious places, disused reel of cotton, interior of eye-piece of polariscope, keyholes, etc. It also takes over the *tron fistulare*, Dahlb. In the forest this bee to construct its nest on the f. The nest varies in the number containing two or three, others as many as are roughly about 1 cm. in length and breadth, with the ends neatly rounded. It is filled with a viscid mixture of honey and in which the larva feeds. The walls of the cells are constructed of some dark vegetable matter, the exterior is sticky. This insect has a rapid flight, and is particularly fond of visiting flowers of a certain species of Solanaceous plant. While the bee is within the nest it emits an exceedingly shrill buzzing sound, which is difficult to locate.

Another very common species of *Euglossa* in British Guiana. It is greatly attracted to the same Solanaceous plant as attracts the first species and has a somewhat slow flight and is not so conspicuous. The nest is constructed of mud in holes in timber, in cavities in the wall, or at times on the top of roof-beams. The nest is thin flakes of bark, or sometimes plaster from houses, exceedingly thin and fragile. Several cells are usually packed together. Each cell is stored with a mass of honey and pollen of moderately coarse texture. The cells are about 1.5 cm. in length, and about 1 cm., ovoid in shape, with a narrow entrance.

Not a particularly common species. It is found on the coast lands. On one occasion it was observed to be nesting within a hollow beam of the hall of one of the largest hotels in

Georgetown. The bees passed to and fro, apparently quite regardless of the proximity of human beings.

E. piliventris, Guér. An uncommon species from Upper Essequibo River. Attracted to flowers of the wild Solanaceous plant previously mentioned.

E. analis, Lep. An uncommon species. A large nest consisting of many ovoid cells was once taken from the soil on the East Coast of Demerara. The cells were hard, dark in colour, and joined together. Only one bee hatched from this nest.

E. cayennensis, Lep. (= *fasciata*, Lep.). A fairly common and widely distributed species throughout the Colony. Nesting habits not observed.

E. decorata, Sm. An uncommon species from the interior. Essequibo River.

Genus EXAERETE, Hoffm.

E. smaragdina, Guér. A common species in some parts of the interior, especially the N.W.D. All of my specimens were collected while flying about piles of cordwood from which a strong-smelling sap was exuding, and on which the bees were feeding. Occasionally seen on the coast lands and in the Botanic Gardens, Georgetown. Nesting habits unobserved.

Subfamily MEGACHILINAE.

Genus MEGACHILE, Latr.

M. lobitarsis, Smith. An uncommon species with a wide distribution.

M. lanata, F. This insect has only once been collected, and that beneath the Government Laboratory, Georgetown. The mud cells were situated within a disused 3/4-inch iron pipe. The nest was cylindrical in shape and contained about four cells. The nest was very firmly cemented together, making the whole structure exceedingly strong and hard to break. This bee is well known in India.*

Subfamily COELIOXYNAE.

Genus COELIOXYS, Latreille.

C. simillima, Smith. Apparently a widely distributed though not common species. Nest observed in cylindrical borings in a wooden post.

* Lefroy, H. Maxwell, "Indian Insect Life," p. 219.

Subfamily BOMBINAE.

Genus BOMBUS, Latr.

B. cayennensis, F. A common species found only in the interior.

Genus MELIPONA, Ill.

M. interrupta, Latr. This species is widely distributed and common. The specimens in the collection were all taken while they were collecting soft mud from paths.

M. pallida, Latr. A fairly common species in the interior. A small nest on one occasion observed in a rotten log of timber with a small circular hole formed of wax for exit. When the nest is disturbed they swarm out and attack by biting the exposed parts of the head and neck, emitting at the same time a shrill buzzing sound. These small bees possess a peculiar odour which is characteristic.

M. favosa, F. A common species on the coast lands, where it is known as the Courida Bee, from a supposed habit of collecting honey from the flowers of the maritime Courida (*Avicennia nitida*). The nests are found in hollow trees, etc., especially in old Courida trees, but they have been found in disused drain-pipes and other unlikely places. This bee possesses no sting, and I have never observed it to attack in any way; when the nest is disturbed the bees swarm out, but do not demonstrate their resentment in any more practical manner. This bee is frequently domesticated, and when the nests are found in the field they are removed, taken home, and put in wooden boxes with a small exit hole. They thrive in captivity. The honey which they produce is thin and of a somewhat insipid flavour; mixed with other ingredients it is utilised by the natives as a cure for cold in the throat or chest. The honey is stored in egg-shaped cells constructed of wax, about $1\frac{1}{2}$ -2 inches in length; similar cells are also constructed containing nothing but wax. The cells containing the larvae are 8 mm. in length and about 5 mm. in breadth.

M. clavipes, F. A common species in most parts of the Colony. The nests are usually constructed in hollow decaying logs. The following notes on the nesting habits of this species were made from a nest which was found in a hollow log of Trysil wood (*Pentaclethra filamentosa*). The extreme length of the hollowed-out portion containing

the nest was 32 inches. Three different kinds of cells were observed on the nest, arranged longitudinally. The layer next to the entrance consisted of wax cells about $\frac{1}{2}$ inch in length and $\frac{1}{4}$ inch in width, ovate in shape. The length of the cells increased towards the interior. These wax cells were yellowish in colour and of solid wax. To the taste the wax had a slightly sweet flavour, and a sour smell which was perceptible throughout the whole nest. The layer next to the honey cells; this layer was 9 inches in length and the cells themselves were slightly smaller than those of the entrance, in shape and dark brown in colour. The wax was thick, very sweet and possessed a pleasant odour, but if the slightest trace of the wax was got mixed with the honey the bitter taste of the honey obliterated the pleasant flavour of the wax. The layer consisted of the cells with embryos in different stages of development; it was 9 inches in length and the cells were about $\frac{1}{2}$ inch in length and $\frac{1}{4}$ inch in width. In shape they were cylindrical. The nest was entirely enclosed, with the exception of a small exit was by no means a conspicuous feature. The surface of the wax constructed of wax much the same as the surrounding bark. It was slightly raised above the surface of the bark, and roughly circular with a small exit hole. Attention was drawn to this nest by the bees hovering about it. This species is by no means so pugnacious as some species of *Melipona*.

M. recurva, Sm. An uncommon species. District.

M. lineata, Lep. Bartica, Essequibo. *M. flavipennis*, Sm. Taken on attending flowers of Guava.

M. varia, Lep. A nest of this species was found in a large greenheart beam supporting the Essequibo River. A non-pugnacious species to the nest is funnel-shaped and conical.

M. guianae, Ckll. A most pugnacious nest is only slightly disturbed the intruder by biting the head and neck, at the same time enjoining the intruder to leave. The nests (which are large) are usually found in the branches of trees at some distance

BOMBINAE.

OMBUS, Latr.

mon species found only in the

ELIPONA, Ill.

is species is widely distributed
mens in the collection were all
ecting soft mud from paths.
fairly common species in the
n one occasion observed in a
a small circular hole formed of
nest is disturbed they swarm
the exposed parts of the head
he same time a shrill buzzing
possess a peculiar odour which

mon species on the coast lands,
Courida Bee, from a supposed
from the flowers of the maritime
. The nests are found in hollow
ld Courida trees, but they have
drain-pipes and other unlikely
ses no sting, and I have never
any way; when the nest is dis-
ut, but do not demonstrate their
practical manner. This bee is
and when the nests are found in
d, taken home, and put in wooden
hole. They thrive in captivity.
roduce is thin and of a somewhat
with other ingredients it is utilised
e for cold in the throat or chest.
g-shaped cells constructed of wax,
length; similar cells are also con-
ning but wax. The cells contain-
n. in length and about 5 mm. in

ommon species in most parts of
are, usually constructed in hollow
lowing notes on the nesting habits.
le from a nest which was found in
l wood (*Pentaclethra filamentosa*).
he hollowed-out portion containing

the nest was 32 inches. Three distinct layers of various kinds of cells were observed on splitting open the log longitudinally. The layer next to the entrance consisted of wax cells about $\frac{1}{2}$ inch in length and $\frac{3}{4}$ inch in diameter; ovate in shape. The length of this layer was 8 inches. These wax cells were yellowish in colour and contained solid wax. To the taste the wax had an exceedingly bitter flavour, and a sour smell which seemed to pervade the whole nest. The layer next to the wax cells contained honey cells; this layer was 9 inches in length. The cells themselves were slightly smaller than the wax cells, same shape and dark brown in colour. The contained honey was thick, very sweet and possessed quite a good flavour, but if the slightest trace of the wax coating of the cell got mixed with the honey the bitterness of the wax entirely obliterated the pleasant flavour of the honey. The next layer consisted of the cells with embryonic bees in various stages of development; it was 9 inches in length. These cells were about $\frac{1}{2}$ inch in length and dull yellow in colour. In shape they were cylindrical. Apparently the nest was entirely enclosed, with the exception of the exit. This exit was by no means a conspicuous object, as it was constructed of wax much the same colour and texture as the surrounding bark. It was slightly raised above the surface of the bark, and roughly cone-shaped with a very small exit hole. Attention was drawn to the presence of this nest by the bees hovering around the exit. This species is by no means so pugnacious as some of the other species of *Melipona*.

M. recurva, Sm. An uncommon species from the N.W. District.

M. lineata, Lep. Bartica, Essequibo River.

M. flavipennis, Sm. Taken on one occasion while attending flowers of Guava.

M. varia, Lep. A nest of this species observed in a large greenheart beam supporting the hotel at Rockstone, Essequibo River. A non-pugnacious species. The entrance to the nest is funnel-shaped and constructed of wax.

M. guianae, Ckll. A most pugnacious species. If the nest is only slightly disturbed the bees swarm out and attack the intruder by biting the exposed parts of the head and neck, at the same time emitting a shrill buzzing. The nests (which are large) are usually constructed in the branches of trees at some distance from the ground. The

bees possess a peculiar smell, due to the character of the substance which they collect on their hind-legs. A large nest observed on one occasion on the branches of a Pimento tree (*Pimenta officinalis*).

M. rufiventris, Lep., var. *flavolineata*, Friese. Tumatumari, Essequibo River.

Melipona amalthea, F. A common and widely distributed species. May be observed on the blossoms of most garden plants, and is particularly fond of feeding on over-ripe fruit. It may also frequently be seen collecting mud from damp paths, creeks, etc. Known locally as "Tarbaby bees." It has a habit of collecting the scrap-rubber from recently tapped trees of *Hevea brasiliensis*.

M. rufiventris, Lep. Rockstone, Essequibo River.

M. dallatorreana, Friese. East Coast, Demerara.

M. mutata, Lep. Upper Demerara River.

Genus *Apis*, Linn.

A. mellifera, L. Only a small number of hives of the domestic bee are kept in the Colony, and these are principally owned by the Chinese and Portuguese. The honey produced is of good quality and very sweet. Fresh stock is usually imported from the United States. They are mostly "Italian bees." The Wax Moth (*Galleria mellonella*, L.) is common and causes much damage.

Family VESPIDAE.

Subfamily VESPINAE.

Genus *Polistes*, Latr.

P. pacificus, F. Essequibo Coast. An uncommon species.

P. analis, F. A fairly common species in the interior.

P. versicolor, Oliv. A common species on the coast lands.

P. goeldii, Ducke. A rare species in the interior.

P. canadensis, L., var. *amazonicus*, Schulz. The commonest species of *Polistes* in the Colony, occurring everywhere. Unless severely molested it rarely attacks people, but its sting is both powerful and painful, and causes a severe swelling. Large nests are rarely seen, the usual number of cells being about twenty. Beneath bridges, houses, on the rafters, under the eaves, beneath the platform of railway stations, behind pictures, etc., are favourite

nesting-places for this species. domestic habitations for its nests. wooden bridges which span the sugar estates are always thickly with nests of these insects, and passing in a small boat is always an exciting a painful experience. In dwelling a source of danger, especially v. The local name is "marabunta." nests with a wad of dried palm pole soaked in kerosene and in operation for an onlooker at a times these wasps will remain qu position on their nests for hours guard. From the economic stand ingly useful, for they may often consuming the larvae of various cially the Rice Worm (*Laphygma*). The nests are irregular in structure for pieces are frequently falling f of this insect is somewhat clumsy

P. crinitus, Felton. An uncommon lands.

Genus *Polybia*

P. fulvofasciata, de G. (= *p*) species on the coast lands; the attached to the under surfaces of the mango tree. Large nests

P. occidentalis, Oliv. An ex throughout the coast lands, an the interior. The nests are fo surfaces of many species of pa large, more or less circular in exceedingly light and fragile is not unduly pugnacious, and nest is damaged; it is thus fr felling the branches of young has no great lasting effects, bu ingly painful. On one occas trimming a hedge of Barbado brought into the laboratory species which he had very ca of the wasps *in situ* on the ou

nect, due to the character of the
flect on their hind-legs. A large
sion on the branches of a Pimento

ar. *flavolineata*, Friese. Tumatu-

A common and widely distributed
ed on the blossoms of most garden
rly fond of feeding on over-ripe
requently be seen collecting mud
, etc. Known locally as "Tarbaby
of collecting the scrap-rubber from
Hevea brasiliensis.

Rockstone, Essequibo River.

ese. East Coast, Demerara.

pper Demerara River.

us APIS, Linn.

ly a small number of hives of the
in the Colony, and these are prin-
hinese and Portuguese. The honey
ality and very sweet. Fresh stock
from the United States. They are

The Wax Moth (*Galleria mellonella*,
uses much damage.

ily VESPIDAE.

amily VESPINAE.

us POLISTES, Latr.

Essequibo Coast. An uncommon

ly common species in the interior.

A common species on the coast

A rare species in the interior.

var. *amazonicus*, Schulz. The com-
istes in the Colony, occurring every-
ly molested it rarely attacks people,
powerful and painful, and causes a
ge nests are rarely seen, the usual
g about twenty. Beneath bridges,
, under the eaves, beneath the plat-
ns, behind pictures, etc., are favourite

nesting-places for this species. It appears to prefer
domestic habitations for its nesting-places. The short
wooden bridges which span the navigation trenches on
sugar estates are always thickly infested beneath with the
nests of these insects, and passing beneath such bridges
in a small boat is always an exciting and quite occasionally
a painful experience. In dwelling-houses they are always
a source of danger, especially when children are about.
The local name is "marabunta." Destroying marabunta
nests with a wad of dried palm leaves attached to a long
pole soaked in kerosene and ignited is an interesting
operation for an onlooker at a respectful distance. At
times these wasps will remain quite motionless in an alert
position on their nests for hours together, as though on
guard. From the economic standpoint they are exceed-
ingly useful, for they may often be seen hunting for and
consuming the larvae of various agricultural pests, espe-
cially the Rice Worm (*Laphygma frugiperda*, S. & A.).
The nests are irregular in structure and not strongly made,
for pieces are frequently falling from the nest. The flight
of this insect is somewhat clumsy.

P. crinitus, Felton. An uncommon species on the coast
lands.

Genus POLYBIA, Lep.

P. fulvofasciata, de G. (= *phthisica*, F.). A common
species on the coast lands; the nests are frequently found
attached to the under surfaces of leaves, especially those
of the mango tree. Large nests are seldom encountered.

P. occidentalis, Oliv. An exceedingly common species
throughout the coast lands, and at times encountered in
the interior. The nests are found attached to the under
surfaces of many species of palms. The nests are never
large, more or less circular in shape, and constructed of
exceedingly light and fragile material. This small wasp
is not unduly pugnacious, and seldom attacks unless the
nest is damaged; it is thus frequently encountered when
felling the branches of young coconut palms. The sting
has no great lasting effects, but the first shock is exceed-
ingly painful. On one occasion a gardener who was
trimming a hedge of Barbados Cherry (*Malpighia glabra*)
brought into the laboratory a good-sized nest of this
species which he had very carefully removed with some
of the wasps *in situ* on the outside of the nest. The local

name is "honey marabunta." A native method of destroying such nests is to seize the nest quickly and firmly with both hands and then crush it. An individual possessed of large and thick-skinned hands is likely to be the most successful.

P. fastidiosuscula, Sauss., var. *sampaioi*, Ducke. Appears to take the place of the foregoing in the interior, where it is common. It has never been taken on the coast lands. The nest is often met with attached to the under surfaces of palm leaves.

P. fasciata, Lep. A species by no means of infrequent occurrence both on the coast lands and interior. A small nest was taken on one occasion attached to the floor-boards beneath a house; the structure is somewhat peculiar (see photo). Length about $2\frac{1}{2}$ inches; length of single cell $\frac{3}{4}$ inch.

P. fuscicornis, Lep. A rare species from the interior.

P. chrysothorax, Web. A fairly common species only encountered on the coast lands. Large pendant nests nearly a foot in length and roughly cylindrical are constructed; the one actually observed was attached to a bush only a short distance from the ground. This nest was unfortunately destroyed in an attempt to secure it.

P. dimidiata, Oliv. Rockstone, Essequibo River.

P. constructrix, Sauss. Tumatumari, Essequibo River.

P. jurinei, Sauss. Issororo, N.W.D.

P. sericea, Oliv. Tumatumari, Essequibo River.

P. sycophanta, Gribodo. An uncommon species with a wide distribution.

P. velutina, Ducke. Issororo, N.W.D.

P. rejecta, F. A species with a wide distribution, but uncommon.

P. obidensis, Ducke. Puruni River.

Genus PROTOPOLYBIA, Ducke.

P. laboriosa, Sauss. Rockstone, Essequibo River.

Genus METAPOLYBIA, Ducke.

M. pediculata, Sauss. A fairly common species which builds its nests on beams beneath houses. The nest, which is constructed of papery material, is usually about 5 inches in diameter and about 1 inch in thickness and of a flattened irregular appearance. It is by no means a

conspicuous object, and is as the beam to which it has no hostile tendencies, and quite a safe undertaking, the interior.

Genus CHAR

C. chartarius, Oliv. A common species in the interior. The nest of this species has been previously described. They are prized by many and fetch a very fair price in Georgetown. They are sometimes seen attached to the rafters of the river. The taking of them is no easy matter, as the insect is very difficult to get from their home. Nests, however, may be taken by getting into the particular tree to which they are attached, and a well-directed blow of a stick will dislodge them, bearing the nest so that it may be eventually retrieved. An unsuccessful blow, however, is a case on record where the unfortunate native was killed by the river many feet below the nest.

Genus CH

C. pallidilineatus, Cam

Genus NE

N. bilineolata, Spin., v. far only taken in the Bo. nest was found attached to the rafters of sugar-cane. Greatest length $\frac{3}{4}$ inch.

N. scutellaris, F. Iss

N. lecheguana, Latr.

Genus S

S. surinama, L. A common species, which occurs at times in great numbers, and is attached to trees in the

bunta." A native method of to seize the nest quickly and then crush it. An individual black-skinned hands is likely to be

, var. *sampaioi*, Ducke. Appears foregoing in the interior, where it has been taken on the coast lands. It is attached to the under surfaces

species by no means of infrequent on the coast lands and interior. A small nest is attached to the floor-boards of a structure is somewhat peculiar about 2½ inches; length of single

a rare species from the interior.

A fairly common species only on the coast lands. Large pendant nests nearly cylindrical are constructed; one was attached to a bush only a few feet from the ground. This nest was unfortunately destroyed by an attempt to secure it.

Rockstone, Essequibo River.

Tumatumari, Essequibo River.

Issororo, N.W.D.

Tumatumari, Essequibo River.

o. An uncommon species with a

Issororo, N.W.D.

species with a wide distribution, but

Puruni River.

OTOPOLYBIA, Ducke.

Rockstone, Essequibo River.

ETAPOLYBIA, Ducke.

A fairly common species which occurs beneath houses. The nest, made of papery material, is usually about 1 inch in thickness and has a peculiar appearance. It is by no means a

conspicuous object, and is often of much the same colour as the beam to which it is attached. This insect shows no hostile tendencies, and the taking of the entire nest is quite a safe undertaking. It has only been observed in the interior.

Genus CHARTERGUS, Lepeletier.

C. chartarius, Oliv. A fairly common species in the interior. The nest of this species is a well-known object and has been previously described by a number of authors. They are prized by many colonists as "curios," and fetch a very fair price in Georgetown. These nests may sometimes be seen attached to the branches of trees overhanging the river. The taking of the nests is by no means an easy matter, as the insects resent any interference with their home. Nests, however, which overhang the river may be taken by getting a native to ascend quietly the particular tree to which the nest is attached and with one well-directed blow of a sharp cutlass sever the branch bearing the nest so that it falls into the river, whence it is eventually retrieved as soon as the wasps have left. An unsuccessful blow, however, spells disaster, and there is a case on record where such an incident occurred, and the unfortunate native rather than suffer jumped into the river many feet below and thus evaded the infuriated wasps.

Genus CHARTERGINUS, Fox.

C. pallidilineatus, Cameron. Issororo, N.W.D.

Genus NECTARINA, Shuckard.

N. bilineolata, Spin., var. *möbiana*, Sauss. A species so far only taken in the Botanic Gardens, Georgetown. The nest was found attached to the end of a dried "arrow" of sugar-cane. Greatest diameter about 1½ inches—depth ¾ inch.

N. scutellaris, F. Issororo, N.W.D.

N. lecheguana, Latr. Issororo, N.W.D.

Genus SYNOECA, Saussure.

S. surinama, L. A common species on the coast lands and occurs at times in the interior. The nests of this species, which are irregular in shape, may usually be seen attached to trees in the Botanic Gardens, Georgetown.

The whole of one side of the nest is attached to the tree, and the external wall exhibits a slight "ribbing." The nests are often a foot or more in length. The sting of this species is particularly formidable, and it does not hesitate to use it when occasion arises. The adult wasps may often be seen feeding on decaying fruit which has fallen on the ground, and they frequently visit certain species of flowers. I have taken specimens of this insect with the "pollinia" of a species of Euphorbiaceous plant attached to its legs; this is by no means a common occurrence.

S. irina, Spin. An uncommon species in the interior.

Genus APOICA, Lepeletier.

A. pallida, Oliv. A common species on the coast lands. The nest, which is invariably attached to a tree and never far from the ground, is disc-shaped. The under surface consists of innumerable cells, the ends of which are exposed. The under surface is invariably crowded with adult wasps, which attach themselves each to a particular cell and remain motionless, an aspect is thus presented of a cluster of wasps adhering to the under surface of the nest; in this position the bright yellow tips of their abdomens are very conspicuous. They do not readily move from this position, and if slightly disturbed will only make a slight movement. Once while observing a large nest of this species I happened to approach rather too near, when one wasp left the nest and stung me on the forehead immediately between the eyes, and returned at once to its original position. The effect of the sting was as though a blow had been delivered, and in a short space of time a swelling the size of a fowl's egg made its appearance, accompanied by considerable pain. The species is readily attracted to artificial light.

A. pallens, F. Issororo, N.W.D.

Genus MISCHOCYTTARUS, Saussure.

M. labiatus, F. A fairly common species on the coast lands. Only small nests are constructed.

Subfamily EUMENIDINAE.

Genus MONTEZUMIA, Sauss.

M. leprieuri, Spin. (1841) (= *M. rodwayi*, Cam. (1911)). An uncommon species taken in the vicinity of Georgetown.

M. nigriceps, Spin. One of the Hymenoptera on the coast lands. Particular economic value, as it hunts larvae of many agricultural pests, in (*Laphygma frugiperda*, S. & A.) and (*Mocis repanda*, F.). This wasp is searching for its prey amongst grass pendant, irregularly-shaped nest is what fragile in its structure. The and active community.

M. infernalis, Spin. An uncommon species on the coast lands. According to C. B. Williams the habits of this wasp, it constructs banks of the canals or trenches to its estates. Green caterpillars are stored in an egg is deposited, which is hidden in a burrow by a long stalk. The operation is small in comparison with the size of the nest.

M. infundibuliformis, F. Issororo.

Genus EUMENES,

E. canaliculata, Oliv. An extremely common species to be met with mostly on the coast lands. The nest is shaped mud cells, in small colonies. Sometimes more, are objects of common to the rafters beneath houses and walls and palings. These mud cells are about $\frac{1}{2}$ inch in height and with a diameter of $\frac{1}{4}$ inch. Some cells possess a kind of "neck" cell, giving it the appearance of a bottle. These cells are stored usually slightly over an inch in diameter, paralysed by the sting of the wasp. From four to six such larvae are stored in each cell. The wasp becomes mature in ten days to two weeks of emergence from the egg. The adult wasp bites a circular hole in one side of the cell, afterwards utilised by a species of spider and stored with spiders. A spider (*Cremastogaster*, sp.) finally colonises the cell. In constructing such cells the wasp uses a mass of moist and plastic mud in position, and the nest is built by means of the two front

of the nest is attached to the tree, exhibits a slight "ribbing." The r more in length. The sting of this formidable, and it does not hesitate on arises. The adult wasps may on decaying fruit which has fallen ey frequently visit certain species ken specimens of this insect with species of Euphorbiaceous plant this is by no means a common

uncommon species in the interior.

APOICA, Lepeletier.

common species on the coast lands. ariably attached to a tree and never is disc-shaped. The under surface e cells, the ends of which are exposed. r variably crowded with adult wasps, res each to a particular cell and repect is thus presented of a cluster of e under surface of the nest; in this llow tips of their abdomens are very o not readily move from this position, bed will only make a slight move- oserving a large nest of this species ch rather too near, when one wasp ng me on the forehead immediately returned at once to its original posi- he sting was as though a blow had a short space of time a swelling the ade its appearance, accompanied by he species is readily attracted to

oro, N.W.D.

CHOCYTARUS, Saussure.

fairly common species on the coast sts are constructed.

ily EUMENIDINAE.

MONTEZUMIA, Sauss.

(1841) (= *M. rodwayi*, Cam. (1911). taken in the vicinity of Georgetown.

M. nigriceps, Spin. One of the commonest species of *Hymenoptera* on the coast lands. Also a species of particular economic value, as it hunts for and destroys the larvae of many agricultural pests, including the Rice Worm (*Laphygma frugiperda*, S. & A.) and the Para-grass Worm (*Mocis repanda*, F.). This wasp may frequently be seen searching for its prey amongst grass and tall rice. A large pendant, irregularly-shaped nest is formed, which is somewhat fragile in its structure. These nests contain a large and active community.

M. infernalis, Spin. An uncommon species on the coast lands. According to C. B. Williams, who has observed the habits of this wasp, it constructs burrows in the clay banks of the canals or trenches to be found on all sugar estates. Green caterpillars are stored up in the nest, and an egg is deposited, which is hung from the roof of the burrow by a long stalk. The opening of the burrow is small in comparison with the size of the wasp.

M. infundibuliformis, F. Issororo, N.W.D.

Genus EUMENES, Latreille.

E. canaliculata, Oliv. An exceedingly common species to be met with mostly on the coast lands. Its dome-shaped mud cells, in small colonies of five or six and sometimes more, are objects of common observation attached to the rafters beneath houses and in sheltered spots on walls and palings. These mud cells (see photo) are usually about ½ inch in height and with a diameter of about ½ inch. Some cells possess a kind of "neck" at the top of the cell, giving it the appearance of a squat-shaped earthen bottle. These cells are stored with Lepidopterous larvae usually slightly over an inch in length and, of course, paralysed by the sting of the wasp at the time of capture. From four to six such larvae are stored. The wasp larva becomes mature in ten days to two weeks from the time of emergence from the egg. The pupal stage lasts slightly over a week. The adult wasp emerges from the cell by biting a circular hole in one side. These empty cells are afterwards utilised by a species of *Trypoxylon* for its nest, and stored with spiders. A small species of black ant (*Cremastogaster*, sp.) finally colonises these disused cells. In constructing such cells the adult wasp holds the piece of moist and plastic mud in position on the half-constructed nest by means of the two front pairs of legs, and the deli-

cate process of moulding the mud around the edge of the nest is performed with the mandibles, which are long and well suited to the work. During the operation the long antennae are bent downwards and kept rapidly moving about the work as though guiding it and preserving the symmetry of the structure. A small species of Chrysid was bred on one occasion from the mud cells of this *Eumenes*.

E. callimorpha, Sauss. Vicinity of Georgetown.

Genus PACHYMENES, Sauss.

P. pallipes, Oliv. Onderneeming, Essequibo.

Genus ZETHUS, Fab.

Z. mexicanus, L., var. *lugubris*, Perty. An uncommon species on the coast lands.

Z. gigas, Spin. Issororo, N.W.D.

Z. sicheliani, Sauss. Inhabiting disused borings in timber, Courantyne Coast, Berbice.

Genus ODYNERUS, Latr.

O. nasidens, Latr. Courantyne Coast, Berbice.

O. clavilineatus, Cameron. An uncommon species in the interior.

FOSSORES.

Family MUTILLIDAE

Genus MUTILLA, L.

M. (Thaumatmutilla) parallela, Klug. A fairly common species on the sandy soils of the interior.

M. mediata, F. From Courantyne Coast, Berbice. Taken while issuing from some disused borings in timber.

Genus THAUMATOMUTILLA, André.

T. ocellaris, Klug. Tumatumari, Essequibo River.

Family SCOLIADAE.

Subfamily TIPHIINAE.

Genus TIPHIA, Fab.

T. parallela, Smith. The larva of this wasp is parasitic on the larva of *Dyscinetus bidentatus*, Burm. (Coleoptera); fairly common on the coast lands.

some British 'G

Genus D

D. dorsata, F. Of common the Colony.

D. hyalina, Lep. (= *D. j* in most parts.

D. variegata, F. Issororo

Genus

E. flavopicta, Smith. T

Family RHOP

Genus RHOP

R. guianense, Schulz. A River.

Family PSAL

Genus PSE

P. comparata, Sm. Tun

P. chlorosoma, Sm. Pur

Subfamily

Genus CRY

C. purpureipes. A com

Genus

P. tinctipennis, Smith.

P. sapphiria, P. de B.

P. jucunda, Mocs. Roc

P. chlorotica, Mocs. Or

P. nigrescens, Smith. H

P. dimidiata, F. Rocks

P. amethystina, F. A c

the Colony.

P. seladonica, Dahlb. I

Subfamily PS

Genus POM

P. philadelphia, Lep. lands.

Genus *DIELIS*, Sauss.

D. dorsata, F. Of common occurrence in most parts of the Colony.

D. hyalina, Lep. (= *D. fallax*, Sauss.). Fairly common in most parts.

D. variegata, F. Issororo, N.W.D.

Genus *ELIS*, F.

E. flavopicta, Smith. Turkeyn, East Coast, Demerara.

Family RHOPALOSOMIDAE.

Genus RHOPALOSOMA, Schulz.

R. guianense, Schulz. At light, Rockstone, Essequibo River.

Family PSAMMOCHARIDAE.

Genus PSEUDAGENIA, Kohl.

P. comparata, Sm. Tumatumari, Essequibo River.

P. chlorosoma, Sm. Puruni River.

Subfamily PEPSINAE.

Genus CRYPTOCHILUS, Panz.

C. purpureipes. A common species on the coast lands.

Genus PEPSIS, Fab.

P. tinctipennis, Smith. Issororo, N.W.D.

P. sapphiria, P. de B. Rockstone, Essequibo River.

P. jucunda, Mocs. Rockstone, Essequibo River.

P. chlorotica, Mocs. Onderneeming, Essequibo.

P. nigrescens, Smith. Rockstone, Essequibo River.

P. dimidiata, F. Rockstone, Essequibo River.

P. amethystina, F. A common species in most parts of the Colony.

P. seladonica, Dahlb. Issororo, N.W.D.

Subfamily PSAMMOCHARINAE.

Genus POMPILOGASTER, Ashm.

P. philadelphica, Lep. A common species on the coast lands.

Subfamily *SPHECINAE*.Genus *SCELIPHRON*, Klug.

S. fistulare, Dahlb. This species is widely distributed. Its mud nests are common objects on palings, beneath houses, and at times within the house itself, behind pictures, etc. The nests are somewhat irregular in shape and may contain as many as eight or ten cells. Sometimes only two cells are constructed together, when the architecture is then better demonstrated. The usual procedure is to build one cell and attach it firmly, and then to build other cells around it. Each cell is stored with spiders, from sixteen to twenty according to the size of the spiders. As soon as the requisite number of spiders has been obtained the cell is closed up. The egg, according to C. B. Williams, is deposited on dorsal surface of the abdomen of the first spider placed in the cell. The following observations by Mr. Williams on the habits of this wasp are of interest. A wasp which was engaged in constructing its mud cells was observed, and its movements recorded as follows—

12.5' 30" p.m. gone for mud,
12.9' 10" p.m. returned,
12.10' 15" p.m. gone,
12.17' 58" p.m. returned,
12.19' 20" p.m. gone again,
12.22' 20" p.m. returned,
12.23' 10" p.m. gone.

At 12.37' 45" the wasp entered its nest and remained inside; 12.38' 50" the wasp flew away, and on inspection the nest was found to contain a spider (Fam. *Argiopidae*—*Gasteracanthinae*) with an egg on dorsal surface of abdomen. This spider was eventually removed by myself. At 12.51' 15" the wasp returned with another spider, and went away 12.52' 15". At 4.10 p.m. the nest was almost completely closed. The egg of this species is sausage-shaped and slightly curved, dull yellowish white in colour, with small almost transparent areas towards each end; length 3.4 mm. Young larvae when first hatched out are almost 4 mm. in length. They soon attach themselves to a spider, which they proceed to consume, leaving only the legs. Development is completed in about two weeks. The larva then commences to spin its cocoon, which occupies several days. The cocoon is dark brown in

colour, and of a papery consistency lasts a little over a week, its way out through the mud neat circular hole. The small *cordata*, L., often utilises the A fungus occasionally destroys finally the young larvae. No has been observed up to the p that only about 60 % of the wasp.

S. figulus, Dahlb. Closely and with similar habits and numerous.

Genus *Sphex*

S. ichneumoneus, L. A common species throughout the Colony it usually chooses a dry piece of a path or other places passing. While constructing nest it is a most conspicuous on the surface of the soil. made by Mr. C. B. Williams instance the insect was found in swampy land at the foot of from the river invariably commenced about 1.40 about $\frac{1}{2}$ inch deep. At 2.1 backwards, remaining below came out again. Burrowing nests are stored with four hoppers, either nymphs or nest is not closed up by grasshopper."

Sphex (Isodontia) costipes

S. (Harpactopus) thomasi which does not occur on burrows in soil and stores which is placed head first in at the base of the grasshopper

Genus

P. denticulatum, Sm. distributed species.

Subfamily SPHECINAE.

Genus SCELIPHRON, Klug.

Dahlb. This species is widely distributed. It is common on palings, beneath eaves within the house itself, behind pictures. The nests are somewhat irregular in shape and many as eight or ten cells. Sometimes they are constructed together, when the architecture is demonstrated. The usual procedure is to dig a hole and attach it firmly, and then to build up the cells. Each cell is stored with spiders, twenty according to the size of the spiders. A large number of spiders has been obtained from the nests. The egg, according to C. B. Williams, is on the dorsal surface of the abdomen of the first cell. The following observations by C. B. Williams on the habits of this wasp are of interest. The wasp was engaged in constructing its mud cells and its movements recorded as follows—

12.5' 30" p.m. gone for mud,
12.9' 10" p.m. returned,
12.10' 15" p.m. gone,
12.17' 58" p.m. returned,
12.19' 20" p.m. gone again,
12.22' 20" p.m. returned,
12.23' 10" p.m. gone.

The wasp entered its nest and remained for 50" the wasp flew away, and on inspection found it to contain a spider (Fam. *Argiopidae*—*Argiope*) with an egg on dorsal surface of abdomen. The egg was eventually removed by myself. At 12.23' the wasp returned with another spider, and at 12.23' 15". At 4.10 p.m. the nest was almost completed. The egg of this species is sausage-shaped, slightly curved, dull yellowish white in colour, with most transparent areas towards each end;

Young larvae when first hatched out are 1/2 inch in length. They soon attach themselves to the mud and proceed to consume, leaving only the pupa. Development is completed in about two weeks. The pupa commences to spin its cocoon, which is completed in about 10 days. The cocoon is dark brown in

colour, and of a papery consistency, easily broken. Pupation lasts a little over a week, and the adult wasp then bites its way out through the mud wall of the nest, leaving a neat circular hole. The small green metallic bee, *Euglossa cordata*, L., often utilises these disused cells for its nest. A fungus occasionally destroys the stored spiders and finally the young larvae. No actual parasite of the species has been observed up to the present. It appears, however, that only about 60% of the stored cells produce an adult wasp.

S. figulus, Dahlb. Closely allied to the above species, and with similar habits and distribution, though not so numerous.

Genus SPHEX, L.

S. ichneumoneus, L. A common and widely distributed species throughout the Colony. For its subterranean nests it usually chooses a dry piece of soil; often in the centre of a path or other places where people are frequently passing. While constructing the initial excavation for its nest it is a most conspicuous object, moving rapidly about on the surface of the soil. The following observations made by Mr. C. B. Williams are of interest. "In this instance the insect was boring into soft pegass soil on swampy land at the foot of a hill. High spring tides from the river invariably cover the land. The burrow was commenced about 1.40 p.m., and at 1.45 p.m. it was about 1/2 inch deep. At 2.19 the wasp entered the hole backwards, remaining below a few minutes, and then came out again. Burrowing continued till 4 p.m. These nests are stored with four or five short-winged grasshoppers, either nymphs or brachypterous species; the nest is not closed up between the insertion of each grasshopper."

Sphex (Isodontia) costipennis, Spin. Issororo, N.W.D.

S. (Harpactopus) thomae, F. An uncommon species which does not occur on the coast lands. Constructs burrows in soil and stores them with a single grasshopper, which is placed head first in the nest. The egg is attached at the base of the grasshopper's hind-leg (C. B. Williams).

Genus PODIUM, F.

P. denticulatum, Sm. An uncommon though widely distributed species.

Subfamily PHILANTHINAE.

Genus CERCERIS, Latr.

C. (Trachypus) mexicanus, Cameron. An uncommon species. Taken while emerging from its burrow in sandy soil at the base of a tree. Plantation Bath, Berbice.

Subfamily BEMBICINAE.

Genus BEMIDULA, Burm.

B. discisa, Tasch. Upper Demerara River.

B. variegata, Ol. Upper Demerara River.

Genus MONEDULA, Latr.

M. pantherina, Handl. A fairly common species in some parts of the interior, especially where sandy soils exist. It is quite the largest and fiercest of these insects to be found in the Colony. Its burrows are stored with *Tabanidae*.

M. punctata, Lep. An uncommon species in the interior.

M. signata, Latr. The commonest and most widely distributed *Bembex* in British Guiana. It may frequently be seen about mules and cattle in the pastures waiting to carry off the *Tabanidae* which are always present about these animals. Their burrows may be seen in sandy soil, and usually occur in colonies of twenty or thirty. A large species of Bombyliid fly may be seen about these burrows, occasionally entering them. Its exact relationship to the wasp has never been actually traced, but it is quite possibly parasitic. The fly belongs to the genus *Anthrax*.

Subfamily GORYTINAE.

Genus GORYTES, Latr.

G. brasiliensis, Shuck. Rockstone, Essequibo River.

Subfamily LARRINAE.

Genus TACHYTES, Panzer.

T. ametina, Cameron. Rockstone, Essequibo River.

Genus LARRA, Fab.

L. rubricata, Smith. Rockstone, Essequibo River.

Subfamily TRYPOX

Genus TRYPOXYL

T. palliditarse, Sauss. A fairly common species on the coast lands.

T. fugax, F. This insect has been taken from the cells of *Eumenes canaliculata*, Ol. occurrence.

T. brevicarinatum, Cameron. A fairly common distributed species all over the Colony. It constructs its nest of mud cells beneath the ground, sometimes as many as twenty-five cells cemented together. The largest cells are about 1/4 of an inch in length and about 1/4 of an inch in diameter. Spiders are stored up by this species.

T. albitarse, F. Bred from a nest. Potaro River.

T. nitidum, Sm. Issororo, N.W.

T. cinereum, Cameron. The habit of this species has been observed by Mr. C. B. Williams. The cell was constructed by one of the workers of the woodwork of a door. When the 'ball' of mud is collected and a nest is gradually drawing it down. A shrill buzzing sound is emitted during the process. Seconds to one minute are occupied in the construction of mud. Spiders are stored."

Family TRIGON

Genus TRIGONALY

T. melanoleuca, Westwood. A fairly common species emerged from the pupae of the *Meloidae*, which had been taken from the soil on a sugar estate.

Family CHRYS

Subfamily CHRY

Genus CHRYS

C. punctatissima, Spin. A common species on the coast lands.

C. distinctissima, Dahlb. Bar

C. parvula, F. East Coast, D

TRANS. ENT. SOC. LOND. 1917.—P

ILANTHINAE.

CRICERIS, Latr.

s, Cameron. An uncommon
ing from its burrow in sandy
Plantation Bath, Berbice.

EMBICINAE.

BIDULA, Burm.

Demerara River.
Demerara River.

VEDULA, Latr.

fairly common species in some
cially where sandy soils exist.
l fiercest of these insects to
Its burrows are stored with

common species in the interior.
commonest and most widely
ish Guiana. It may frequently
ttle in the pastures waiting to
hich are always present about
ows may be seen in sandy soil,
nies of twenty or thirty. A
fly may be seen about these
ing them. Its exact relation-
er been actually traced, but it
The fly belongs to the genus

GORYTINAE.

ORYTES, Latr.

Rockstone, Essequibo River.

LARRINAE.

HYTES, Panzer.

Rockstone, Essequibo River.

LARRA, Fab.

ckstone, Essequibo River.

Subfamily TRYPOXYLINAE.

Genus TRYPOXYLON, Latr.

T. palliditarse, Sauss. A fairly common species on the
coast lands.

T. fugax, F. This insect has been bred from the disused
cells of *Eumenes canaliculata*, Ol. It is of fairly common
occurrence.

T. brevicarinatum, Cameron. A common and widely
distributed species all over the Colony. It usually con-
structs its nest of mud cells beneath a palm leaf. Some-
times as many as twenty-five cylindrical cells are found
cemented together. The largest cells are nearly an inch
in length and about $\frac{1}{4}$ of an inch in diameter. Small
spiders are stored up by this species.

T. albitarse, F. Bred from a large-sized rounded mud
nest. Potaro River.

T. nitidum, Sm. Issororo, N.W.D.

T. cinereum, Cameron. The habits of this species have
been observed by Mr. C. B. Williams. "An elongate mud
cell was constructed by one of these wasps in the angle
of the woodwork of a door. When making these nests a
'ball' of mud is collected and applied to the side of the
nest, gradually drawing it downwards as it adheres. A
shrill buzzing sound is emitted the whole time. Thirty
seconds to one minute are occupied in applying one 'ball'
of mud. Spiders are stored."

Family TRIGONALIDAE

Genus TRIGONALYS, Westw.

T. melanoleuca, Westwood. A number of these insects
emerged from the pupae of the Noctuid moth *Amyra octo*,
Guen., which had been taken from just beneath the surface
of the soil on a sugar estate.

Family CHRYSIDIDAE

Subfamily CHRYSIDINAE.

Genus CHRYSIS, L.

C. punctatissima, Spin. A common species on the coast
lands.

C. distinctissima, Dahlb. Bartica, Essequibo River.

C. parvula, F. East Coast, Demerara.

TRANS. ENT. SOC. LOND. 1917.—PARTS II, III, IV. (MAY '18) Y

Family BRACONIDAE.

Subfamily CHELONINAE.

Genus FORNICIA, Brullé.

F. clathrata, Brullé. A larval parasite of the Limacodid moth, *Sibine fusca*, Stoll. This is a very abundant moth on the coast lands and attacks a large variety of economic plants. The parasite emerges before the larva is half developed. The percentage of parasitism is never very high.

Subfamily BRACONINAE.

Genus LASIOPHORUS, Hal.

L. fortispinus, Cameron. A fairly common species on the coast lands.

Genus IPHIAULAX, Forster.

I. medianus, Cameron. This is the well-known parasite of the several species of *Diatraea* (Lepidoptera) which bore into sugar-cane in British Guiana. The species is a common one, it being an easy matter to capture either sex in the cane-fields. The life-history of the species has been observed in the field by Mr. Harold Moore, and the following is his own description: *—"On the morning of the 12th, in a field at Non Pareil, I noticed a female alight near the top of a cane-stump, run down the shoot, tapping it the while with her antennae, till she got to a *Diatraea* hole near the base. Into the tunnel she began pushing her ovipositor, when a slight movement on my part, in an endeavour to obtain a closer view of the proceedings, disturbed her. Off she flew, but merely to wheel round and alight again on the shoot, when she found the hole the same way, and began inserting her ovipositor. I lay prostrate on the trash bank, as motionless as possible, and watched. It was 9.28 a.m. Soon after her ovipositor had been inserted she gave a slight quick backward jerk with her abdomen, showing she had evidently felt something. Back down went her abdomen and again a similar jerk. This occurred three or four times, after which she kept quite still for several minutes, her ovipositor being well in the caterpillar's tunnel. She withdrew her ovipositor at 9.38, thus occupying about ten minutes in parasitising the caterpillar, of the presence of which I had no doubt.

* Published in the "Daily Argosy," Demerara, March 1913.

I cut out the cane-shoot and examine it till next morning tained a caterpillar, and in the egg of the parasite. It is at the widest part—whitish, narrow, elongated, tapering where there is a small curve at the end is evidently for the in place. From the above is paralysed by the parasite. From observations made on it would appear that paralysis not always, occur immediately. The firm texture of the egg should the caterpillar happen 3.30 p.m. on the same day (had hatched and that the feeding, attached to the side caterpillar. It had probably The egg-period is therefore On the morning of the sixteenth almost the whole posterior half the parasite larva. By the parasite larva had completed and was spinning a cocoon larval period, too, is very shortness of both the egg and readily understood, when it requires fresh food, and has which to feed. It of necessity it rapidly, before extreme do all its growing in that shoot at the angle between the shoot the tunnel, probably on account through drying of the shoot know that the larval period. On the morning of the 31st Pupal period therefore about egg-laying to adult is the The cocoon is transparent woven silk, giving a membrane cylindrical, blunted at both ends 2.1 cm., diameter .4 cm.

I. obscuricarinatus, Cameron

BRACONIDAE.

Family CHELONINAE.

FORNICIA, Brullé.

A larval parasite of the Limacodid
boll. This is a very abundant moth
attacks a large variety of economic
emerges before the larva is half
percentage of parasitism is never very

Family BRACONINAE.

LASIOPHORUS, Hal.

erion. A fairly common species on

IPHIAULAX, Forster.

on. This is the well-known parasite
of *Diatraea* (Lepidoptera) which
in British Guiana. The species is a
g an easy matter to capture either
The life-history of the species has
field by Mr. Harold Moore, and the
description: *—"On the morning of
Non Pareil, I noticed a female alight
-stump, run down the shoot, tapping
antennae, till she got to a *Diatraea*
Into the tunnel she began pushing
a slight movement on my part, in
ain a closer view of the proceedings,
he flew, but merely to wheel round
the shoot, when she found the hole
egan inserting her ovipositor. I lay
a bank, as motionless as possible, and
8 a.m. Soon after her ovipositor had
ve a slight quick backward jerk with
ng she had evidently felt something.
r abdomen and again a similar jerk.
or four times, after which she kept
l minutes, her ovipositor being well
unnel. She withdrew her ovipositor
ing about ten minutes in parasitising
e presence of which I had no doubt.

"Daily Argosy," Demerara, March 1913.

I cut out the cane-shoot and took it home. I could not
examine it till next morning (13th), when I found it con-
tained a caterpillar, and in the same tunnel with it was
the egg of the parasite. It is about 3 mm. by about 1 mm.
at the widest part—whitish, translucent, firm or leathery—
narrow, elongated, tapering to a long point posteriorly,
where there is a small curved blackish hook. The hook
at the end is evidently for the purpose of keeping the egg
in place. From the above it seems that the caterpillar
is paralysed by the parasite at the time of oviposition.
From observations made on previous occasions, however,
it would appear that paralysis does not, or at least does
not always, occur immediately, but comes on gradually.
The firm texture of the egg prevents it from being injured
should the caterpillar happen to wriggle against it. At
3.30 p.m. on the same day (13th) I observed that the egg
had hatched and that the young larva was peacefully
feeding, attached to the side of the ninth segment of the
caterpillar. It had probably hatched during the forenoon.
The egg-period is therefore very short—about one day.
On the morning of the sixteenth the caterpillar was dead,
almost the whole posterior half having been devoured by
the parasite larva. By the morning of the 17th the
parasite larva had completely devoured the caterpillar,
and was spinning a cocoon in order to pupate. The
larval period, too, is very short—about four days. The
shortness of both the egg and larval stages can be very
readily understood, when it is remembered that the larva
requires fresh food, and has only a single caterpillar on
which to feed. It of necessity therefore has to get through
it rapidly, before extreme decomposition is reached, and
do all its growing in that short time. The larva pupated
at the angle between the shoot and the glass jar. It left
the tunnel, probably on account of the latter shrinking
through drying of the shoot. From other observations I
know that the larval period is sometimes certainly a week.
On the morning of the 31st the perfect insect emerged.
Pupal period therefore about fourteen days. Time from
egg-laying to adult is therefore about nineteen days."
The cocoon is transparent and constructed of closely
woven silk, giving a membranous appearance. In shape
cylindrical, blunted at both ends. Colour whitish. Length
2.1 cm., diameter .4 cm.

I. obscuricarinatus, Cameron. Issororo, N.W.D.

Subfamily RHOGADINAE.

Genus MEGARHOGAS, Szep.

M. fuscipalpis, Cam. Rockstone, Essequibo River.

Subfamily AGATHINAE.

Genus SPILOMICRODUS, Cam.

S. nigriceps, Cam. Bartica, Essequibo River.

Family CHALCIDIDAE.

Subfamily TRICHOGRAMMATINAE.

Genus TRICHOGRAMMA, Westw.

T. minutum, Riley. This well-known parasite is of common occurrence in the Colony as an egg-parasite of the Cane Borers (*Diatraea*) and several other species of Lepidoptera. Its life-history and habits when parasitising the ova of *Diatraea* have been fully worked out by myself and published in the Journal of the Board of Agriculture, British Guiana, vol. vi, No. 4.

Subfamily CHALCIDINAE.

Genus CHALCIS, F.

C. pandora, Cwfd. Bred from the pupa case of a species of Hesperid (sp. indet.) butterfly feeding on cane blades. An uncommon species.

C. annulata, F. A common parasite of *Brassolis sophorae*, L. (Lepidoptera) and also *Caligo ilioneus ilioneus*, Cramer (Lepidoptera). It is widely distributed and the commonest Chalcid in the Colony.

Genus SMICRA, Spin.

S. fulvomaculata, Cam. Bred from the pupa of a Hesperid butterfly (sp. indet.). An uncommon species.

Family EVANIIDAE.

Subfamily EVANIINAE.

Genus EVANIA, F.

E. erythrasis, Cameron. Issororo, N.W.D.

E. appendigaster, L. This insect emerged from a large box containing a quantity of seed rice packed in bags. Innumerable Blattid egg-cases (*Periplaneta americana* L.), as well as adult insects, were in the box.



Trans

NOTICE TO

Sketch Map facing p. 164, Part I.



BRITISH GUIANA HYM.