

8-22-1996

The Role of Calcium in the Regulation of Apoptosis

David J. McConkey

The University of Texas M.D. Anderson Cancer Center, David_McConkey@isqm.mda.uth.tmc.edu

Follow this and additional works at: <https://digitalcommons.usu.edu/microscopy>



Part of the [Biology Commons](#)

Recommended Citation

McConkey, David J. (1996) "The Role of Calcium in the Regulation of Apoptosis," *Scanning Microscopy*. Vol. 10 : No. 3 , Article 14.

Available at: <https://digitalcommons.usu.edu/microscopy/vol10/iss3/14>

This Article is brought to you for free and open access by the Western Dairy Center at DigitalCommons@USU. It has been accepted for inclusion in Scanning Microscopy by an authorized administrator of DigitalCommons@USU. For more information, please contact digitalcommons@usu.edu.



THE ROLE OF CALCIUM IN THE REGULATION OF APOPTOSIS

David J. McConkey

Department of Cell Biology - 173, The University of Texas M.D. Anderson Cancer Center,
1515 Holcombe Boulevard, Houston, Texas 77030

Telephone number: (713) 792-8591 / FAX number (713) 792-8747

E.mail: David_McConkey@isqm.mda.uth.tmc.edu

(Received for publication May 8, 1995 and in revised form August 22, 1996)

Abstract

The recognition that apoptosis is regulated by an evolutionarily conserved set of polypeptides from the nematode *Caenorhabditis elegans* to humans suggests that a conserved set of biochemical mechanism(s) may also be involved in the response. Early evidence suggested that the endogenous endonuclease implicated in apoptosis in most model systems is Ca^{2+} -dependent, and subsequent work from a number of independent laboratories suggests that alterations in cytosolic Ca^{2+} homeostasis are one of the conserved biochemical pathways regulating the response. Molecular targets for Ca^{2+} are now being identified and include signal transduction intermediates, endonuclease(s) and proteases, and the enzymes involved in the maintenance of phospholipid asymmetry in the plasma membrane. Furthermore, interesting preliminary work suggests that BCL-2 suppresses apoptosis via a mechanism that is linked to intracellular Ca^{2+} compartmentalization, and it appears that Ca^{2+} alterations in some examples of apoptosis occur as the result of changes within the mitochondria. This review will summarize what is known about the role of Ca^{2+} in the regulation of apoptosis and discuss how Ca^{2+} might interact with some of the other biochemical signals implicated in cell death.

Key Words: Apoptosis, calcium, endonuclease, oxidative stress, BCL-2.

Introduction

Apoptosis (programmed cell death) is a highly regulated process of selective cell deletion involved in development, normal cell turnover, hormone-induced tissue atrophy, cell-mediated immunity, tumor regression, and a growing number of pathological disorders, typified by acquired immune deficiency syndrome (AIDS) and Alzheimer's diseases (Thompson, 1995; Wyllie *et al.*, 1980). The response is characterized by stereotyped morphological alterations, including plasma and nuclear membrane blebbing, organelle relocalization and compaction, chromatin condensation, and the formation of membrane-enclosed structures termed "apoptotic bodies" that are extruded into the extracellular milieu (Savill *et al.*, 1993; Wyllie *et al.*, 1980). Uptake of apoptotic debris is carefully controlled, as apoptotic cells and bodies are specifically recognized and cleared by neighboring epithelial cells and professional phagocytic cells (macrophages) before their contents can be released into the extracellular milieu, thereby allowing for cell death to occur in the absence of inflammation (Savill *et al.*, 1993).

Apoptosis has historically been characterized biochemically by endogenous endonuclease activation, resulting first in the production of domain-sized large (50-300 kilobase) DNA fragments (Brown *et al.*, 1993; Filipinski *et al.*, 1990; Oberhammer *et al.*, 1993) and subsequently into oligonucleosomal cleavage products commonly referred to as "DNA ladders" (Wyllie, 1980). It appears that DNA ladders are derived from the larger DNA fragments but that these events may be mediated by different enzymatic activities that can be distinguished by their divalent cation requirements (Sun and Cohen, 1994). In addition to endonuclease activation, more recent work has demonstrated that a family of cysteine proteases homologous to the *Caenorhabditis elegans* (*C. elegans*) cell death gene *ced-3* and human interleukin-1 converting enzyme (ICE) are also critically involved in the response, as inhibitors of these enzymes block both endonuclease activation and cell death (Fernandes-Alnemri *et al.*, 1994, 1995; Kumar *et al.*, 1994; Martin




Figure 1. Schematic representation of phospholipase C-dependent and damage-mediated pathways involved in promoting Ca^{2+} elevations in apoptotic cells.

and Green, 1995; Nicholson *et al.*, 1995; Tewari *et al.*, 1995; Wang *et al.*, 1994; Yuan *et al.*, 1993). The important substrates for these proteases remain largely unidentified but include poly (ADP-ribose) polymerase (PARP), the lamins, and a viral inhibitor of their activity (baculovirus p35) (Bump *et al.*, 1995; Kaufmann *et al.*, 1993; Lazebnik *et al.*, 1993; Tewari *et al.*, 1995; Xue and Horvitz, 1995). Although many of the important biochemical alterations associated with apoptosis occur within the nucleus, recent work indicates that apoptosis can also be initiated in enucleated cells (cytoplasts) (Jacobson *et al.*, 1994; Schulze-Osthoff *et al.*, 1994), and may lead investigators to suspect that the process is initiated in the cytoplasm.

At the molecular level, apoptosis is regulated by many familiar oncogenes (*bcl-2*, *myc*, *ras*, *abl*, *fos*) and tumor suppressor genes (*p53*, *Rb*) (Thompson, 1995). However, the family of polypeptides homologous to the *bcl-2* appear to collectively be the most important. Regulation of apoptosis susceptibility is their only established function to date. One class of *bcl-2* homologs (including *bcl-2*, *bcl-xL*, *mcl-1*, and several viral proteins) suppress apoptotic cell death, while another group (*bax*, *bcl-xs* and *bak*) promotes apoptosis sensitivity (Oltvai

and Korsmeyer, 1994). Thus, overexpression of BCL-2 or BCL-XL blocks apoptosis induced by very diverse stimuli, including growth factor withdrawal, tumor necrosis factor, engagement of the Fas antigen, ionizing radiation, oncogenes such as *myc*, and chemical chemotherapeutic agents (Thompson, 1995). Moreover, the apoptosis-regulatory functions of BCL-2 and its homologs are evolutionarily conserved, as the *C. elegans* cell death suppressor *ced-9* is a structural and functional homolog of human *bcl-2* (Hengartner *et al.*, 1992; Hengartner and Horvitz, 1994). Together, these observations have suggested to many investigators that BCL-2 regulates a central biochemical signal in the pathway to cell death, the identity of which is still unknown. However, several strong candidates are emerging. Work from our laboratory (McConkey *et al.*, 1989a, 1989b, 1994) and others (Shi *et al.*, 1989; Martikainen and Isaacs, 1990; Story *et al.*, 1992) over the past five years has demonstrated that alterations (Baffy *et al.*, 1993; Lam *et al.*, 1994) in intracellular Ca^{2+} homeostasis are commonly involved in promoting apoptosis, and more recent work suggests that one aspect of BCL-2 function involves preventing these alterations (Baffy *et al.*, 1993; Lam *et al.*, 1994). Here, we will discuss the role of

Ca²⁺ in regulating apoptosis and compare it with other general mediators of apoptotic cell death.

Calcium Alterations in Apoptosis

Early studies by Kaiser and Edelman (1977) demonstrated that glucocorticoid-stimulated apoptosis is associated with enhanced Ca²⁺ influx, work that provided the first evidence that increases in intracellular Ca²⁺ might be involved in triggering apoptosis. We have since confirmed that glucocorticoids induce elevations in the cytosolic Ca²⁺ concentration in thymocytes via enhanced Ca²⁺ influx across the plasma membrane (McConkey *et al.*, 1989b). Recent results from Synder's laboratory have implicated the type 3 receptor for inositol trisphosphate (IP₃R) in Ca²⁺ influx, DNA fragmentation, and cell death, suggesting that these physiologically relevant Ca²⁺ channels play a central role in the response (Khan *et al.*, 1996). However, intracellular Ca²⁺ storage sites also appear to be affected, as the Ca²⁺ pool located in the endoplasmic reticulum is depleted in a lymphoid cell line in response to glucocorticoid treatment (Lam *et al.*, 1993), and a similar phenomenon has been documented in an interleukin 3 (IL-3)-dependent myeloid cell line undergoing apoptosis following IL-3 withdrawal (Baffy *et al.*, 1993). Circumstantial evidence suggests that the mitochondrial Ca²⁺ pool may also be affected (Richter, 1993), as mitochondrial membrane potential drops very early during apoptosis (Petit *et al.*, 1995; Zamzani *et al.*, 1995a, 1995b), and it is well known that the maintenance of mitochondrial Ca²⁺ homeostasis is dependent upon mitochondrial membrane potential (Richter, 1993). Apoptosis in other systems also appears to involve elevations in the cytosolic Ca²⁺ concentration. For example, rapid, sustained Ca²⁺ increases precede the cytotoxicity of the cellular targets of cytotoxic T lymphocytes (Allbritton *et al.*, 1988) and natural killer (NK) cells (McConkey *et al.*, 1990). In developing T lymphocytes, high affinity engagement of the T cell receptor induces apoptosis (McConkey *et al.*, 1989a; Murphy *et al.*, 1990; Shi *et al.*, 1989; Smith *et al.*, 1989) that involves a sustained Ca²⁺ elevation (McConkey *et al.*, 1989a; Nakagama *et al.*, 1992).

Both second messenger- and damage-mediated mechanisms can be involved in promoting Ca²⁺ increases in apoptotic cells (Fig. 1). In an example of the former, T cell receptor engagement on thymocytes leads to a sustained increase in the cytosolic Ca²⁺ concentration that involves protein tyrosine kinase activation, phosphorylation of the γ -isoform of phospholipase C, phosphoinositide hydrolysis leading to the production of inositol trisphosphate (IP₃), and mobilization of Ca²⁺ from the endoplasmic reticulum and extracellular milieu that promote cell death (McConkey *et al.*, 1989a, 1994).

Similarly, surface antigen receptor engagement on B cells leads to Ca²⁺ increases that promote cell death (Norvell *et al.*, 1995; Parry *et al.*, 1994; Tsubata *et al.*, 1993; Yao and Scott, 1993). Thus, in these examples of apoptosis, Ca²⁺ increases occur via a controlled, physiological mechanism that is also utilized in alternative responses such as cellular activation leading to proliferation.

Work from our laboratory (Fernandez *et al.*, 1995) and others (Richter, 1993) has revealed another mechanism that is involved in promoting sustained cytosolic Ca²⁺ increases in apoptotic cells. It is well known that cytosolic Ca²⁺ concentration is maintained at roughly 100 nM in resting cells, whereas the concentrations in the extracellular milieu, and the ER, are much higher (in the millimolar range). Early work on the biochemical mechanisms underlying the cytotoxicity of agents that generate reactive oxygen species in cells (oxidative stress) indicated that the Ca²⁺ transport systems localized to the ER, mitochondria, and plasma membrane can be damaged by oxygen radicals (Orrenius *et al.*, 1989). This leads to diffusion of Ca²⁺ down its concentration gradient, a disruption of intracellular Ca²⁺ homeostasis, and sustained Ca²⁺ increases. Oxidative stress is now known to be commonly involved in apoptosis (Buttke and Sandstrom, 1994; Dypbukt *et al.*, 1994; Fang *et al.*, 1995; Hockenbery *et al.*, 1993; Lennon *et al.*, 1990), and it is therefore possible that oxidative disruption of intracellular Ca²⁺ homeostasis is involved in these systems. Supporting this idea, Fernandez *et al.* (1995) have recently shown that the glucocorticoid-induced Ca²⁺ increase observed in thymocytes is blocked by antioxidants, and Kroemer's laboratory has presented evidence that oxidative stress leads to disruption of mitochondrial Ca²⁺ stores (Zamzani *et al.*, 1995b).

Direct evidence that Ca²⁺ increases can mediate apoptotic endonuclease activation and cell death has been obtained from experiments with intracellular Ca²⁺ buffering agents and extracellular Ca²⁺ chelators. Our group (Aw *et al.*, 1990; Bellomo *et al.*, 1992; McConkey *et al.*, 1989a, 1989b, 1990, 1994; Zhivotovsky *et al.*, 1993) and others (Robertson *et al.*, 1993; Story *et al.*, 1992) have shown that these agents can inhibit both DNA fragmentation and death in apoptotic cells. The Ca²⁺-dependent regulatory cofactor calmodulin may link these Ca²⁺ alterations to the effector machinery, as we and others have shown that calmodulin antagonists can interfere with apoptosis in some of these systems (Dowd *et al.*, 1991; McConkey *et al.*, 1989b) and increases in calmodulin expression are linked to apoptosis in glucocorticoid-treated thymoma cells (Dowd *et al.*, 1991) and in prostatic epithelial cells following withdrawal of androgen (Furuya and Isaacs, 1993). Independent evidence for the involvement of Ca²⁺ influx in the trigger-

ing of apoptosis has come from studies with specific Ca^{2+} channel blockers, which abrogate apoptosis in the regressing prostate following testosterone withdrawal (Martikainen and Isaacs, 1990) and in pancreatic β -cells treated with serum from patients with type I diabetes (Juntti-Berggren *et al.*, 1993).

Other support for the involvement of Ca^{2+} in apoptosis comes from the observation that agents which directly mobilize Ca^{2+} can trigger apoptosis in diverse cell types. Early work by Kaiser and Edelman (1978) demonstrated the cytolytic effects of glucocorticoids on lymphoid cells can be mimicked by treating the cells with Ca^{2+} ionophores. Subsequently, Wyllie *et al.* (1984) demonstrated that Ca^{2+} ionophores cause endonuclease activation as well as many of the morphological changes that are typical of apoptosis in thymocytes. Calcium ionophores also trigger apoptosis in prostate tumor cells (Martikainen and Isaacs, 1990) and in non-metastatic melanoma lines (McConkey, unpublished). Independent evidence for the general relevance of this mechanism has come from studies with the endoplasmic reticular Ca^{2+} -ATPase inhibitor thapsigargin, the product of the plant, *Thapsa garganica*, which can also trigger all of the morphological and biochemical events of apoptosis in thymocytes (Jiang *et al.*, 1994) and some other cell types (Choi *et al.*, 1995; Kaneko and Tsukamoto, 1994; Levick *et al.*, 1995).

A final argument for a central role for Ca^{2+} in regulating apoptosis comes from recent and ongoing work on the biochemical mechanisms of apoptosis suppression by the BCL-2 oncoprotein (Baffy *et al.*, 1993; Lam *et al.*, 1994; Zornig *et al.*, 1995; Marin *et al.*, 1996). The possible relationship between Ca^{2+} and BCL-2 was first suggested by work by Baffy *et al.* (1993), who showed that BCL-2 can block the depletion of the endoplasmic reticular Ca^{2+} pool in transfectants of an IL-3-dependent cell line (32D). Interestingly, Baffy *et al.* (1993) also demonstrated that constitutive levels of Ca^{2+} in mitochondria (measured following treatment with an uncoupler that promotes rapid and selective depletion of this intracellular Ca^{2+} store) were significantly lower in BCL-2-expressing cells compared to vector control transfectants, consistent with the notion that BCL-2 may also regulate Ca^{2+} compartmentalization in mitochondria. More recently, Lam *et al.* (1994) have shown that overexpression of BCL-2 interferes with thapsigargin-induced Ca^{2+} mobilization from the ER in the WEHI7.2 T lymphoma cell line, an effect that is associated with preservation cell viability. Other work has shown that BCL-2 prevents the Ca^{2+} increase induced by hydrogen peroxide in B cells from BCL-2 transgenic mice (Zornig *et al.*, 1995). Finally, a screen for polypeptides capable of physically interacting with BCL-2 has identified a protein (Nip-2) containing a putative Ca^{2+} -binding

domain that colocalizes with BCL-2 in the ER and perinuclear region (Boyd *et al.*, 1994). Precisely how BCL-2 regulates intracellular Ca^{2+} is still unclear, although given its co-localization with Ca^{2+} transport sites in mitochondria, the ER, and the nuclear envelope (deJong *et al.*, 1994), a direct effect of BCL-2 on Ca^{2+} channel(s) is possible. Alternatively, given the tight interrelationship between Ca^{2+} and oxidative stress, BCL-2 could be influencing Ca^{2+} homeostasis via effects on cellular redox status (Hockenbery *et al.*, 1993; Kane *et al.*, 1993). It is also possible that BCL-2 regulates the localization and/or enzymatic activities of other polypeptides implicated in the control of apoptosis, especially nuclear proteins such as p53, Myc, AP-1 and the catalytic component of protein kinase A (Marti *et al.*, 1994).

Calcium Coupling to the Effector Pathway

An important aspect of ongoing research involves defining the biochemical consequences of Ca^{2+} mobilization in apoptotic cells, and at present there are two models to explain how these alterations might trigger apoptosis. In one, depletion of intracellular stores and possibly influx of Ca^{2+} across the plasma membrane promote a sustained Ca^{2+} increase that acts as a signal for apoptosis, perhaps in part by activating key catabolic enzymes that make up the effector machinery (Fig. 2). In the second, it is not the Ca^{2+} increase but the emptying of intracellular Ca^{2+} stores that triggers apoptosis, perhaps by disrupting intracellular architecture and allowing key elements of the effector machinery to gain access to their substrates. These models are certainly not mutually exclusive. Evidence for both models will be presented below, but it should be emphasized at the outset that definitive proof for either one is lacking at present.

Possible Targets for Ca^{2+} Elevations

Signal transduction intermediates

Activation of Ca^{2+} -dependent protein kinases and/or phosphatases leading to alterations in gene transcription represents one possible way that Ca^{2+} might regulate apoptosis. The most convincing support for this hypothesis has come from experiments with the immunosuppressant cyclosporin A, a compound that binds a family of cytosolic receptors termed cyclophilins and in so doing, forms a composite molecular surface that binds to and inhibits the Ca^{2+} /calmodulin-dependent protein serine/threonine phosphatase, calcineurin (Liu *et al.*, 1991). Studies by several independent laboratories have shown that cyclosporin A can block Ca^{2+} -dependent apoptosis in lymphoid model systems (Amendola *et al.*, 1994; Bonnefoy-Berard *et al.*, 1994; Makrigiannis *et al.*,

Calcium regulation of apoptosis




Figure 2. Targets for Ca^{2+} in the regulation of apoptosis. The figure schematically presents the processes involved in apoptosis for which a role for Ca^{2+} is strongly supported by recent investigation. PS: phosphatidylserine; PLA_2 : phospholipase A_2 ; NS: nuclear scaffold.

1994; Shi *et al.*, 1989), indicating that calcineurin activation may be required for these responses. Recent work (Jiang, Chow, Orrenius, manuscript submitted) confirms that cyclosporin A and FK506 block Ca^{2+} stimulated apoptosis in T cell hybridomas but they are without effect on Ca^{2+} -dependent apoptosis in immature rodent thymocytes *in vitro*. Induction of the orphan steroid receptor Nur77 and the Fas ligand represent at least two of the confirmed molecular targets of calcineurin in mature T cells and T cell hybridomas (Anel *et al.*, 1994; Yazdanbakhsh *et al.*, 1995). The involvement of calcineurin in Ca^{2+} stimulated apoptosis could also potentially explain, in part, the sensitivity of various apoptotic pathways to inhibition by calmodulin antagonists.

Phospholipase A_2 (PLA_2) is another signaling intermediate that has been suggested to be a target for Ca^{2+} in apoptosis. Stimulation of T cell clones with retroviral or bacterial superantigens is associated with the generation of reactive oxygen intermediates and PLA_2 activation (Wever *et al.*, 1995). Similarly, several groups have shown that tumor necrosis factor-induced apoptosis is associated with PLA_2 activation (Hayakawa *et al.*, 1993; Hollenbach *et al.*, 1992; Mutch *et al.*, 1992; Voelkel-Johnson *et al.*, 1995), and that the sensitivity of a large panel of melanoma cell lines to TNF plus inhibitors of macromolecular synthesis is associated with the level of cytosolic PLA_2 (Voelkel-Johnson *et al.*, 1995). Arachidonic acid release has been documented in several

other models of apoptosis (Agarwal *et al.*, 1993), most notably radiation-induced apoptosis in thymocytes (Korystov *et al.*, 1993; Shaposhnikova *et al.*, 1994), and inhibitors of PLA₂ block both arachidonic acid release and cell death (Korystov *et al.*, 1993). Radiation and TNF-induced apoptosis may actually be linked by PLA₂ activation, as irradiation triggers TNF α production via a PLA₂-dependent mechanism (Hallahan *et al.*, 1994). Direct effects of X-rays on PLA₂ activation in isolated membrane preparations have also been reported (Cohen and DeLeo, 1993). It has also been suggested that BCL-2 and BCL-X may inhibit TNF- and Fas-induced apoptosis in breast carcinoma cells via a mechanism that involves inhibition of PLA₂ activation (Jaattela *et al.*, 1995). Therefore, Ca²⁺ may act in these and other systems to promote PLA₂ activation, which may in turn result in oxidative stress and DNA damage via release and metabolism of arachidonic acid.

Ca²⁺-activated proteases

There is some evidence that Ca²⁺-sensitive protease(s) might represent direct targets for Ca²⁺ elevations in apoptosis. Recent work has shown that the Ca²⁺-dependent neutral protease, calpain, is rapidly activated in T lymphocytes following treatment with glucocorticoids or exposure to γ -irradiation, and that calpain antagonists can block DNA fragmentation associated with the response (Squier *et al.*, 1994). Similarly, Henkart's group has demonstrated that some (but not all) pathways of apoptosis in mature T lymphocytes can be inhibited by calpain antagonists (Sarin *et al.*, 1993; 1995). The cytoskeletal protein, fodrin, is at least one substrate for calpain that is cleaved in T cells following treatment with glucocorticoids or engagement of the Fas antigen (Martin *et al.*, 1995). Similarly, we have recently obtained evidence that the cytoskeletal protein, vimentin, is also cleaved in apoptotic cells by a calpain-sensitive mechanism (J. Kiefer *et al.*, unpublished observation). Precisely how fodrin and vimentin cleavage participate in the apoptosis effector mechanism is unclear, although they may be involved in cellular shrinkage, membrane blebbing, or other structural alterations associated with apoptosis.

Other Ca²⁺-activated proteases may also participate in the process. Previous work has demonstrated that incubation of isolated nuclei in the presence of Ca²⁺ promotes the rapid degradation of a family of nuclear matrix proteins, the lamins (Clawson *et al.*, 1992; Tokes and Clawson, 1989). The protease responsible for lamin cleavage in this system is directly associated with the nuclear matrix and is activated by Ca²⁺. In parallel, independent work from several laboratories has shown that lamins are also degraded in cells undergoing apoptosis (Kaufmann, 1989; Lazebnik *et al.*, 1993; Neamati *et al.*,

1995; Oberhammer *et al.*, 1994), and in thymocytes lamin cleavage occurs via a Ca²⁺-dependent mechanism (Neamati *et al.*, 1995), inspiring investigations into the possible involvement of the lamin protease in the response. Thus, we have found that a specific peptide inhibitor of the putative lamin protease (also known as "nuclear scaffold (NS) protease") blocks cellular shrinkage and DNA fragmentation in thymocytes exposed to antibodies to the T cell receptor, thapsigargin and glucocorticoids, but not in cells treated with etoposide, a cancer chemotherapeutic agent that acts via induction of DNA damage (Zhivotovsky *et al.*, 1995; McConkey, 1996). We have also found that inhibitors of this protease selectively block both lamin cleavage and DNA fragmentation in isolated nuclei incubated with Ca²⁺, while inhibitors of the ICE family of proteases do not (McConkey, 1996). Importantly, however, the NS protease is not the only protease that can cleave the lamins, as indicated by recent work by Lazebnik *et al.* (1995), who have shown that an inhibitor of the ICE/ced-3 family of cysteine proteases blocks lamin cleavage in another isolated nuclei system, whereas treatment with inhibitors of the NS protease does not. Moreover, in this system, lamin cleavage, chromatin condensation, and DNA fragmentation do not require exogenous Ca²⁺. Whether Ca²⁺ is involved in generating the apoptosis-promoting activity found within the extracts these authors use to promote nuclear lamin cleavage is not clear at present. In addition, whether BCL-2 is capable of blocking the nuclear alterations induced by these extracts is also not known, and it is therefore possible that the activity present in them is an irreversibly activated component of the effector machinery (i.e., an ICE protease) that is no longer subject to Ca²⁺ regulation.

Ca²⁺-activated endonuclease(s)

As introduced above, endonuclease activation resulting in the formation of oligonucleosome-length DNA fragments (DNA ladders) remains the most characteristic biochemical feature of apoptotic cell death. Early work by Hewish and Burgoyne (1973) and later by Vanderbilt *et al.* (1982) demonstrated that a Ca²⁺/Mg²⁺-dependent enzyme activity, capable of generating characteristic apoptotic chromatin cleavage patterns, is constitutively present within nuclei of a variety of different cell types. Subsequent work by Cohen and Duke (1984), and Wylie *et al.* (1984), demonstrated the involvement of this activity in the DNA fragmentation observed in thymocytes undergoing apoptosis, and it is now thought that it mediates DNA fragmentation in a variety of other model systems as well. The search for, and purification of potential Ca²⁺-dependent apoptotic nucleases has subsequently been undertaken by several laboratories. Thus, Gaido and Cidlowski (1991) have described a low-

molecular weight nuclease (NUC18) with Ca^{2+} and Mg^{2+} dependence activity in apoptotic lymphoid cells in response to several kinds of apoptotic stimuli. Interestingly, the purified NUC18 shares amino acid sequence homology with cyclophilin, and human recombinant cyclophilin A has biochemical and pharmacological properties identical to native NUC18 (Montague *et al.*, 1994). NUC18 is also present in untreated thymocytes in precursor form or as part of a higher molecular weight complex (> 100 kDa), suggesting that the nuclease dissociates in response to apoptotic signals. Although the precise mechanism of liberation of active enzyme from its precursors is unknown, an attractive possibility is that it may involve proteolysis.

The Ca^{2+} -dependent endonuclease DNase I is another excellent candidate apoptotic nuclease (Peitsch *et al.*, 1993). Addition of the enzyme to isolated nuclei and other reconstitution systems promotes the formation of DNA strand breaks that possess the same 5'- PO_4 and 3'-OH end groups found in DNA fragments isolated from apoptotic cells. Although the enzyme is localized within the rough endoplasmic reticulum, the Golgi complex, and small (secretory) vesicles in viable cells, it is also found within the perinuclear space of apoptotic cells, and it is possible that structural alterations in the ER and/or nuclear envelope associated with apoptosis may promote the entry of DNase I into the nucleus (see below). A similar mechanism may promote entry of an ER-localized fraction of the NS lamin protease into the nucleus. Several other proteins with $\text{Ca}^{2+}/\text{Mg}^{2+}$ endonuclease activity have been isolated (Ishida *et al.*, 1974; Nikonova *et al.*, 1993; Ribeiro and Carson, 1993; Wylie *et al.*, 1992), but to date proof that any one of these activities is directly involved in oligonucleosomal DNA fragmentation in apoptosis is lacking.

Transglutaminase activation

Transglutaminases are a group of Ca^{2+} -dependent enzymes that catalyze the post-translational coupling of amines (including polyamines) into proteins and the crosslinking of proteins via gamma glutamyl lysine bridges when the amine is a peptide-bound lysine residue. Tissue transglutaminase has been implicated in a number of physiological processes, including crosslinking of integral plasma membrane proteins with the cytoskeleton. Recent work indicates that tissue transglutaminase is also involved in induction of apoptosis (Fesus *et al.*, 1987, 1989). Expression of transglutaminase mRNA and protein levels increase markedly in dying cells. The enzyme appears to be activated by elevations of the cytosolic Ca^{2+} concentration, which are involved in apoptosis in many different systems. Isolation of apoptotic bodies from a number of different tissues has

shown that they are resistant to dissolution by detergents and chaotropic agents; this may in part be explained by the fact that surface polypeptides in these structures are crosslinked via gamma glutamyl lysine isopeptide bonds (Taresa *et al.*, 1992). The resistance of these structures to proteolysis may allow them to accumulate, and they can be detected in the media of cell cultures containing high rates of apoptotic cell death (Fesus *et al.*, 1991). Isodi-peptide can also be detected in normal plasma, and its concentration increases following induction of apoptosis in various organs, including the thymus and liver.

The role of transglutaminase promoting cell death and/or phagocytosis is still poorly understood. One possibility is that protein crosslinking stabilizes apoptotic cells and bodies, preventing leakage of intracellular contents into the extracellular milieu (which can trigger inflammation). Alternatively, transglutaminase modification may target proteins for subsequent degradation. Intriguingly, overexpression of the enzyme has been reported to trigger apoptotic cell death (Melino *et al.*, 1994), suggesting that transglutaminase may be a component of the death effector pathway. Further efforts are required to identify the substrates for transglutaminase in apoptotic cells and to determine the consequences of their modification.

Exposure of phosphatidylserine and macrophage recognition

Recent work indicates that the movement of phosphatidylserine (PS) from the inner to the outer surface of the plasma membrane, a process that functions in the removal of apoptotic cells and bodies by both professional phagocytic cells (macrophages) and neighboring cells in tissues (Fadok *et al.*, 1992a, 1992b), is another component of apoptosis that is regulated by alterations in cytosolic Ca^{2+} . Phospholipids are known to be distributed asymmetrically across the plasma membrane, with phosphatidylcholine and sphingomyelin localized primarily to the extracellular surface and PS and phosphatidylethanolamine restricted almost exclusively to the intracellular surface under normal conditions (Verkleij *et al.*, 1973; Zwaal *et al.*, 1975). Most of the work on plasma membrane lipid asymmetry has been conducted with red blood cells, where it is known that PS localization is regulated by energy-dependent processes involving specific lipid transporters ("flipases" and "flopases") that move PS to the outside or inside surface, respectively. Transport in both directions is ATP-dependent and sensitive to sulfhydryl modifying agents. Interestingly, inhibition of lipid movement with reagents that abrogate the activities of both the flipase and flopase does not result in loss of membrane asymmetry (Connor and Schroit, 1990), suggesting that these enzymes may primarily function to restore lipid asymmetry following its disruption.

tion. However, transport can also be inhibited by increasing the cytosolic Ca^{2+} concentration, which does result in rapid non-specific redistribution of all phospholipids (Beyers *et al.*, 1990). The mechanism of Ca^{2+} -mediated PS exposure is not yet clear, but in red blood cells, it closely parallels formation of cytoskeleton-free lipid microvesicles (Sims *et al.*, 1989) and other phenomena, such as, calpain activation (Fox *et al.*, 1991) and the formation of phosphatidylinositol(4,5)-bisphosphate- Ca^{2+} complexes (Sulpice *et al.*, 1994). However, neither direct Ca^{2+} effects, calpain-mediated protein degradation (Basse *et al.*, 1993; Orrenius *et al.*, manuscript submitted), $\text{Ins}(4,5)\text{P}_2$ accumulation (Beyers *et al.*, 1995), nor inhibition of the flippase can singularly accommodate the membrane rearrangements that occur. More recent work suggests that Ca^{2+} -mediated redistribution is mediated by a Ca^{2+} and sulfhydryl-sensitive, energy-dependent lipid scramblase (Williamson *et al.*, 1995). Therefore, elevations in the cytosolic Ca^{2+} concentration probably promote PS exposure and allow for macrophage recognition primarily by inactivating the PS translocase and by activating the scramblase (Verhoven *et al.*, 1995). Interestingly, PS exposure on aged red blood cells is associated with increased cytosolic Ca^{2+} levels and increased cell density, suggesting that these events may be mechanistically related in both red cells and apoptotic cells. In addition, given its regulation by Ca^{2+} and calpain-dependent mechanisms, it is tempting to speculate that red blood cell microvesiculation may be functionally related to the formation of apoptotic bodies by nucleated cells undergoing apoptosis.

Plasma membrane lipid asymmetry is critically involved in several other important physiological functions. For example, surface-exposed PS serves as the point of assembly for the coagulation factors Va and Xa into the prothrombinase complex (Rosing *et al.*, 1985). In addition, PS exposure enhances membrane fusion events and appears to be involved in the initiation of microvesiculation in red blood cells (Schewe *et al.*, 1992). Finally, surface PS is also detectable on certain tumor cells (Utsugi *et al.*, 1991). Although the mechanisms underlying the latter have not been defined, it is possible that PS exposure is involved in the prominent macrophage infiltration observed in most solid tumors and that surface PS may represent a potential target for anti-tumor therapies.

Possible Consequences of Intracellular Ca^{2+} Pool Depletion

In some cellular systems, extracellular or intracellular Ca^{2+} chelators can actually promote DNA fragmentation (Treves *et al.*, 1994), even though other triggers of apoptosis in these systems (i.e., glucocorticoids,

growth factor withdrawal) have been shown to deplete the ER Ca^{2+} store. These observations have led Baffy *et al.* (1993) and more recently Lam *et al.* (1993) to propose that depletion of the ER Ca^{2+} store may itself serve as a signal for apoptosis. How could this occur? At least two of the catabolic enzymes proposed to be involved in the effector mechanism of apoptosis (DNase I and an extranuclear pool of the NS protease) are localized to the ER, and it is therefore possible that loss of Ca^{2+} leads to release of these factors into the perinuclear region or into the nuclear matrix itself. In addition, it is known that ER Ca^{2+} pool depletion results in the release of a small biomolecule that participates in a retrograde signal for plasma membrane Ca^{2+} influx, and it is possible that it or another molecule released in a similar fashion can also promote cell death.

Depletion of mitochondrial Ca^{2+} stores may also participate in the signal for apoptosis. Mitochondrial Ca^{2+} uptake is driven by mitochondrial membrane potential ($\Delta\Psi$) (Richter, 1993). In de-energized mitochondria, Ca^{2+} can be released by a reversal of the uptake pathway. Under conditions of oxidative stress, mitochondrial Ca^{2+} cycling can reach critical levels, leading to increased energy expenditure and a dramatic fall in $\Delta\Psi$. Recent work has shown that a fall in mitochondrial $\Delta\Psi$ is an early event in apoptosis (Petit *et al.*, 1995; Zamzani *et al.*, 1995a, 1995b), and ruthenium red, an inhibitor of the mitochondrial Ca^{2+} uptake pathway, blocks apoptosis in L929 fibroblasts (Hennet *et al.*, 1993) and inhibits the progression of apoptosis in glucocorticoid-treated splenocytes (Zamzani *et al.*, 1995a), suggesting that mitochondrial Ca^{2+} release is involved. Again, further efforts are required to determine the relationship between this event and the activation of the proteases and nucleases of the effector pathway.

Alternative Signals for Apoptosis

Oxidative stress

Several lines of evidence indicate that reactive oxygen species are involved in promoting apoptosis in diverse model systems. Treatment of cells with low to moderate doses of exogenous oxidants (i.e., hydrogen peroxide, tert-butyl peroxide, menadione) can trigger apoptosis (Buttke and Sandstrom, 1994; Hennet *et al.*, 1993; Hockenbery *et al.*, 1993; Kane *et al.*, 1993; Lennon *et al.*, 1990, 1991; McConkey *et al.*, 1988). Moreover, apoptosis induced by agents that are not direct oxidants (TNF, glucocorticoids, thapsigargin, chemotherapeutic agents) is associated with oxygen radical production and depletion of intracellular antioxidants (i.e., reduced glutathione) (Fernandez *et al.*, 1995; Hennet *et al.*, 1993; Hockenbery *et al.*, 1993; Kane *et al.*, 1993; Mayer and Noble, 1994; Richter, 1993; Slater *et al.*,

Calcium regulation of apoptosis




Figure 3. Candidate downstream mediators of ceramide-induced apoptosis. Production of ceramide via sphingomyelinase activation or ceramide synthase stimulation (daunorubicin) can lead to elevations in ceramide levels that can trigger cell death. Signaling intermediates known to be directly regulated by ceramide include ceramide-activated protein phosphatase (CAPP) and ceramide-activated protein kinase (CAPK). More recent work suggests that the former can promote the dephosphorylation and activation of the p110 retinoblastoma protein, resulting in cell cycle arrest (Dbaibo *et al.*, 1995), whereas the latter has been shown to interact with and regulate the protein serine/threonine kinase Raf-1 (Yao *et al.*, 1995). We (Trent *et al.*, 1996) and others (Gulbins *et al.*, 1995) have obtained evidence that activation of Ras is also involved, and preliminary work suggests that the ζ isoform of protein kinase C (PKC ζ) represents another target for ceramide (Lozano *et al.*, 1994). Importantly, diacylglycerol and phorbol esters can block ceramide-induced apoptosis, presumably by activating one or more PKC isozymes. How cell cycle arrest and apoptosis are differentially regulated by ceramide is not known at present.

1995; Wolfe *et al.*, 1994; Zamzani *et al.*, 1995a). Although the identities of the oxygen radicals involved in each system are still under active investigation, roles for superoxide (Hennet *et al.*, 1993; Zamzani *et al.*, 1995b), lipid peroxides (Hockenberry *et al.*, 1993), nitric oxide (Xie *et al.*, 1995), and hydroxyl radicals (Wolfe *et al.*, 1994) have been proposed. In these systems, exogenous antioxidants such as N-acetyl cysteine and free radical scavengers block DNA fragmentation and cell death (Fernandez *et al.*, 1995; Mayer and Noble, 1994; Slater *et al.*, 1995; Wolfe *et al.*, 1994). Finally, several laboratories have now presented strong evidence that BCL-2 (Hockenberry *et al.*, 1993; Kane *et al.*, 1993) and BCL-XL (Fang *et al.*, 1995) possess antioxidant properties that may be involved in their abilities to inhibit cell death. Thus, oxidative stress is another good candidate for a central cell death signal, one which may affect intracellular Ca^{2+} homeostasis. However, it should be noted that BCL-2 and BCL-XL can still inhibit apoptosis under conditions of low oxygen (Jacobson and Raff, 1995; Shimizu *et al.*, 1995), which has been raised as an argument against a universal role for oxidative stress in the response.

Intracellular acidification

Eastman's group (Barry and Eastman, 1993) have proposed another alternative to Ca^{2+} that may serve as a general signal for cell death. In their efforts to characterize the biochemical mechanisms underlying chemotherapy-induced apoptosis, they determined that isolated nuclei from their cellular models possessed an endonuclease activity that was stimulated by acidic pH (Dnase II) (Barry and Eastman, 1993). Interestingly, Mary Collins and her co-workers (personal communication) have recently identified an acidic nuclease in IL-3-dependent hematopoietic cells that can also be activated by Ca^{2+} under the appropriate conditions, suggesting that intracellular acidification and alterations in intracellular Ca^{2+} homeostasis may represent independent ways of arriving at the same endpoint (endonuclease activation) in apoptotic cells. Other efforts have shown that a drop in cytoplasmic pH precedes the morphological and biochemical features of apoptosis in certain models (Barry and Eastman, 1992; Li and Eastman, 1995), observations that have since been confirmed by other investigators in other systems (Gottlieb *et al.*, 1995). Moreover, the protective effects of certain survival factors and agents that activate protein kinase C have been linked to activation of the Na^+/H^+ antiporter and intracellular alkalinization (Gottlieb *et al.*, 1995; Rajotte *et al.*, 1992). Finally, the drop in pH is prevented by overexpression of BCL-2 (A. Eastman, personal communication), suggesting that acidification may represent another cellular target for this family of apoptosis suppressors.

It will be interesting to determine whether intracellular acidification is related to oxidative stress and intracellular Ca^{2+} alterations in these systems.

Ceramide production

The generation of bioactive signal transduction regulators via the hydrolysis of plasma membrane phospholipids is emerging as an important general means of regulating apoptotic cell death. Of all of the second messengers shown to be involved, most recent attention has been focussed on ceramide as a possible ubiquitous trigger of apoptosis (Obeid *et al.*, 1993; Pushkareva *et al.*, 1995) (Fig. 3). Two major mechanisms have been identified that appear to contribute to the formation of ceramide under different circumstances. The most common pathway involves activation of the enzyme sphingomyelinase, which catalyzes the hydrolysis of sphingomyelin to form ceramide and diacylglycerol, a response that is involved in apoptosis induced by tumor necrosis factor, engagement of the Fas antigen, and ionizing radiation (Cifone *et al.*, 1994; Gulbins *et al.*, 1995; Haimovitz-Friedman *et al.*, 1994; Jarvis *et al.*, 1994b; Obeid *et al.*, 1993; Pushkareva *et al.*, 1995; Tepper *et al.*, 1995). Ceramide can also be formed *de novo* via activation of ceramide synthase, and it has been shown that the cytotoxic effects of the chemotherapeutic agent daunorubicin on leukemic cell lines are mediated by this pathway (Bose *et al.*, 1995). Notably, exogenous hydrolysis-resistant ceramide analogs or sphingomyelinase can mimic the effects of TNF and the other apoptosis-inducing agents to trigger endonuclease activation and cell death, indicating that ceramide production is sufficient to induce apoptosis (Bose *et al.*, 1995; Cifone *et al.*, 1994; Gulbins *et al.*, 1995; Haimovitz-Friedman *et al.*, 1994; Jarvis *et al.*, 1994a, 1994b; Obeid *et al.*, 1993; Tepper *et al.*, 1995). In addition, the possibility that ceramide is an evolutionarily conserved trigger of apoptosis is suggested by the fact that overexpression of the *Drosophila* cell death protein, Reaper, leads to upregulation of cellular ceramide levels (Pronk *et al.*, 1996). Interestingly, phorbol esters and diacylglycerol (DAG) antagonize the death-promoting effects of ceramide in diverse models (Jarvis *et al.*, 1994a; Obeid *et al.*, 1993), suggesting that a balance between ceramide and DAG may determine the outcome of death signals. The downstream targets for ceramide remain unclear, but candidates include a ceramide-dependent protein phosphatase (CAPP) (Wolff *et al.*, 1994), a ceramide-activated protein kinase (CAPK) (Kolesnick and Golde, 1994), the ζ isoform of protein kinase C (Lozano *et al.*, 1994), and the H-ras protooncogene (Gulbins *et al.*, 1995; Trent *et al.*, 1996). Whether ceramide acts upstream, downstream, or independently of the other candidate biochemical mediators of apoptosis remains to be determined.

Conclusions and Future Directions

The independent efforts of many laboratories over the past several years have established that alterations in intracellular Ca^{2+} homeostasis are commonly involved in initiating apoptosis. Ongoing work suggests that activation of Ca^{2+} stimulated signalling networks and catabolic enzymes represents one way these signals are translated into responses. In addition, preliminary evidence indicates that the depletion of intracellular Ca^{2+} pools can itself serve as a signal for cell death, perhaps by promoting relocalization of some of the key catabolic enzymes involved and by enhancing oxidative stress in mitochondria. In some systems, other signals have been defended as key regulators of the response, and further efforts are therefore required to determine if and how all of these signals are interrelated.

An important component of the defense of Ca^{2+} as a central cell death regulator is the idea that BCL-2 and its homologs specifically regulate intracellular Ca^{2+} compartmentalization, an idea that has strong preliminary support but requires a good deal of additional investigation. In particular, further efforts are required to determine how BCL-2 exerts its effects on Ca^{2+} and whether these effects are required for its cell death-suppressing function. Similarly, it is maintained at present that any central cell death signal would directly or indirectly activate one or more members of the ICE family of cysteine proteases, and at present there is no evidence available that any of the candidate central cell death signals are capable of this. Elucidation of these relationships over the next few years should provide new targets for therapeutic intervention that may aid in the treatment of the expanding number of diseases, including cancer, AIDS, and neurodegenerative diseases, in which apoptosis is thought to play a central role in their pathologies.

References

- Agarwal ML, Larkin HE, Zaidi SI, Mukhtar H, Oleinick NL (1993) Phospholipase activation triggers apoptosis in photosensitized mouse lymphoma cells. *Cancer Res* **53**, 5897-5902.
- Allbritton NL, Verret CR, Wolley RC, Eisen HN (1988) Calcium ion concentrations and DNA fragmentation in target cell destruction by murine cloned cytotoxic T lymphocytes. *J Exp Med* **167**, 514-527.
- Amendola A, Lombardi G, Oliverio S, Colizzi V, Piacentini M (1994) HIV-1 gp120-dependent induction of apoptosis in antigen-specific human T cell clones is characterized by "tissue" transglutaminase expression and prevented by cyclosporin A. *FEBS Lett* **339**, 258-264.
- Anel A, Buferne M, Boyer C, Schmitt-Verhulst AM, Golstein P (1994) T cell receptor-induced Fas ligand expression in cytotoxic T lymphocyte clones is blocked by protein tyrosine kinase inhibitors and cyclosporin A. *Eur J Immunol* **24**, 2469-2476.
- Aw TY, Nicotera P, Manzo L, Orrenius S (1990) Tributyltin stimulates apoptosis in rat thymocytes. *Arch Biochem Biophys* **283**, 46-50.
- Baffy G, Miyashita T, Williamson JR, Reed JC (1993) Apoptosis induced by withdrawal of interleukin-3 (IL-3) from an IL-3-dependent hematopoietic cell line is associated with repartitioning of intracellular calcium and is blocked by enforced BCL-2 oncoprotein production. *J Biol Chem* **268**, 6511-6519.
- Barry MA, Eastman A (1992) Endonuclease activation during apoptosis: The role of cytosolic Ca^{2+} and pH. *Biochem Biophys Res Comm* **186**, 782-789.
- Barry MA, Eastman A (1993) Identification of deoxyribonuclease II as an endonuclease involved in apoptosis. *Arch Biochem Biophys* **300**, 440-450.
- Basse F, Gaffet P, Rendu F, Bienvenue A (1993) Translocation of spin-labeled phospholipids through the plasma membrane during thrombin- and ionophore A23187-induced platelet aggregation. *Biochemistry* **32**, 2337-2344.
- Bellomo G, Perotti M, Taddei F, Mirabelli F, Finardi G, Nicotera P, Orrenius S (1992) Tumor necrosis factor α induces apoptosis in mammary adenocarcinoma cells by an increase in intranuclear free Ca^{2+} concentration and DNA fragmentation. *Cancer Res* **52**, 1342-1346.
- Bevers EM, Verahllen PF, Visser AJ, Comfurius P, Zwaal RF (1990) Bidirectional transbilayer lipid movement in human platelets as visualized by the fluorescent membrane probe 1[4-(trimethylammonio)phenyl]-6-phenyl-1,3,5-hexatriene. *Biochemistry* **29**, 5132-5137.
- Bevers EM, Wiedmer T, Comfurius P, Zhao J, Smeets EF, Schlegel RA, Schroit AJ, Weiss HJ, Williamson P, Zwaal RF, Sims PJ (1995) The complex of phosphatidylinositol 4,5-bisphosphate and calcium ions is not responsible for Ca^{2+} -induced loss of phospholipid asymmetry in the human erythrocyte: A study of Scott syndrome, a disorder of calcium-induced phospholipid scrambling. *Blood* **86**, 1983-1991.
- Bonnefoy-Berard N, Genestier L, Flacher M, Revillard JP (1994) The phosphoprotein phosphatase calcineurin controls calcium-dependent apoptosis in B cell lines. *Eur J Immunol* **24**, 325-329.
- Bose R, Verheij M, Haimovitz-Friedman A, Scotto K, Fuks Z, Kolesnick R (1995) Ceramide synthase mediates daunorubicin-induced apoptosis: An alternative mechanism for generating death signals. *Cell* **82**, 405-414.
- Boyd JM, Malstrom S, Subramanian T, Venkatesh LK, Schaeper U, Elangovan B, D'Sa-Eipper C, Chinn-

durai G (1994) Adenovirus E1B 19 kDa and BCL-2 proteins interact with a common set of cellular proteins. *Cell* **79**, 341-351.

Brown DG, Sun XM, Cohen GM (1993) Dexamethasone-induced apoptosis involves cleavage of DNA to large fragments prior to internucleosomal fragmentation. *J Biol Chem* **268**, 3037-3039.

Bump NJ, Hackett M, Hugunin M, Seshagiri S, Brady K, Chen P, Ferez C, Franklin S, Ghayur T, Li P, Licari P, Mankovich M, Shi L, Greenberg AH, Miller LK, Wong WW (1995) Inhibition of ICE family proteases by baculovirus antiapoptotic protein p35. *Science* **269**, 1885-1888.

Buttke TM, Sandstrom PA (1994) Oxidative stress as a mediator of apoptosis. *Immunol Today* **15**, 7-10.

Choi MS, Boise LH, Gottschalk AR, Quintans J, Thompson CB (1995) The role of BCL-XL in CD40-mediated rescue from anti- μ -induced apoptosis in WEHI-231 B lymphoma cells. *Eur J Immunol* **25**, 1352-1357.

Cifone MG, Maria RD, Roncaioli P, Rippon RM, Azuma M, Lanier LL, Santoni A, Testi R (1994) Apoptotic signaling through CD95 (Fas/Apo-1) activates acidic sphingomyelinase. *J Exp Med* **180**, 1547-1552.

Clawson GA, Norbeck LL, Hatem CL, Rhodes C, Amiri P, McKerrow JH, Patierno SR, Fiskum G (1992) Ca^{2+} regulated serine protease associated with the nuclear scaffold. *Cell Growth Differentiation* **3**, 827-838.

Cohen D, DeLeo VA (1993) Ultraviolet radiation-induced phospholipase A2 activation occurs in mammalian cell membrane preparations. *Photochem Photobiol* **57**, 383-390.

Cohen JJ, Duke RC (1984) Glucocorticoid activation of a calcium-dependent endonuclease in thymocyte nuclei leads to cell death. *J Immunol* **132**, 38-42.

Connor J, Schroit AJ (1990) Aminophospholipid translocation in erythrocytes: Evidence for the involvement of a specific transporter and an endofacial protein. *Biochemistry* **29**, 37-43.

Dbaibo GS, Pushkareva MY, Jayadev S, Schwarz JK, Horowitz JM, Obeid LM, Hannun YA (1995) Retinoblastoma gene product as a downstream target for a ceramide-dependent pathway of growth arrest. *Proc Natl Acad Sci USA* **92**, 1347-1351.

deJong D, Prins FA, Mason DY, Reed JC, Ommen GBv, Kluin PM (1994) Subcellular localization of the bcl-2 protein in malignant and normal lymphoid cells. *Cancer Res* **54**, 256-260.

Dowd DR, MacDonald PN, Komm BS, Maussler MR, Miesfeld R (1991) Evidence for early induction of calmodulin gene expression in lymphocytes undergoing glucocorticoid-mediated apoptosis. *J Biol Chem* **266**, 18423-18426.

Dypbukt JM, Ankarcona M, Burkitt M, Sjöholm A, Strom K, Orrenius S, Nicotera P (1994) Different

pro-oxidant levels stimulate growth, trigger apoptosis, or produce necrosis of insulin-secreting RINm5F cells. The role of intracellular polyamines. *J Biol Chem* **269**, 30553-30560.

Fadok VA, Voelker DR, Campbell PA, Cohen JJ, Bratton DL, Henson PM (1992a) Exposure of phosphatidylserine on the surface of apoptotic lymphocytes triggers specific recognition and removal by macrophages. *J Immunol* **148**, 2207-2216.

Fadok VA, Savill JS, Haslett C, Bratton DL, Doherty DE, Campbell PA, Henson PM (1992b) Different populations of macrophages use either the vitronectin receptor or the phosphatidylserine receptor to recognize and remove apoptotic cells. *J Immunol* **149**, 4029-4035.

Fang W, Rivard JJ, Ganser JA, LeBien TW, Nath KA, Mueller DL, Behrens TW (1995) BCL-xL rescues WEHI 231 B lymphocytes from oxidant mediated death following diverse apoptotic stimuli. *J Immunol* **155**, 66-75.

Fernandes-Alnemri T, Litwack G, Alnemri ES (1994) CPP32, a novel human apoptotic protein with homology to *Caenorhabditis elegans* cell death protein ced-3 and mammalian interleukin-1 β -converting enzyme. *J Biol Chem* **269**, 30761-30764.

Fernandes-Alnemri T, Litwack G, Alnemri ES (1995) Mch2, a new member of the apoptotic Ced-3/Ice cysteine protease gene family. *Cancer Res* **55**, 2737-2742.

Fernandez A, Kiefer J, Fosdick L, McConkey DJ (1995) Oxygen radical production and thiol depletion are required for Ca^{2+} -mediated endogenous endonuclease activation in apoptotic thymocytes. *J Immunol* **155**, 5133-5139.

Fesus L, Thomazy V, Falus A (1987) Induction and activation of tissue transglutaminase during programmed cell death. *FEBS Lett* **224**, 104-108.

Fesus L, Thomazy V, Autuori F, Ceru MP, Tarcsa E, Piacentini M (1989) Apoptotic hepatocytes become insoluble in detergents and chaotropic agents as a result of transglutaminase action. *FEBS Lett* **245**, 150-154.

Fesus L, Tarcsa E, Kedei N, Autuori F, Piacentini M (1991) Degradation of cells dying by apoptosis leads to accumulation of epsilon (gamma-glutamyl) lysine isodipeptide in culture fluid and blood. *FEBS Lett* **284**, 109-112.

Filipski J, Leblanc J, Youdale T, Sikorska M, Walker PR (1990) Periodicity of DNA folding in higher order chromatin structures. *EMBO J* **9**, 1319-1327.

Fox JE, Austin CD, Reynolds CC, Steffen PK (1991) Evidence that agonist-induced activation of calpain causes the shedding of procoagulant-containing microvesicles from the membrane of aggregating platelets. *J Biol Chem* **266**, 13289-13295.

Furuya Y, Isaacs JT (1993) Differential gene regu-

lation during programmed death (apoptosis) versus proliferation of prostatic glandular cells induced by androgen manipulation. *Endocrinology* **133**, 2660-2666.

Gaido ML, Cidlowski JA (1991) Identification, purification, and characterization of a calcium-dependent endonuclease (NUC 18) from apoptotic rat thymocytes. *J Biol Chem* **266**, 18580-18585.

Gottlieb RA, Giesing HA, Zhu JY, Engler RL, Babior BM (1995) Cell acidification in apoptosis: Granulocyte colony-stimulating factor delays programmed cell death in neutrophils by up-regulating the vacuolar H⁺-ATPase. *Proc Natl Acad Sci USA* **92**, 5965-5968.

Gulbins E, Bissonnette R, Mahboubi A, Martin S, Nishioka W, Brunner T, Baier G, Baier-Bitterilch G, Byrd C, Lang F, Kolesnick R, Altman A, Green D (1995) FAS-induced apoptosis is mediated via a ceramide-initiated RAS signaling pathway. *Immunity* **2**, 341-351.

Haimovitz-Friedman A, Kan CC, Ehleiter D, Persaud RS, McLoughlin M, Fuks Z, Kolesnick RN (1994) Ionizing radiation acts on cellular membranes to generate ceramide and initiate apoptosis. *J Exp Med* **180**, 525-535.

Hallahan DE, Virudachalam S, Kuchibhotla J, Kufe DW, Weichselbaum RR (1994) Membrane-derived second messenger regulates X-ray-mediated tumor necrosis factor alpha gene induction. *Proc Natl Acad Sci USA* **91**, 4897-4901.

Hayakawa M, Ishida N, Takeuchi K, Shibamoto S, Hori T, Oku N, Ito F, Tsujimoto M (1993) Arachidonic acid-selective cytosolic phospholipase A2 is crucial in the cytotoxic action of tumor necrosis factor. *J Biol Chem* **268**, 11290-11295.

Hengartner MO, Horvitz HR (1994) *C elegans* cell death gene *ced-9* encodes a functional homolog of mammalian proto-oncogene *bcl-2*. *Cell* **76**, 665-676.

Hengartner MO, Ellis RE, Horvitz HR (1992) *Caenorhabditis elegans* gene *ced-9* protects cells from programmed cell death. *Nature* **356**, 494-499.

Hennet T, Richter C, Peterhans E (1993) Tumor necrosis factor-alpha induces superoxide anion production in mitochondria of L929 cells. *Biochem J* **289**, 587-592.

Hewish DR, Burgoyne LA (1973) Chromatin substructure. The digestion of chromatin DNA at regularly spaced sites by a nuclear deoxyribonuclease. *Biochem. Biophys Res Comm* **52**, 504-510.

Hockenbery DM, Oltvai ZN, Yin XM, Millman CL, Korsmeyer SJ (1993) BCL-2 functions in an antioxidant pathway to prevent apoptosis. *Cell* **75**, 241-251.

Hollenbach PW, Zilli DL, Laster SM (1992) Inhibitors of transcription and translation act synergistically with tumor necrosis factor to cause the activation of phospholipase A2. *J Biol Chem* **267**, 39-42.

Ishida R, Akiyoshi H, Takahashi T (1974) Isolation and purification of calcium and magnesium dependent endonuclease from rat liver nuclei. *Biochem Biophys Res Comm* **56**, 703-708.

Jaattela M, Benedict M, Tewari M, Shayman JA, Dixit VM (1995) BCL-X and BCL-2 inhibit TNF and Fas-induced apoptosis and activation of phospholipase A2 in breast carcinoma cells. *Oncogene* **10**, 2297-2305.

Jacobson MD, Raff MC (1995) Programmed cell death and Bcl-2 protection in very low oxygen. *Nature* **374**, 814-816.

Jacobson MD, Burne JF, Raff MC (1994) Programmed cell death and BCL-2 protection in the absence of a nucleus. *EMBO J* **13**, 1899-1910.

Jarvis WD, Jarvis FA, Fornari J, Browning JL, Gewirtz DA, Kolesnick RN, Grant S (1994a) Attenuation of ceramide-induced apoptosis by diglyceride in human myeloid leukemia cells. *J Biol Chem* **269**, 31685-31692.

Jarvis WD, Kolesnick RN, Fornari FA, Traylor RS, Gewirtz DA, Grant S (1994b) Induction of apoptotic DNA damage and cell death by activation of the sphingomyelin pathway. *Proc Natl Acad Sci USA* **91**, 73-77.

Jiang S, Chow SC, Nicotera P, Orrenius S (1994) Intracellular Ca²⁺ signals activate apoptosis in thymocytes: Studies using the Ca²⁺ ATPase inhibitor thapsigargin. *Exp Cell Res* **212**, 84-92.

Juntti-Berggren L, Larsson O, Rorsman P, Ammala C, Bokvist K, Wahlander K, Nicotera P, Dybukt JM, Orrenius S, Hallberg A, Berggren P (1993) Increased activity of L-type Ca²⁺ channels exposed to serum from patients with type I diabetes. *Science* **261**, 86-90.

Kaiser N, Edelman IS (1977) Calcium dependence of glucocorticoid-induced lymphocytolysis. *Proc Natl Acad Sci USA* **74**, 638-642.

Kaiser N, Edelman IS (1978) Further studies on the role of calcium in glucocorticoid-induced lymphocytolysis. *Endocrinology* **103**, 936-942.

Kane DJ, Sarafian TA, Anton R, Hahn H, Gralla EB, Valentine JS, Ord T, Bredesen DE (1993) BCL-2 inhibition of neural death: Decreased generation of reactive oxygen species. *Science* **262**, 1274-1277.

Kaneko Y, Tsukamoto A (1994) Thapsigargin-induced persistent intracellular calcium pool depletion and apoptosis in human hepatoma cells. *Cancer Lett* **79**, 147-155.

Kaufmann SH (1989) Induction of endonucleolytic DNA cleavage in human acute myelogenous leukemia cells by etoposide, camptothecin, and other cytotoxic anticancer drugs: A cautionary note. *Cancer Res* **49**, 5870-5878.

Kaufmann SH, Desnoyers S, Ottaviano Y, Davidson NE, Poirier GG (1993) Specific proteolytic cleavage of poly (ADP-ribose) polymerase: An early marker of che-

- motherapy-induced apoptosis. *Cancer Res* **53**, 3976-3985.
- Khan AA, Soloski MJ, Sharp AH, Schilling G, Sabatini DM, Li SH, Ross CA, Snyder SH (1996) Lymphocyte apoptosis: Mediation by increased type 3 nitro 1,4,5-trisphosphate receptor. *Science* **273**, 503-508.
- Kolesnick R, Golde DW (1994) The sphingomyelin pathway in tumor necrosis factor and interleukin-1 signaling. *Cell* **77**, 325-328.
- Korystov YN, Shaposhnikova VV, Dobrovinskaya OR, Eidus LK (1993) Intercellular interactions in the interphase death of irradiated thymocytes. *Radiat Res* **134**, 301-306.
- Kumar S, Kinoshita M, Noda M, Copeland NG, Jenkins NA (1994) Induction of apoptosis by the mouse Nedd2 gene, which encodes a protein similar to the product of *Caenorhabditis elegans* cell death gene ced-3 and the mammalian IL-1 β -converting enzyme. *Genes Dev* **8**, 1613-1626.
- Lam M, Dubyak G, Distelhorst CW (1993) Effect of glucocorticoid treatment on intracellular calcium homeostasis in mouse lymphoma cells. *Mol Endocrinol* **7**, 686-693.
- Lam M, Dubyak G, Chen L, Nunez G, Miesfeld RL, Distelhorst CW (1994) Evidence that bcl-2 represses apoptosis by regulating endoplasmic reticulum-associated Ca²⁺ fluxes. *Proc Natl Acad Sci USA* **91**.
- Lazebnik YA, Cole S, Cooke CA, Nelson WG, Earnshaw WC (1993) Nuclear events of apoptosis *in vitro* in cell-free mitotic extracts: A model system for analysis of the active phase of apoptosis. *J Cell Biol* **123**, 7-22.
- Lazebnik YA, Takahashi A, Moir RD, Goldman RD, Poirier GG, Kaufmann SH, Earnshaw WC (1995) Studies of the lamin proteinase reveal multiple parallel biochemical pathways during apoptotic execution. *Proc Natl Acad Sci USA* **92**, 9042-9046.
- Lennon SV, Martin SJ, Cotter TG (1990) Induction of apoptosis (programmed cell death) in tumour cells by widely diverging stimuli. *Biochem Soc Trans* **18**, 343-345.
- Lennon SV, Martin SJ, Cotter TG (1991) Dose-dependent induction of apoptosis in human tumour cell lines by widely divergent stimuli. *Cell Prolif* **24**, 203-204.
- Levick V, Coffey H, D'Mello SR (1995) Opposing effects of thapsigargin on the survival of developing cerebellar granule neurons in culture. *Brain Res* **676**, 325-335.
- Li J, Eastman A (1995) Apoptosis in an interleukin-2-dependent cytotoxic T lymphocyte cell line is associated with intracellular acidification. *J Biol Chem* **270**, 3203-3211.
- Liu J, Liu JD, Farmer J, Lane WS, Friedman J, Weissman I, Schreiber SL (1991) Calcineurin is a common target of cyclophilin-cyclosporin A FKBP-FK506 complexes. *Cell* **66**, 807-815.
- Lozano J, Berra E, Municio MM, Diaz-Meco MT, Dominguez I, Sanz L, Moscat J (1994) Protein kinase C zeta isoform is critical for kB-dependent promoter activation by sphingomyelinase. *J Biol Chem* **269**, 19200-19202.
- Makrigiannis AP, Blay J, Hoskin DW (1994) Cyclosporin A inhibits 2-chloroadenosine-induced DNA cleavage in mouse thymocytes. *Int J Immunopharmacol* **16**, 995-1001.
- Marin MC, Fernandez A, Bick RJ, Brisban S, Buja M, Snuggs M, McConkey DJ, von Eschenbach AC, Keating MJ, McDonnell TJ (1996) Apoptosis suppression by BCL-2 is correlated with the regulation of nuclear and cytosolic Ca²⁺. *Oncogene* **12**, 2259-2266.
- Marti A, Jehn B, Costello E, Keon N, Ke G, Martin F, Jaggi R (1994) Protein kinase A and AP-1 (c-Fos/JunD) are induced during apoptosis of mouse mammary epithelial cells. *Oncogene* **9**, 1213-1223.
- Martikainen P, Isaacs J (1990) Role of calcium in the programmed cell death of rat ventral prostatic glandular cells. *Prostate* **17**, 175-187.
- Martin SJ, Green DR (1995) Protease activation during apoptosis: Death by a thousand cuts? *Cell* **82**, 349-352.
- Martin SJ, O'Brien GA, Nishioka WK, McGahon AJ, Mahboubi A, Saido TC, Green DR (1995) Proteolysis of fodrin (non-erythroid spectrin) during apoptosis. *J Biol Chem* **270**, 6425-6428.
- Mayer M, Noble M (1994) N-acetyl-L-cysteine is a pluripotent protector against cell death and enhancer of trophic factor-mediated cell survival *in vitro*. *Proc Natl Acad Sci USA* **91**, 7496-7500.
- McConkey DJ (1996) Calcium-dependent, interleukin 1 β -converting enzyme (ICE) inhibitor-insensitive degradation of lamin B, and DNA fragmentation in isolated thymocyte nuclei. *J. Biol. Chem.* in press.
- McConkey DJ, Hartzell P, Nicotera P, Wyllie AH, Orrenius S (1988) Stimulation of endogenous endonuclease activity in hepatocytes exposed to oxidative stress. *Toxicol Lett* **42**, 123-130.
- McConkey DJ, Hartzell P, Amador-Perez JF, Orrenius S, Jondal M (1989a) Calcium-dependent killing of immature thymocytes by stimulation via the CD3/T cell receptor complex. *J Immunol* **143**, 1801-1806.
- McConkey DJ, Nicotera P, Hartzell P, Bellomo G, Wyllie AH, Orrenius S (1989b) Glucocorticoids activate a suicide process in thymocytes through an elevation of cytosolic Ca²⁺ concentration. *Arch Biochem Biophys* **269**, 365-370.
- McConkey DJ, Chow SC, Orrenius S, Jondal M (1990) NK cell-induced cytotoxicity is dependent on a

- Ca²⁺ increase in the target. *FASEB J* **4**, 2661-2664.
- McConkey DJ, Fosdick L, D'Adamio L, Jondal M, Orrenius S (1994) Co-receptor (CD4/CD8) engagement enhances CD3-induced apoptosis in thymocytes. Implications for negative selection. *J Immunol* **153**, 2436-2443.
- Melino G, Annicchiarico-Petruzzelli M, Piredda L, Candi E, Gentile V, Davies PJ, Piacentini M (1994) Tissue transglutaminase and apoptosis: Sense and anti-sense tranfection studies with human neuroblastoma cells. *Mol Cell Biol* **14**, 6584-6596.
- Montague JW, Gaido ML, Frye C, Cidlowski JA (1994) A calcium-dependent nuclease from apoptotic rat thymocytes is homologous with cyclophilin. Recombinant cyclophilins A, B, and C have nuclease activity. *J Biol Chem* **269**, 18877-18880.
- Murphy K, Heimberger A, Loh DY (1990) Induction by antigen of intrathymic apoptosis of CD4+CD8+ thymocytes *in vivo*. *Science* **250**, 1720-1723.
- Mutch DG, Powell CB, Kao MS, Collins JL (1992) Resistance to tumor necrosis factor alpha in malignant gynecological cell lines is associated with the expression of protein(s) that prevent the activation of phospholipase A2 by tumor necrosis factor alpha. *Cancer Res* **52**, 866-872.
- Nakagama T, Ueda Y, Yamada H, Shores EW, Singer A, June CH (1992) *In vivo* calcium elevations in thymocytes with T cell receptors that are specific for self ligands. *Science* **257**, 96-99.
- Neamati N, Fernandez A, Wright S, Kiefer J, McConkey DJ (1995) Degradation of lamin B1 precedes oligonucleosomal DNA fragmentation in apoptotic thymocytes and isolated thymocyte nuclei. *J Immunol* **154**, 3788-3795.
- Nicholson DW, Ali A, Thornberry NA, Vaillancourt JP, Ding CK, Gallant M, Gareau Y, Griffin PR, Labelle M, Lazebnik YA, Munday NA, Raju SM, Smulson ME, Yamin TT, Yu VL, Miller DK (1995) Identification and inhibition of the ICE/ced-3 protease necessary for mammalian apoptosis. *Nature* **376**, 37-43.
- Nikonova LV, Beletsky IP, Umansky SR (1993) Properties of some nuclear nucleases of rat thymocytes and their changes in radiation-induced apoptosis. *Eur J Biochem* **215**, 893-901.
- Norvell A, Mandik L, Monroe JG (1995) Engagement of the antigen receptor on immature murine B lymphocytes results in death by apoptosis. *J Immunol* **154**, 4404-4413.
- Obeid LM, Linardic CM, Karolak LA, Hannun YA (1993) Programmed cell death induced by ceramide. *Science* **259**, 1769-1771.
- Oberhammer F, Wilson JW, Dive C, Morris ID, Hickman JA, Wakeling AE, Walker PR, Sikorska M (1993) Apoptotic death in epithelial cells: Cleavage of DNA to 300 and/or 50 kb fragments prior to, or in the absence of internucleosomal fragmentation. *EMBO J* **12**, 3679-3684.
- Oberhammer FA, Hochegger K, Froschl G, Tiefenbacher R, Pavelka M (1994) Chromatin condensation during apoptosis is accompanied by degradation of lamin A+B without enhanced activation of cd/c2 kinase. *J Cell Biol* **1267**, 827-837.
- Oltvai ZN, Korsmeyer SJ (1994) Checkpoints of dueling dimers foil death wishes. *Cell* **79**, 189-192.
- Orrenius S, McConkey DJ, Bellomo G, Nicotera P (1989) Role of Ca²⁺ in toxic cell killing. *Trends Pharmacol Sci* **10**, 281-285.
- Parry SL, Holman MJ, Hasbold J, Klaus GGB (1994) Plastic-immobilized anti-m or anti-d antibodies induce apoptosis in mature murine B lymphocytes. *Eur J Immunol* **24**, 974-979.
- Peitsch MC, Polzar B, Stephan H, Crompton T, MacDonald HR, Mannherz HG, Tschopp J (1993) Characterization of the endogenous deoxyribonuclease involved in nuclear DNA degradation during apoptosis (programmed cell death). *EMBO J* **12**, 371-377.
- Petit PX, Lecoecur H, Zorn E, Dauguet C, Mignotte B, Gougeon ML (1995) Alterations in mitochondrial structure and function are early events of dexamethasone-induced thymocyte apoptosis. *J Cell Biol* **130**, 157-167.
- Pronk GJ, Ramer K, Amiri P, Williams LT (1996) Requirement of an ICE-like protease for induction of apoptosis and ceramide generation by REAPER. *Science* **271**, 808-810.
- Pushkareva M, Obeid LM, Hannun YA (1995) Ceramide: An endogenous regulator of apoptosis and growth suppression. *Immunol Today* **16**, 294-297.
- Rajotte D, Haddad P, Haman A, Haman EJ, Cragoe J, Hoang T (1992) Role of protein kinase C and the Na⁺/H⁺ antiporter in suppression of apoptosis by granulocyte macrophage colony-stimulating factor interleukin-3. *J Biol Chem* **267**, 9980-9987.
- Ribeiro JM, Carson DA (1993) Ca²⁺/Mg²⁺ dependent endonuclease from human spleen: Purification, properties, and role in apoptosis. *Biochemistry* **32**, 9129-9136.
- Richter C (1993) Pro-oxidants and mitochondrial Ca²⁺: Their relationship to apoptosis and oncogenesis. *FEBS Lett* **325**, 104-107.
- Robertson LE, Chubb S, Meyn RE, Story M, Ford R, Hittelman WN, Plunkett W (1993) Induction of apoptotic cell death in chronic lymphocytic leukemia by 2-chloro-2'-deoxyadenosine and 9-beta-D-arabinosyl-2'-fluoroadenine. *Blood* **81**, 143-150.
- Rosing J, van Rijn JL, Bevers EM, van Diejen G, Comfurius P, Zwaal RF (1985) The role of activated human platelets in prothrombin and factor X activation. *Blood* **65**, 319-332.

- Sarin A, Adams DH, Henkart PA (1993) Protease inhibitors selectively block T cell receptor-triggered programmed cell death in a murine T cell hybridoma and activated peripheral T cells. *J Exp Med* **178**, 1693-1700.
- Sarin A, Nakajima H, Henkart PA (1995) A protease-dependent TCR-induced death pathway in mature lymphocytes. *J Immunol* **154**, 5806-5812.
- Savill J, Fadok V, Henson P, Haslett C (1993) Phagocytic recognition of cells undergoing apoptosis. *Immunol Today* **14**, 131-136.
- Schewe M, Muller P, Korte T, Herrmann A (1992) The role of phospholipid asymmetry in calcium phosphate-induced fusion of human erythrocytes. *J Biol Chem* **267**, 5910-5915.
- Schulze-Osthoff K, Walczak H, Droge W, Krammer PH (1994) Cell nucleus and DNA fragmentation are not required for apoptosis. *J Cell Biol* **127**, 15-20.
- Shaposhnikova VV, Dobrovinskaya OR, Eidus LK, Korystov YN (1994) Dependence of thymocyte apoptosis on protein kinase C and phospholipase A2. *FEBS Lett* **348**, 317-319.
- Shi Y, Sahai BM, Green DR (1989) Cyclosporin A inhibits activation-induced cell death in T cell hybridomas and thymocytes. *Nature* **339**, 625-626.
- Shimizu S, Eguchi Y, Kosaka H, Kamiike W, Matsuda H, Tsujimoto Y (1995) Prevention of hypoxia-induced cell death by Bcl-2 and Bcl-xL. *Nature* **374**, 811-813.
- Sims PJ, Wiedmer T, Esmon CT, Weiss HJ, Shattil SJ (1989) Assembly of the platelet prothrombinase complex is linked to vesiculation of the platelet plasma membrane. Studies in Scott syndrome: An isolated defect in platelet procoagulant activity. *J Biol Chem* **264**, 17049-17057.
- Slater AF, Nobel CS, Maellaro E, Bustamante J, Kimland M, Orrenius S (1995) Nitron spin traps and a nitroxide antioxidant inhibit a common pathway of thymocyte apoptosis. *Biochem J* **306**, 771-778.
- Smith CA, Williams GT, Kingston R, Jenkinson EJ, Owen JJT (1989) Antibodies to CD3/T cell receptor complex induce death by apoptosis in immature T cells in thymic cultures. *Nature* **337**, 181-184.
- Squier MKT, Miller ACK, Malkinson AM, Cohen JJ (1994) Calpain activation in apoptosis. *J Cell Physiol* **159**, 229-237.
- Story MD, Stephens LC, Tomosovic SP, Meyn RE (1992) A role for calcium in regulating apoptosis in rat thymocytes irradiated *in vitro*. *Int J Radiat Biol* **61**, 243-251.
- Sulpice JC, Zachowski A, Devaux PF, Giraud F (1994) Requirement for phosphatidylinositol 4,5-bisphosphate in the Ca^{2+} -induced phospholipid redistribution in the human erythrocyte membrane. *J Biol Chem* **269**, 6347-6354.
- Sun XM, Cohen GM (1994) Mg^{2+} -dependent cleavage of DNA into kilobase pair fragments is responsible for the initial degradation of DNA in apoptosis. *J Biol Chem* **269**, 14857-14860.
- Taresa E, Kedei N, Thomazy V, Fesus L (1992) An involucrin-like protein in hepatocytes serves as a substrate for tissue transglutaminase during apoptosis. *J Biol Chem* **267**, 25648-25651.
- Tepper CG, Jayadev S, Liu B, Bielawska A, Wolff R, Yonehara S, Hannun YA, Seldin MF (1995) Role for ceramide as an endogenous mediator of Fas-induced cytotoxicity. *Proc Natl Acad Sci USA* **92**, 8443-8447.
- Tewari M, Quan LT, O'Rourke K, Desnoyers S, Salvesen GS, Dixit VM (1995) Yama/ CPP32b, a mammalian homolog of ced-3, is a crmA-inhibitable protease that cleaves the death substrate poly (ADP-ribose) polymerase. *Cell* **81**, 801-809.
- Thompson CB (1995) Apoptosis in the pathogenesis and treatment of disease. *Science* **267**, 1456-1462.
- Tokes ZA, Clawson GA (1989) Proteolytic activity associated with the nuclear scaffold. The effect of self-digestion on lamins. *J Biol Chem* **264**, 15059-15065.
- Trent JC, McConkey DJ, Loughlin SM, Harbison MT, Fernandez A, Ananthaswamy HN (1996) Ras signaling in tumor necrosis factor-induced apoptosis. *EMBO J*, in press.
- Treves S, Trentini PL, Ascanelli M, Bucci G, Di Virgilio F (1994) Apoptosis is dependent on intracellular zinc and independent of intracellular calcium in lymphocytes. *Exp Cell Res* **211**, 339-343.
- Tsubata T, Wu J, Honjo T (1993) B-cell apoptosis induced by antigen receptor crosslinking is blocked by a T cell signal through CD40. *Nature* **364**, 645-648.
- Utsugi T, Schroit AJ, Connor J, Bucana CD, Fidler IJ (1991) Elevated expression of phosphatidylserine in the outer membrane leaflet of human tumor cells and recognition by activated human blood monocytes. *Cancer Res* **51**, 3062-3066.
- Vanderbilt JN, Bloom KS, Anderson JN (1982) Endogenous nuclease: Properties and effects on transcribed genes in chromatin. *J Biol Chem* **257**, 13009-13017.
- Verhoven B, Schlegel RA, Williamson P (1995) Mechanisms of phosphatidylserine exposure, a phagocyte recognition signal, on apoptotic T lymphocytes. *J Exp Med* **182**, 1597-1601.
- Verkleij AJ, Zwaal RF, Roelofsens B, Comfurius P, Kastelijn D, van Deenen LL (1973) The asymmetric distribution of phospholipids in the human red cell membrane. A combined study using phospholipases and freezeetch electron microscopy. *Biochem Biophys Acta* **323**, 178-193.
- Voelkel-Johnson C, Thorne TE, Laster SM (1995) Susceptibility to TNF in the presence of inhibitors of

transcription or translation is dependent on the activity of cytosolic phospholipase A2 in human melanoma tumor cells. *J Immunol* **156**, 201-207.

Wang L, Miura M, Bergeron L, Zhu H, Yuan J (1994) Ich-1, an ICE/ced-3-related gene, encodes both positive and negative regulators of programmed cell death. *Cell* **78**, 739-750.

Wever GF, Abromson-Leeman S, Cantor H (1995) A signaling pathway coupled to T cell receptor ligation by MMTV superantigen leading to transient activation and programmed cell death. *Immunity* **2**, 363-372.

Williamson P, Bevers EM, Smeets EF, Comfurius P, Schlegel RA, Zwaal RF (1995) Continuous analysis of the mechanism of activated transbilayer lipid movement in platelets. *Biochemistry* **34**, 10448-10455.

Wolfe JT, Ross D, Cohen GM (1994) A role for metals and free radicals in the induction of apoptosis in thymocytes. *FEBS Lett* **352**, 58-62.

Wolff RA, Dobrowsky RT, Bielawska A, Obeid LM, Hannun YA (1994) Role of ceramide-activated protein phosphatase in ceramide-mediated signal transduction. *J Biol Chem* **269**, 19605-19606.

Wyllie AH (1980) Glucocorticoid-induced thymocyte apoptosis is associated with endogenous endonuclease activation. *Nature* **284**, 555-556.

Wyllie AH, Kerr JFR, Currie AR (1980) Cell death: The significance of apoptosis. *Int Rev Cytol* **68**, 251-305.

Wyllie AH, Morris RG, Smith AL, Dunlop D (1984) Chromatin cleavage in apoptosis: Association with condensed chromatin morphology and dependence on macromolecular synthesis. *J. Pathol* **142**, 67-77.

Wyllie AH, Arends MJ, Morris RG, Walker SW, Evan G (1992) The apoptosis endonuclease and its regulation. *Sem Immunol* **4**, 389-397.

Xie K, Huang S, Dong Z, Juang SH, Gutman M, Xie QW, Nathan C, Fidler IJ (1995) Transfection with the inducible nitric oxide synthase gene suppresses tumorigenicity and abrogates metastasis by K-1735 murine melanoma cells. *J Exp Med* **181**, 1333-1343.

Xue D, Horvitz HR (1995) Inhibition of the *Caenorhabditis elegans* cell-death protease CED-3 by a CED-3 cleavage site in baculovirus p35 protein. *Nature* **377**, 248-251.

Yao B, Zhang Y, Delikat S, Mathias S, Basu S, Kolesnick R (1995) Phosphorylation of Raf by ceramide-activated protein kinase. *Nature* **378**, 307-310.

Yao XR, Scott DW (1993) Antisense oligodeoxynucleotides to the blk tyrosine kinase prevent anti-mediated growth inhibition and apoptosis in a B-cell lymphoma. *Proc Natl Acad Sci USA* **90**, 7946-7950.

Yazdanbakhsh K, Choi JW, Li Y, Lau LF, Choi Y (1995) Cyclosporin A blocks apoptosis by inhibiting the DNA binding activity of the transcription factor Nur77.

Proc Natl Acad Sci USA **92**, 437-441.

Yuan J, Shaham S, Ledoux S, Ellis HM, Horvitz HR (1993) The *C elegans* cell death gene *ced-3* encodes a protein similar to mammalian interleukin-1 β -converting enzyme. *Cell* **75**, 641-652.

Zamzani N, Marchetti P, Castedo M, Decaudin D, Macho A, Hirsch T, Susin SA, Petit PX, Mignotte B, Kroemer G (1995a) Sequential reduction of mitochondrial transmembrane potential and generation of reactive oxygen species in early programmed cell death. *J Exp Med* **182**, 367-377.

Zamzani N, Marchetti P, Castedo M, Zanin C, Vayssiere JL, Petit PX, Kroemer G (1995b) Reduction in mitochondrial potential constitutes an early irreversible step of programmed lymphocyte death *in vivo*. *J Exp Med* **181**, 1661-1672.

Zhivotovsky B, Nicotera P, Bellomo G, Hanson K, Orrenius S (1993) Ca²⁺ and endonuclease activation in radiation-induced lymphoid cell death. *Exp Cell Res* **207**.

Zhivotovsky B, Gahm A, Andarcrona M, Nicotera P, Orrenius S (1995) Multiple proteases are involved in thymocyte apoptosis. *Exp Cell Res* **221**, 404-412.

Zornig M, Busch G, Beneke R, Gulbins E, Lang F, Ma A, Korsmeyer SJ, Moroy T (1995) Survival and death of prelymphomatous B-cells from N-myc/bcl-2 double transgenic mice correlates with the regulation of intracellular Ca²⁺ fluxes. *Oncogene* **11**, 2165-2174.

Zwaal R. F, Roelofsen B, Comfurius P, van Deenen LL (1975) Organization of phospholipids in human red cell membranes as detected by the action of various purified phospholipases. *Biochem Biophys Acta* **406**, 83-96.

Discussion with Reviewers

K.M. Kim: Abnormal increases in Ca²⁺ play a key role in regulation of apoptosis by activation of enzymes that leads to DNA fragmentation. How does the increase in Ca²⁺ come about in the first place? What initially triggers apoptosis?

Author: It appears that Ca²⁺ evaluations in apoptotic cells occur via one of two general mechanisms. The first involves receptor-coupled activation of phospholipase(s) C leading to polyphosphoinositide hydrolysis and production of inositol triphosphate, which promotes release of Ca²⁺ from the endoplasmic reticulum and Ca²⁺ influx across the plasma membrane. This mechanism is analogous to the one involved in growth factor-induced cell proliferation and can therefore be regarded as "physiological." The second involves damage, resulting in redistribution of Ca²⁺ down the concentration gradients maintained between the cytosol and intra- and extracellular Ca²⁺ down the concentration gradients

maintained between the cytosol and intra- and extracellular Ca^{2+} pools. Oxidative stress represents at least one common trigger for the damage-dependent Ca^{2+} increase pathway, and depletion of cellular ATP represents an attractive (hypothetical) second. The initial trigger for apoptosis is unknown, although Zamzani *et al.* (1995b) have advanced the hypothesis that alterations in mitochondrial membrane potential and/or function are involved which could initiate the irreversible disruption of intracellular Ca^{2+} homeostasis.

K.M. Kim: Activation of ICE/CED-3 is said to be involved in the earliest stage of apoptosis. What is the mechanism?

Author: The biochemical mechanisms involved in the activation of the ICE family are not known but are under active investigation. Importantly, the ICE family proteases are not directly Ca^{2+} dependent.

K.M. Kim: Abrogation of apoptosis by BCL-2 is said to involve inhibition of Ca^{2+} transports by membranes. Would the author elaborate on the phenomenon?

Author: Changes in intracellular Ca^{2+} compartmentalization occur in many examples of apoptosis. In a number of cellular systems the protective effects of BCL-2 have been linked to preventing these Ca^{2+} compartmentalization changes, but the mechanism(s) involved have not been elucidated to date. It is possible that BCL-2 serves as a channel regulator or that its effects on Ca^{2+} are secondary to its primary action on mitochondrial membrane potential and/or oxidative stress.

A.R. Eastman: This review focuses on the possible activation of calcium-dependent endonucleases in apoptosis. However, this conclusion is questionable when one considers the potential role of calcium in activation of proteases, calmodulin and tissue transglutaminase. Given these multiple functions, can one really suggest that calcium is required for any one particular step? I agree that evidence clearly supports the presence of a calcium endonuclease in many nuclei, but is there really evidence for its involvement in apoptosis? Most of the evidence is based upon experimentally increasing or decreasing calcium, both these manipulations will equally modify proteases, calmodulin and tissue transglutaminase. The author ignores protein kinase C, yet, this is a calcium-dependent kinase that frequently has been shown to protect cells from apoptosis. This is yet another step in apoptosis that can be modified by changes in calcium.

In a similar vein, the author suggests that calcium is required for lamin degradation, but is lamin degradation required for apoptosis?

Author: Dr. Eastman is absolutely correct in emphasizing the pleiotropic effects of Ca^{2+} in apoptotic cells (and indeed in those protected from apoptosis), and I agree that Ca^{2+} is most likely not acting at one step in the apoptotic pathway but at many. Ongoing investigation aimed at directly identifying the molecular targets for Ca^{2+} in apoptotic cells should help to clarify this issue.

Our hypothesis is that lamin degradation promotes DNA fragmentation. Since Lazebnik has shown that DNA fragmentation can occur under certain circumstances where lamin degradation is blocked (Lazebnik *et al.*, 1995), we would not even state that lamin degradation is absolutely required for endonuclease activation. However, our preliminary work indicates that inhibitors of the lamin protease dramatically delay apoptotic cell death in many of our models; in fact, they appear more potent than are inhibitors of the ICE family. Therefore, we would suggest that activation of the lamin protease may be required for cell death, although its "death substrates" may be quite numerous and diverse.

A.R. Eastman: The author describes an apparent paradox that increasing calcium sometimes induces apoptosis but at other times inhibits it. What are the levels of calcium in each case? Upon withdrawal of IL3, I believe that there is a drop in intracellular calcium, and that A23187 permits recovery, not really an increase in calcium. In contrast, in other cases A23187 causes a dramatic influx of calcium. Could the answer be that cells undergo apoptosis at either low or high calcium, and that in the IL-3 situation, A23187 helps restore normal calcium homeostasis?

Author: It is definitely possible that it is the magnitude of the change in Ca^{2+} concentration, rather than the absolute level achieved in the cytoplasm, that determines whether cell undergoes apoptosis or not. How this $\Delta[\text{Ca}^{2+}]_i$ would engage the enzymatic machinery for apoptosis is not clear, but disruption of subcellular structure (i.e., endoplasmic reticular permeability changes) could be postulated.

A.R. Eastman: Is there evidence for a calcium-dependent and independent mechanism of apoptosis in a single cell line, or does the calcium dependence vary with cell line?

Author: The only clear example I know of involves work with an IL-3-dependent hematopoietic cell line, where Mary Collins and her co-workers (personal communication) have found evidence for the co-expression of both $\text{Ca}^{2+}/\text{Mg}^{2+}$ and acidic endonucleases in isolated nuclei.