

Utah State University

DigitalCommons@USU

---

Bu

Bee Lab

---

1-1-1902

## On the Hymenoptera collected by Mr. Robert Shelford at Sarawak, and on the Hymenoptera of the Sarawak Museum

Peter Cameron

Follow this and additional works at: [https://digitalcommons.usu.edu/bee\\_lab\\_bu](https://digitalcommons.usu.edu/bee_lab_bu)



Part of the [Entomology Commons](#)

---

### Recommended Citation

Cameron, Peter, "On the Hymenoptera collected by Mr. Robert Shelford at Sarawak, and on the Hymenoptera of the Sarawak Museum" (1902). *Bu*. Paper 186.

[https://digitalcommons.usu.edu/bee\\_lab\\_bu/186](https://digitalcommons.usu.edu/bee_lab_bu/186)

This Article is brought to you for free and open access by the Bee Lab at DigitalCommons@USU. It has been accepted for inclusion in Bu by an authorized administrator of DigitalCommons@USU. For more information, please contact [digitalcommons@usu.edu](mailto:digitalcommons@usu.edu).



11, 235

On the Hymenoptera collected by Mr. <sup>Shelford</sup>  
Robert Shelford at Sarawak, and  
on the Hymenoptera of the  
Sarawak Museum.

BY P. CAMERON, OF NEW MILLS, DERBYSHIRE.

This paper is based on material collected at Sarawak, by Mr. Robert Shelford of Cambridge University and on the species in the Sarawak Museum brought home by Mr. Shelford for the purpose of being named. In addition to many known species, the two collections contain many noteworthy undescribed genera and species. Since the publication of the paper by the late Mr. F. Smith (Jour. Linn. Soc. 1857) on the Hymenoptera collected by A. R. Wallace, very little has been written on the Bornean species, of which an immense number must still remain to be discovered in all the families, but more particularly among the smaller parasitic tribes—Ichneumonidæ, Braconidæ, Oxyura and Chalcididæ.

TENTHREDINIDÆ.

*Hylotoma pruinosa*, sp. nov.

Coerulea, dense albo pruinosa; alis hyalinis, macula sub-stigmatali fusca, ♀.

Long: 10 mm.

Hab. Sarawak, Borneo (Shelford).

Bright metallic blue densely covered with a white pile. The flagellum of the antennæ is black, the hinder tibiæ are broadly fuscous in the middle. The frontal fovea is deep, its sides oblique, it extends from the ocelli to shortly below the antennæ and is open above and below; the lateral furrows of the face are wide and deep; the labrum has a slight violet tinge. The vertex and the mesonotum have a slight purple tinge. The cloud on the fore wing occupies all the radial cellule; the upper half



of the first cubital and the greater part of the second, the third cellule being also slightly clouded; the second cubital cellule is distinctly longer at the top and bottom than the third; both the recurrent nervures are received shortly behind the middle of the cellules; the third transverse cubital nervure is angled outwardly above the middle and from the angle a short nervure issues; the upper and lower parts are straight and have an oblique slope. Abdomen coloured like the body; the apex of the first and the base laterally of the second segment are fuscous.

Allied to *H. janthina* Kl. and *H. maculipennis* Cam. Characteristic is the third transverse cubital nervure with its distinct nervure issuing from the angle above the middle.

#### EVANIIDÆ.

##### *Evania borneana*, sp. nov.

Nigra, capite thoraceque albopilosis; mesonoto sparse punctato; alis hyalinis, nervis nigris, ♂.

Long: 8 mm.

Hab. Sarawak (Shelford).

Antennæ longer than the body: the scape not dilated, narrow covered with a pale pile and as long as the following two joints united: the third and fourth joints are about equal in length. Head shining, smooth, almost impunctate, and covered with short white pubescence. Clypeus on the lower side bounded by a distinct curved furrow. On the front, outside the antennæ, is a narrow covered keel. The ocelli are in a curve; the hinder are separated from each other by a distinctly greater distance than they are from the eyes. There is a narrow, but distinct, keel between the antennæ. The mesonotum bears some large scattered punctures; the lateral furrows are distinct, deep and curved; there is a distinct, longitudinal furrow opposite the tegulæ. The scutellum has scattered punctures in four irregular rows. The median segment is regularly reticulated, except in the middle above, where it bears large, deep scattered punctures. The apex of the propleuræ is irregularly furrowed above. The upper part of the mesopleuræ is smooth; the lower regularly punctured. The breast is sparsely, and not very strongly, punctured. Wings clear hyaline: the

stigma and nervures black; the second transverse cubital nervure is obsolete; as is also the cubitus from the first transverse cubital nervure, which is interstitial with the recurrent.

The radial cellule is wide at the apex, through the radius having an oblique downward slope at the base; the apical abscissa is straight and oblique; the transverse basal nervure is almost interstitial. Legs black; the calcaria fuscous; the tibiae without spines.

The metasternal forks are roundly curved. In Schletterer's Monograph (Ann. K. K. H. of Mus. Wien. 1889) this species would come near *E. appendigaster* Linn.

*Megiseleus longicollis*, sp. nov.

Black, the head yellowish-red; the four front legs tinged with rufous; the wings clear hyaline, the cubital and the transverse cubital nervures obliterated; the radius incomplete, ♀.

Long: 18; terebra 17-18 mm.

Hab. Sarawak, Borneo (Shelford).

Antennae black; the basal three joints rufous. Head pale rufous, the orbits with a yellowish tinge; the anterior three tubercles are longer and sharper than the posterior; the front is coarsely, closely striated, obliquely above, transversely below; the vertex, behind the ocelli, is indistinctly furrowed in the middle, and closely obliquely striated on either side of it; the outer orbits are smooth. The prothorax has a brownish tinge; it is distinctly longer than the mesothorax, is deeply and widely incised at the base, the apex of the incision being rounded; the basal half is closely transversely striated, the apex is indistinctly striated; the dilated apex is smooth. Mesonotum coarsely, irregularly reticulated at the base; the apex is widely depressed in the middle, the raised sides are irregularly punctured. Scutellum smooth, its sides punctured. Mesopleurae smooth; the base pilose. Median segment closely and regularly reticulated. The propleurae in front of the tegulae are strongly striated. The hinder coxae are closely, but not strongly, striated; the coxal teeth are irregular and not very prominent. Wings clear hyaline; the nervures and stigma black; the basal abscissa of the radius is straight and oblique and forms an angle with the apical branch which is about equal in length to it; only the basal two

cellules are enclosed or complete, and the only apical nervure is the abbreviated radius.

A species easily known by the abbreviated and obsolete ulur nervures. The pronotum, too, is longer and narrower than it is with the other Oriental species.

*Megiseleus maculifrons*, sp. nov.

Black, the head red; the vertex and the upper part of the front black; the outer orbits dull red, narrowly yellowish on the inner side close to the eyes; there is a broad red mark immediately behind the ocelli; legs black; the four front tarsi dull testaceous, the basal joint of the hinder tarsi white; the wings hyaline with a slight fuscous tinge; the stigma and nervures black, ♂.

Long: 12 mm.

Hab. Baram District. Low country (Hose).

Antennæ black; the scape and pedicle rufous. Head rufous, the outer orbits and the face duller, more yellowish in tint; the vertex and the upper part of the front and the upper part of the outer orbits, black; behind the ocelli is a red mark, which is broader than long. The front tubercle is longer, more sharply pointed than the others and is directed backwards; the hinder pair are shorter and broader than the middle. The vertex is narrowly rugose in the middle, the sides are striated transversely. Mandibles rufous, black at the apex, the palpi black, paler towards the apex. Prothorax short; the pleuræ depressed in the middle.

BRACONIDÆ.

*Iphiaulax*, Foer.

The following three species of *Iphiaulax* are similarly coloured—luteous with the wings fuscous, yellow at the base. They may be separated as follows:

- (a) The keel on the centre of second segment not reaching to the middle of the segment, not much longer than broad; *acragas*.
- (b) The keel on the centre of the second segment reaching beyond the middle.

- The keel reaching to the apex, of equal width throughout; the segment at its sides not depressed, nor strongly transversely striated; *astriochus*.  
 (c) The keel not reaching to the apex; the segment at its sides depressed and strongly transversely striated; *ceressus*.

*Iphiaulax acragas*, sp. nov.

Long: 11 mm; terebra 6 mm.

Hab. Borneo (Shelford).

Antennae black, the scape luteous beneath. Head smooth and shining; the face sharpened, sparsely covered with long fuscous hair; the clypeus is bordered by oblique furrows laterally; apex of mandibles black; the palpi luteous. Thorax smooth and shining sparsely covered with pale pubescence. Legs coloured like the body; the apical joint of the four hinder tarsi fuscous. Wings fuscous, with a slight violaceous tint; the base behind the transverse basal nervure yellowish-hyaline; the first cubital and the discoidal cellule are lighter coloured in the middle; the stigma is black, with a luteous spot on its base. The central part of the petiole is rugose and is stoutly longitudinally striated in the middle; the second is closely rugously punctured; the lateral depressions are wide, deep and closely longitudinally striated; the oblique apical depressions are narrower, are deep and closely striated; the suturiform articulation is deep and rather strongly and regularly striated; the third segment is closely punctured; the basal furrow is wide, deep and closely striated in the centre; it becomes narrowed and curved at the sides; its apical furrow is narrow. The apical segments are smooth; the furrows on the fifth segment are narrow and striated.

*Iphiaulax ceressus*, sp. nov.

Long: 12 mm.

Hab. Matang, 3000 feet (Shelford).

Antennae entirely black. The face is more yellowish in tint than the vertex; its centre is irregularly striated; the sides punctured; the clypeus is distinctly raised and clearly separated from the face. Mandibular teeth black. Thorax smooth and

shining, sparsely pilose. Wings smoothy-fuscos from the transverse basal nervure; behind it yellowish-hyaline; the lower part of the stigma to the transverse cubital nervure luteous. The central part of the petiole is rugose and longitudinally keeled down the centre; the four following segments are closely rugosely punctured; the basal keel extends to near the apex; it is of nearly equal width to near the apex; its sides are keeled and it is distinctly raised; the segment on either side of it is depressed and bears three stout, irregularly curved keels, the lateral depression is shallow; the sides of the segment outside it forms three ridges. The furrows on the other segments are longitudinally striated; the basal branch of the suturiform articulation is larger, broader, and more oblique and more distinctly striated than the apical.

*Iphiaulax astiochus*, sp. nov.

Long: 9; terebra 1-2 mm.

Hab. Sarawak, Borneo (R. Shelford).

Antennae entirely black. Front and vertex smooth, shining and bare; the face irregularly rugose; the mandibular teeth are black. Thorax smooth and shining; the median segment is thickly covered with long pale hair. Wings dark fuscous; yellowish-hyaline behind the transverse basal nervure. The central area of the petiole is rugosely punctured; it is slightly narrowed towards the apex. The middle segments are closely rugosely punctured; the central keel extends to the suturiform articulation; it is of equal width throughout; is longitudinally striated, with the sides raised and irregular; the basal lateral depression is narrow at the base, much wider at the apex and is stoutly striated at the base of the dilated part, which turns inwardly at the apex; the suturiform articulation is wide and deep; its basal lateral fork is more oblique and narrower than the apical; the other furrows are closely striated; the basal more strongly than the apical; the apical segments are smooth and are more yellowish in tint than the others.

*Spinaria curvispina*, sp. nov.

Rufa, thorace spina collaris curvata, abdominis dorso strigoso; capite, thorace pedibusque anterioribus rufis; abdominis



dorso pedibusque posticis nigris; abdominis apice pallide flava; alis fuscis, ♀.

Long 12-13 mm.

Hab. Borneo (Shelford).

Antennae longer than the body, black, tapering towards the apex. Head and antennae rufous, smooth and shining; the median segment irregularly reticulated; more strongly and regularly at the middle than at the base. On the back of the pronotum, at the apex, is a long curved spine, which tapers towards the apex and is directed towards the head; between this and the middle is a large leaf-like plate, hollowed above, its sides curved and expanded outwardly; the lateral wings are narrowed towards the apex; at the base, running up the spine, is a stout keel. At the apex of the middle lobe of the mesonotum is a stout keel; the part on either side of it is striated. On the depression behind the scutellum are six stout longitudinal keels; from the middle of the sixth runs a stout, transverse keel; behind this are three short keels. The sides of the median segment at the apex project into stout teeth. The four front legs are paler in tint than the thorax; the apical joint of the tarsi is black; the hind legs are entirely black. Wings uniformly dark fuscous; the stigma and nervures are deep black; abdomen deep black; the sides of the first and second segments and the base of the first and the apical segment are pale yellow; the dorsum is closely and strongly longitudinally striated; the first two transverse furrows are strongly striated; the sides of the third and fourth segments project into sharp, stout spines; the spine on the fourth is the larger; on the centre of the third at the apex, is a stout triangular tooth, on the middle of the fourth is a larger, sharper one; the apical segment ends in a longish, sharp tooth.

Comes nearest apparently to *S. spinator*, Guer, from Bengal, but differs in the colouration of the wings and legs.

*Shelfordia*, gen. nov.

Median segment with a narrow area in the centre, extending from the base to the apex; second, third and fourth segments of the abdomen closely longitudinally striated; the second with four converging keels; there is a transverse crenulated furrow

on the base of the third segments only, and no oblique ones. Malar space large; the oral region widely open. Front depressed in the middle; the centre with a stout keel; the sides of the depression keeled. The third and fourth joints of the antennae are equal in length and somewhat longer than the fifth. Legs of moderate length; the fore tarsi twice the length of the tibiae. Sheaths of the ovipositor densely pilose.

The middle lobe of the mesonotum is distinctly separated; the scutellar depression is shallow and longitudinally striated; the furrow on the metapleuræ is long, wide and deep; above its apex is a longitudinal keel. The occiput is not margined; the temples are moderately large and are rounded behind. The legs are pilose, but not so thickly as in *Myosoma*. The clypeus is bordered by a wide oblique depression. The second and third segments are distinctly longer than their width at the apex; the abdomen is fully twice the length of the thorax. The inner orbits above are distinctly keeled.

A distinct genus easily known by the area on the median segment, by the longitudinally striated abdominal segments, by the margined upper inner orbits and by the keeled front.

*Shelfordia ruficeps*, sp. nov.

Nigra, capite, pedibus anticis, prothorace mesothoraceque rufis; alis fumatis, ♀.

Long: 11; terebra 22 mm.

Hab. Sarawak (Shelford).

Antennae black; the scape rufous in the middle beneath. Head rufous; the face and clypeus closely punctured and sparsely covered with long black hair; the centre of the face slightly raised and smooth. The clypeus is clearly separated, especially at the sides; the top, in the middle, is transverse, its sides rounded; at the sides is a smooth depression. Mandibles rufous, their teeth black. The vertex is sparsely punctured, especially at the sides; the depressed front is more shining and smooth. Pro- and mesothorax with the scutellum smooth and impunctate; the scutellar depression is narrow and is stoutly longitudinally crenulated. The median segment is smooth and is sparsely covered with longish black hair. The anterior wings are uniformly smoky-fuscous; the hinder have the basal half milk-



white; the nervures and stigma are black; in the anterior is an oblique white cloud on the base of the first cubital cellule and extending obliquely into the discoidal. The front legs are rufous like the thorax; the base of the middle coxæ rufous; the rest of them and femora piceous; the hinder legs are entirely black; the coxæ impunctate. The petiole is raised in the centre; the raised part is bordered by an irregular stout keel; the centre towards the apex is irregularly transversely striated. The centre of the second segment is raised, the raised part is bordered by a keel and becomes gradually narrowed towards the apex which is sharply pointed; it is irregularly striated; bordering this is an area of equal width extending from the base to the apex of the segment; it is stoutly transversely striated, with some irregular longitudinal striae in the middle; outside this is an area which becomes gradually wider towards the apex and bears some stout, irregular striae. The third segment and the basal two-thirds of the fourth are stoutly longitudinally striated; the transverse furrow on the base of the third segment is stoutly striated; the apical segments are smooth; the basal four ventral segments are milk-white.

*Myosoma*, Brulé

The species of this genus here described are black; with the head, thorax and the two or four anterior legs rufous, and the wings dark fuscous. The species may be divided into two sections:—

- (a) Head, thorax and the four anterior legs rufous. The third segment with a deep wide transverse striated furrow near the centre, the base stoutly, longitudinally striated; *forticarinata*.  
The third segment without a transverse furrow; the base not stoutly striated; *brevicarinata*.
- (b) Head, pro- and mesothorax rufous, the two anterior legs rufous. The ovipositor not much longer than the body.
- (a') The keel on the second segment narrow extending to the apex of the segment; its base and apex dilated smooth and shining; *longicarinata*.

- (b) The keel on the second segment broad, not extending to the apex of the segment, its base not smooth.

The second segment at the base on either side of the keel deep black, striated, the middle segment deep black;

*fuscipennis*.

The second segment at the base on either side of the kee brownish, not striated; the middle segments brownish;

*trichiura*.

*Myosoma forticarinata*, sp. nov.

Long : 13; terebra 19 mm. ♀.

Hab. Sarawak, Borneo (Shelford).

Antennæ black, the scape rufous. Head and thorax rufous, smooth and shining; the face rugosely punctured, the punctures running into reticulations and sparsely covered with longish black hair; there is a narrow furrow on the front. Mandibles and palpi rufous; the mandibular teeth black. The four front legs are rufous; the middle pair darker in tint, especially on the tibiae and tarsi; the tibiae and tarsi are thickly covered with longish dark hair. Wings dark fuscous; the nervures and stigma black. Petiole broadly depressed, on the sides the central part is bordered by stout keels and there is a stout keel down the centre; the basal part bears some stout irregularly curved keels. The second segment bears a complete central and a shorter lateral keel, which converge slightly towards the apex; the basal area is triangular, smooth and shining; the base in the centre is depressed, the centre bears some stout irregular keels; the outer parts bear stout, oblique keels; the inner parts at the apex being smooth and more deeply depressed than the outer. The two transverse depressions are wide, deep and stoutly longitudinally striated; the part between the two is stoutly longitudinally striated. The apical segments are thickly covered with long black hair. The sheaths of the ovipositor are stout and are thickly covered with black hair. The tarsal spines are rufous.

*Myosoma brevicarinata*, sp. nov.

Long : 12; terebra 21 mm. ♀

Hab. Borneo (Shelford).

Scape of antennæ rufous; the flagellum is brownish beneath, especially towards the apex. Head rufous; the face slightly

paler, more yellowish in tint; rugosely punctured and covered with pale hair; the front and vertex are more sparsely covered with longer darker hair. Mandibles black, broadly rufous at the base. Thorax smooth and shining; the middle of the mesonotum behind is broadly expressed; the central lobe is clearly defined; the median segment is thickly covered with longish fuscous hair. Wings dark fuscous, with a violaceous tinge at the base; the nervures and stigma are black. Metapleuræ widely and deeply furrowed down the middle. The front legs are rufous with a yellowish tinge; the middle pair are brownish; the femora paler; the tarsal spines are black. The petiole is broadly keeled down the centre; the sides on the apical half are stoutly irregularly striated; the keel on the second segment extends to the transverse furrow; its dilated base is finely striated; the sides and centre are irregularly, longitudinally striated to near the apex which is smooth and laterally expressed; the third segment at the base is longitudinally striated; the striae in the middle reaches to the apex; the apical segments are brownish.

*Myosoma trichiura*, sp. nov.

Long: 14; terebra 14-15 mm.

Hab. Sarawak (Shelford).

Antennæ black, the base and apex of the scape rufous. The sides of the head are punctured; in the centre of the face are three stout, transverse keels. Mandibles rufous, their apex broadly black. The face is rather thickly covered with long black hair; the frons and vertex sparsely so. Pro- and mesothorax rufous, smooth and shining; the median segment black. Legs black; the anterior pair rufous; the apex of the median segment is sparsely, longitudinally striated. The apex of the petiole is coarsely irregularly reticulated the lateral furrows are depressed and transversely striated. The second segment is irregularly longitudinally striated; its basal area is broad at the base, becomes gradually and sharply narrowed towards the apex, is longitudinally striated and extends beyond the middle; on either side is an oblique, closely striated furrow. The third and fourth segments are closely striated; the basal furrow on the third extends to the sides; from it issues a curved furrow; the furrow on the fourth does not extend to the

sides in a straight line but curves broadly backwards to it; there is a transverse, striated furrow on the apex of the third and fourth segments. The sheaths of the ovipositor are broad and thickly covered with stiff black hairs.

This species is an *Iphiaulax* with the hairy ovipositor and legs of a *Myosoma*.

*Myosoma longicarinata*, sp. nov.

Long: 11-12 mm ♀.

Hab. Sarawak. (Shelford).

Antennæ entirely black, head rufous; the face more yellowish in colour; it is smooth, sparsely punctured on the sides and bearing some longish black hair; the clypeus is irregularly rugose. Mandibles black, the base broadly rufous. Pro- and mesothorax rufous; the median segment, the base of the pleuræ and the lower part of the furrow black. The two front legs are coloured like the thorax; the middle have a piceous tinge; the tarsal spines are for the most part black. Wings dark fuscous, with a violaceous hue; the stigma and nervures are black. There is a stout longitudinal keel on the basal half of the petiole; the apical part is irregularly longitudinally, stoutly striated; the lateral depression bears stout, transverse keels. The central part of the second segment is obliquely narrowed towards the apex; the central keel reaches to the apex; there is a smooth triangular area at its base and a semicircular one at its apex; the central part is reticulated somewhat strongly; the lateral is closely longitudinally striated. The two transverse furrows are longitudinally striated; the second more finely and closely than the first. The third segment is closely longitudinally striated and is broadly depressed in the centre; the fourth is more closely and finely striated, the third and fourth segments are brownish.

*Myosoma fuscipennis*, sp. nov.

Long: 16 mm. terebra 16 mm.

Hab. Borneo (Shelford).

Antennæ black; the scape piceous in the middle below. Head mahogany coloured, smooth and shining; the centre of the face marked with some irregular transverse keels; the sides are

sparsely punctured. Mandibles black, broadly rufous at the base; the palpi rufo-testaceous. Thorax smooth and shining rufous; the median segment black. The anterior legs rufous; the posterior four black; the middle trochanters rufous. Wings brownish-smoky, with a slight violaceous tinge at the base; the stigma and basal nervures black; the apical nervures fuscous. The sides and ventral surface of the petiole are pale testaceous; its base is smooth; the rest of it coarsely reticulated. The second and third segments are closely, irregularly longitudinally punctured, at the apex almost reticulated; the fourth is closely reticulated, the centre more strongly than the sides; the other segments are smooth. The keel on the base of the second segment is not very distinctly defined; it is broad at the base, becomes gradually obliquely narrowed towards the apex, is more strongly striated than the rest of the segments and reaches to shortly beyond its middle; the lateral furrows are wide, oblique and stoutly striated; the lateral furrows on the third and fourth segments are narrower and more roundly curved; the transverse furrows on the base and apex of the third and fourth segments are closely longitudinally striated.

*Holcotroticus*, gen. nov.

Claws of all the legs bifid; the inner claw smaller than the other. Median segment completely areolated. Apex of the scutellum bifurcate, bordered by stout keels; the apex of the median segment obliquely depressed. Malar space moderately large, the oral region not greatly lengthened. Wings longer than the body; the areolet narrowed at the top; the nervures without a stump of a nervure. The oblique mesopleuræ furrow is wide and deep; a narrow oblique one runs from its middle. Pro- and mesopleuræ above with an oblique keel; between the tops of the middle and hinder coxæ is a stout keel. The fore tarsi are longer than the tibiæ; all the joints are longer than broad; the middle three are much shorter than the others. The hinder legs are much longer than the others; their coxæ are large, more than twice longer than wide. The joints of the maxillary palpi are elongated, the three penultimate joints are not short, compressed or lenticular. The occiput is margined, the sides more distinctly than the upper part; the two hinder ocelli are flat, not convex; there is a



small single keel between the antennæ; the scape of the antennæ is stout, about twice longer than broad; the front is keeled in the centre; the keel is stout with oblique sides; it is not hollowed, and there is no keel on each side. The first cubital cellule is clearly separated from the first discoidal; all the apical nervures in the hind wings are obliterated; there is only the brachial cellule defined and there are no transverse nervures; the radial cellule in the fore wing is elongate, narrow.

The affinities of this genus are with *Agathis* and *Troticus*. From the former it may be separated by the cleft claws; in the latter peculiarity it agrees with *Troticus*; but it wants the abnormal palpi of that genus. The hinder legs are longer than they are with *Agathis*, and this is also the case with the wings. The middle lobe of the mesonotum is distinctly raised; the scutellar depression is wide and narrow; the scutellum is keeled at the base.

*Holcotroticus ruficollis*, sp. nov.

Niger, capite, prothorace, mesothorace, pedibusque anticis rufis; alis fumatis, nervis stigmatæque nigris ♂.

Long: 8-9 mm.

Hab. Sarawak, Borneo (Shelford).

Antennæ black; the scape rufous. Face and clypeus shining, sparsely and slightly punctured and thickly covered with short, blackish hair. Thorax rufous; the median segment, except the pleuræ at the base which are rufous, black. Mesonotum sparsely and indistinctly, the scutellum more distinctly, punctured; its apex stoutly keeled and depressed laterally above. Mesopleuræ, below the oblique furrow, closely and distinctly punctured; at the base above are two stout oblique keels; the furrow is wide and bears some stout oblique keels; below the middle behind is an oblique furrow; the apical bordering furrow bears some stout keels. The basal furrow on the metapleuræ is wide and deep; on it is an upper and three lower stout, widely separated, keels. On the base of the median segment are three, on the apex five areæ; the central basal area is smaller than the lateral and is stoutly transversely striated; the central apical one is hollowed and is gradually narrowed towards the apex; there is also an outer

large spiracular area. The two front legs are of a paler rufous colour than the thorax; the middle pair have a piceous hue. The wings are of a uniform dark fuscous-violaceous colour, with black nervures and stigma. Abdomen deep black, smooth and shining; the petiole is furrowed along the sides.

The malar space is not quite so long as in typical *Agathidini*, the head not being so rostriform as usual. It differs also from *Agathis* in the hinder claws having a tooth near the base.

ICHNEUMONIDÆ.

*Siphimedia*, gen. nov.

Wings without an areolet; the recurrent nervure interstitial, or almost touching the transverse cubital; the transverse basal nervure interstitial; the transverse median nervure in hind wings broken shortly above the middle. Mesonotum trilobate; the middle lobe transverse at the base, sharply pointed at the apex; the parapsidal furrows deep. Post-scutellum deeply bifoveate at the base. Median segment areolated more or less with three or four basal areæ; the areola distinct; the spiracles are linear, longish and are placed in the middle. Legs, and especially the posterior, stout; the hinder femora thickened, shorter than the tibiæ; the four front claws are cleft, the hinder simple; the hind coxæ are twice as long as thick, the basal joint only as long as the following three united and not much longer than the apical. The antennæ stout, shorter than the body. The petiole is twice longer than the width at the apex; it is broad at the base and becomes gradually wider towards the apex; the spiracles are placed shortly behind the middle; the abdominal segments are smooth, without furrows or depressions; the second and following segments are wider than long; the hypopygium is very large, plough-share-shaped and projects beyond the dorsal segment; the ovipositor is stout and is about as long as the body. The front legs have one, the four hinder two, spurs. There is a curved furrow on the sides of the meso-sternum.

The antennæ are placed well up on the face, above the middle of the eyes, which are parallel; the apex of the clypeus is rounded; the mandibles have two subequal teeth; the clypeu



is not distinctly separated from the face. The head is wider than long, is not much developed behind the eyes, and is obliquely narrowed there; the labrum does not project. The transverse cubital nervure has the stump of a nervure in the middle. This genus is founded on the species which I doubtfully referred to *Macrogaster* (*Manchester Memoirs*, XLIII, 193). Its real position is with the *Acvenitini*, and comes, in Ashmead's arrangement, nearest to *Arotes*, which may be known from it by the long and slender hinder tibiae and tarsi, the tibiae being almost twice the length of the femora. From *Acvenites* it may be separated by the deep parapsidal furrows, and by the areolated median segment. To this genus may be referred the following species.

*Siphimedia bifasciata*, sp. nov.

Nigra, facie clypeoque flavis; alis hyalinis, fusco-bifasciatis.  
Long: 10 mm.

Hab. Sarawak (R. Shelford).

Head black, the face, except for a triangular black mark in the middle above, and the greater part of the clypeus, pale yellow. The face is strongly, transversely punctured; the apex smooth and depressed; the upper part of the head smooth and shining, and has a distinct plumbeous hue, as have also the pro- and meso-thorax which are smooth and shining; the apex of the middle lobe is stoutly transversely striated. There are three stout longitudinal keels on the scutellar depression. The areola is longer than wide, becomes slightly narrowed towards the apex, which is rounded; the apical central area is long and extends to the apex, its apical half is widely dilated and has the sides stoutly striated; on the top is a stout, curved transverse keel; the outer central area has two stout longitudinal keels in the middle. The median segment is irregularly punctured and coarsely irregularly longitudinally striated in the middle, especially at the base. The mesosternal furrow is wide and is coarsely transversely striated, and is thickly covered with white pubescence. Wings hyaline; there is a fuscous cloud, extending from the stigma to the cubitals and there is a similar cloud at the apex. Legs black; the anterior tibiae and femora testaceous in front, and fuscous behind. The apical dorsal segments of the abdomen are lined with white.

This species is nearly allied to *S. nigricans*, it is smaller, has the mesonotum impunctate, not strongly punctured; there are only two central areae on the median segment, and the basal area is longer than wide, not wider than long.

*Rhyssa nigratarsis*, sp. nov.

Nigra, late flavo-maculata; abdominis apice late brunnea; pedibus flavis, tibiis femoribusque nigro lineatis; tarsis posterioribus nigris; alis hyalinis, stigmate testaceo ♂.

Long: 17 mm.

Hab. Borneo (Shelford).

Antennae black, the scape yellow beneath. Head black, the face entirely; the inner upper orbits to the ocelli, and the outer entirely, yellow. Face almost smooth, sparsely pilose, shining. Clypeus brownish; in the centre, at the apex, is a short, broad tooth with an indistinct tubercle on either side of it; above, on either side, is a fovea. On the thorax the following parts are lemon-yellow; the sides and base of the pronotum; the basal two thirds of the scutellum, the post-scutellum, the median segment, except at the base and sides; the tubercles and a large oblique line below them. The transverse striation, on the mesonotum does not extend to the apex; the scutellum is closely rugose; the median segment is smooth and shining, its base is depressed and black. The fore legs are yellow; the femora are brownish behind; the fore tibiae are as long as the basal joint of the tarsi; the middle coxae, femora and base of tibiae are lined behind with black; the four hinder tarsi are black; the hinder coxae are broadly black below and laterally; the femora are broadly black above; the tibiae black, with a narrow yellow band near the base and a broader one at the apex. The basal two dorsal segments of the abdomen are black, lined with yellow down the centre; the third is brownish-black, banded with black near the middle; the others are brownish; on the fourth segment is a broad yellow band near the middle; the fifth is indistinctly yellow near the apex. The basal half of the ventral surface is pale yellow, marked with black; the apical brownish. The pedicle of the areolet is nearly as long as the basal abscissa.

*Xanthopimpla labiata*, sp. nov.

Lutea, basi late apiceque mesonoti nigris; thorace laevo; abdomine late nigro-maculato; alis hyalinis, apice fusco-violaceo ♀.

Long: 13; terebra 2 mm.

Hab. Sarawak.

Antennae black; the scape yellow below. Head pallid yellow; above smooth, the face and clypeus closely but not strongly, punctured; the vertex in the centre is black from shortly behind the ocelli and this black mark is continued half way down the front, it becoming gradually narrower as it does so; there is an irregular black transverse mark on the centre of the occiput. The thorax is deeper in tint than the head; and is quite smooth, without any punctures; near the base of the mesonotum is a large irregular mark which extends to the sides; it is broadly rounded at the base; with short blunt projections in the middle; the sides at apex project, the projections becoming gradually narrower towards the apex; the apex of the mesonotum is broadly black; the black mark behind being continued into the scutellar depression. Scutellum smooth, broadly rounded, smooth; the sides and apex keeled. On the basal half of the median segment are four lateral and one large central area; the central area does not project beyond the lateral at the apex, which is transverse; its basal third is obliquely narrowed; the apical two thirds are also obliquely narrowed; the apical lateral areae are quadrangular and of equal width throughout; the apical half is keeled on the outside. Wings clear hyaline, except at the apex which is narrowly fusco-violaceous; the areolet is small and is distinctly appendiculated; the upper half of the recurrent nervure is sharply angled. The abdomen has the four middle segments closely punctured; the basal segment has a broad irregular black mark in the centre; the second and fifth segments have two large marks, those on the third and fourth being larger than the others; there are two minute marks on the sixth, and two broad, transverse ones on the seventh; the last segment is immaculate.

The labrum is longer than usual, being longer than the clypeus, it becomes gradually narrowed from the base to the

apex; the eyes are large, parallel; the malar space is small; the spiracular area is obsolete; below the spiracles, in the middle of the metapleuræ, is a curved longitudinal keel, which originates shortly behind the middle, but not extending to the base; there are two transverse lines on the base of the median segment; the bases of the tibiæ are black. The basal segment is longer than the width of its apex; the other segments are all wider than long; the transverse and oblique furrows are distinct; the head is only very slightly developed behind the eyes.

Comes nearest perhaps to *X. punctata* Fah. The edge of the pronotum is more sharply raised than usual; the base of the middle lobe of the mesonotum more distinctly separated, and the labium more projecting and sharper pointed.

*Xanthopimpla nigrobalteata*, sp. nov.

Lutea, nigro-balteata; mesonoto lævo, nigro, flavo-bilineato; pedibus flavis, late nigro-lineatis; alis hyalinis, stigmatibus fusco.

Long: 12 mm.

Hab. Borneo (Shelford).

This species forms a section with three large areas on the median segment and only one apical.

Antennæ brownish on the under side; the scape yellow beneath; head luteous; the ocellar region and the greater part of the occiput black. Face sparsely punctured, slightly keeled down the middle. Clypeus smooth and shining, its apex broadly rounded. Mandibles black at the apex. Mesonotum smooth and shining, base covered with a short pile, black; the sides and two lines in the middle, extending from the base to near the apex, bright yellow. Scutellum thickly covered with long black hair; its apex, from shortly behind the middle, black. The base of the median segment between the stigmas and to near the apex of the area is black; the central area is nearly square; the lateral are longer and narrowed towards the apex and are longer on the outer than on the inner side, the apex being oblique. The apex of the propleuræ, the base of the meso-, the top below the tubercles and the apex except a large spot above, and the basal half of the metapleuræ, are black. Legs coloured like

the body; the top of the four front femora, the extreme base of the hinder, an oblique mark near the apex of the hinder and their extreme apex; the basal two thirds of the four anterior tibiae, the base of the hinder pair and a broad band on their middle and the four hinder tarsi, black. The middle of the petiole broadly, the sides at the base and the base of the other segments broadly, black; the bands on the apical segments occupy more of the segments than on the basal; the segments and the transverse furrows are smooth, impunctate; there is an oblique furrow on the sides of the second and third segments and a narrower one on the fourth. The abdominal segments are all distinctly broader than long; the middle of the mesonotum is distinctly raised at the base; the scutellum is roundly convex and is not raised above the level of the mesonotum; its sides are keeled.

A species not difficult to separate from any of the described species by the smooth thorax and abdomen, by the black abdomen, banded with yellow, by the black mesonotum with two yellow lines, and by the three large areas on the base of the median segment. It has a very similar appearance to *Chrysopimpla ornatipes* Cam.

*Charitopimpla*, gen. nov.

Apex of the clypeus obliquely depressed and with semicircular emargination; its top separated from the face by a straight furrow. Areolet small, triangular; the recurrent nervure is received at the apex. Abdominal segments closely punctured; segments 2-4 with transverse depressions, which are prolonged obliquely backwards; there is also a shallow transverse furrow at the apex. Median segment smooth without keels. Tarsi spinose; the claws simple. The transverse median nervure in hind wings is broken far below the middle.

The areolet is straight, not oblique: the transverse basal nervure is interstitial; the eyes are large, parallel and reach quite close to the base of the mandibles. The second to fourth abdominal segments are, if anything, wider than long, almost square; the metathoracic spiracles are small, oval. The last joint of the antennae is not longer than the preceeding two united. The legs are short; the hinder tarsi are shorter than



the tibiae; the anterior are longer. The antennae are stout and taper towards the apex.

Characteristic of this genus is the obliquely depressed semi-circularly emarginate apex of clypeus. It comes near to *Erythropimpla*, Ashmead.

*Charitopimpla flavo-balteata*, sp. nov.

Nigra, abdomine flavo-balteato; pedibus flavis; posticis nigro alboque maculatis; alis hyalinis, stigmatibus testaceo, nervis nigris ♀.

Long: 13; terebra 7 mm.

Hab. Sarawak, Borneo (Shelford).

Antennae about two thirds of the length of the body; black, distinctly tapering towards the apex; the scape yellow beneath. The face and clypeus are shining, have a plumbeous hue and are uniformly but not strongly punctured; the face is covered sparsely with white, the clypeus with dark hair; the front and vertex are bare, smooth and shining. Mandibles black; the palpi lemon-yellow. Pro- and mesothorax smooth and shining, covered closely with short fuscous hair; the scutellum bears longer and paler hair; the median segment is, especially towards the apex, thickly covered with long white hair. The lower part of the metapleuræ is smooth and bare. The four front legs are yellow, with the femora suffused with fulvous; the hinder legs white; black are the coxæ, except above the base and apex of the femora, a ring near the base of the tibiae, a broader band on their apex and the apical joints of the tarsi; the femora have the sides and lower surface rufous; the coxæ are bright yellow above. Wings hyaline, with a slight fulvous tinge. Abdomen black; the apices of the basal five segments bright lemon-yellow; the sixth is yellow on the sides; the seventh broadly in the middle above; the segments are closely punctured; the petiole has the middle smooth and slightly depressed; the apical two segments are impunctate, the transverse depressions are shallow except on the fifth where they are wider and deeper and the basal one is striated.

## OPHIONIDES.

*Anomalon perornatum*, sp. nov.

Nigrum, abdomine late ferrugineo; pedibus anterioribus, basi tibiæ posticarum late tarsisque posticis flavis; alis fulvo-hyalinis, stigmate testaceo ♀.

Long: 22 mm.

Hab. Borneo (Shelford).

Antennæ black; the scape yellow beneath. Head black; the face clypeus, labrum, the inner orbits, the malar space and the mandibles, except at the apex, bright lemon-yellow. The face broadly projects in the centre; the sides and upper part are irregularly wrinkled; the clypeus is smooth; the front, especially above, is coarsely, irregularly striated; above, the striæ are oblique; below, they almost form reticulations. Thorax entirely black; the median segment is coarsely, irregularly reticulated; the scutellum is coarsely reticulated and striated; above, it is flat in the centre and has an oblique slope; the mesonotum is rugosely punctured and irregularly reticulated; the apex is somewhat strongly transversely striated. The upper half of the propleuræ is closely rugose; the lower in the middle stoutly, longitudinally striated; the mesopleuræ above the middle coarsely striated, at the base reticulated. The four front legs are bright yellow; the apical three joints of the tarsi black; the hinder legs black; the basal two thirds of the tibiæ and the tarsi yellow; the trochanters beneath and the base of the femora fuscous. Abdomen ferruginous; the petiole and the second segment above black.

*Anomalon fuscicorne*, sp. nov.

Long: 15 mm. ♀.

Hab. Borneo (Shelford).

This species resembles closely the preceding species. The differences between the two may be expressed thus:

Face not raised in the middle; the front with oblique striæ; the upper part of the mesopleuræ coarsely longitudinally striated, the lower smooth.

Face not raised in the middle; the front not striated; the upper part of the mesopleuræ closely reticulated.



Antennae brownish beneath; the scape yellow below. The face, clypeus, the inner orbits to near the ocelli, an oblique mark on the top of the eyes, the malar space, the lower orbits and the mandibles except at the apex, lemon-yellow. The front and lower part of the vertex are stoutly longitudinally striated; the striae are curved and form almost reticulations; the face in the middle is irregularly longitudinally striated. The middle lobe of the mesonotum is coarsely longitudinally reticulated; the lateral are closely rugose. Scutellum coarsely rugose. Median segment coarsely reticulated, the top more distinctly than the sides, which have the reticulations less distinct on the lower part. The upper part of the propleurae is coarsely reticulated, as is also the upper part of the meso-pleurae, but less closely and not so distinctly. The four front legs are yellow; the femora are more fulvous in tint; the hinder legs are black; the apex of the coxae, the basal joint of the trochanters and the basal third of the tibiae, dark rufous; the hinder tarsi yellow. Abdomen ferruginous; the petiole, the second segment above and the apical segment black.

*Anisobas cincticornis*, sp. nov.

Ruf. flagello antennarum nigro, medio albo annulato; alis hyalinis, nervis stigmatumque nigris, ♀.

Long: 10 mm.

Hab. Sarawak, Borneo (R. Shelford).

Antennae black; the base rufous; the seventh to the fourteenth joints for the greater part white; the basal joints of the flagellum are rufous below. The front is obscurely punctured; the face is distinctly but not very closely punctured; the clypeus is obscurely punctured above, below smooth and shining; the labrum is fringed with long hair. The mesonotum is darker coloured than the rest of the thorax and is shagreened; the scutellum is thickly covered with longish black pubescence. The basal three areas of the median segment, are smooth and shining; the others are closely, rugosely punctured; the posterior median is smooth, with the sides slightly striated; the lateral teeth are large, and narrowed gradually towards the apex. Propleurae punctured above; the apex irregularly striated in the middle;

the base and the lower half of the mesopleuræ closely, but not strongly punctured; the middle longitudinally striated; the metapleuræ punctured at the base, the rest closely longitudinally striated. Legs coloured like the body, the hinder tarsi black. The base of the wings have a fulvous tinge. Abdomen shining, the middle segments aciculated; the gastrocoeli are smooth, shallow.

It is doubtful if this is a true *Anisobas*. The antennæ are stout and are slightly thickened towards the apex; the basal joints of the flagellum are all much longer than broad; the face is obliquely narrowed from the top to the bottom; the labrum projects and is narrowed towards the apex; the keel on the propleuræ (characteristic of typical *Anisobas*) is stout; the scutellum is obliquely raised, the sides stoutly keeled and the apex at the top depressed; the median segment is completely areolated; the areola is longer than wide, is not much narrowed towards the apex and rounded backwards at the base and apex; the sides are stoutly spined. The wings are as in *Ichneumon*. The abdomen is not much longer than the head and thorax united; there are seven segments; the last is large, above nearly as long as the sixth; the ovipositor largely projects.

*Bodargus*, gen. nov.

Eyes placed high up, separated by their own length from the base of the mandibles. Face and clypeus forming almost one piece; the suture separating them being almost obsolete; the foveæ are shallow; apex of clypeus transverse, its sides broadly rounded. Occiput deeply emarginate. Antennæ shorter than the body, serrate. Scutellum roundly convex; the sides stoutly keeled. Median segment depressed at the base; the areola is faintly indicated, is twice longer than broad, is open at the base and is gradually narrowed towards the apex; the other areæ are obsolete, except on the apical slope where there are three. Areolet much angled, narrowed at the top. The transverse basal nervure is interstitial; there is a short nervure on the cubital-disco nervure and a longer more distinct one on the recurrent; the transverse median nervure in the hind wings is broken far below the middle. Legs short; the hinder femora do not reach much beyond the apex of the second segment. The

middle segments of the abdomen slightly project at the apex; they are closely punctured; the last segment is as long as the penultimate. The main characteristics of this genus are the flat face, continuous with the clypeus, the indistinctness of the keels on the median segment; the short legs and the large, roundly convex, sharply keeled scutellum.

*Bodargus rufus*, sp. nov.

Ferruginea, apice femorum posticorum, basi tibiae, apice late apiceque tarsorum posticorum nigris ♂.

Long: 15 mm.

Hab. Sarawak, Borneo (Shelford).

Antennae dark rufous, darker towards the apex; the scape is yellow. Head uniformly coloured; the face and base of clypeus closely punctured; the vertex is more closely and strongly punctured; the ocellar region black. Mesonotum closely and distinctly punctured; the scutellum is more strongly and not quite so closely punctured, except on the basal slope. The median segment is rugosely punctured, except on the basal slope; on the apex it is transversely striated. Legs coloured like the body; the apical fourth of the hinder femora, the base of the tibiae narrowly and their apex broadly, the apex of the metatarsus, the apical half of the second joint and the whole of the others, black. Wings hyaline, the stigma and nervures fuscous-black. Abdomen coloured like the thorax; the fifth, sixth and base of seventh blue-black; the apical half of seventh white.

*Diapetus*, gen. nov.

Median segment smooth and shining; its base broadly depressed in the middle; there are two stout, transverse keels. Prothorax with a stout, oblique keel above the middle; its base sharply keeled. Areolet minute, not clearly defined through the cubital and radial nervures uniting; the apical abscissæ of the radius and the cubitus spread out obliquely from it. Parapsidal furrows deep, uniting at the apex into one short, wide furrow. Metathoracic spiracles large, linear. Petiole curved, not much narrowed at the base; the spiracles are placed close to the middle, nearer the apex than to the base.

The antennæ are longer than the body, are filiform, and have the third joint distinctly longer than the fourth; the clypeus is roundly convex and is separated by a deep furrow from the face; the mandibles have two unequal teeth; the meso-pleural furrow is wide and deep and is interrupted in the centre. The transverse median nervure is received behind the transverse basal; the stigma is narrow, lanceolate in the hind wings; the cubital nervure is broken above the middle. The legs are long and slender; the claws moderate in size, the hinder tarsi are longer than the others. The abdomen is bluntly pointed and the last segment is larger than the penultima.

This genus may be referred to the *Cryptinas*, but it does not quite agree with that group, as the spiracles on the petiole are placed near the middle. The small, or more correctly, obsolete areolet might place it in the *Mesostenini*; but there is no known genus in that group in which it could be placed.

*Diapetus nigroplagiatus*, sp. nov.

Rufo-fulvo, vertice, mesonoto, metanoto, pleuris abdomineque late nigro-maculatis; alis flavo-hyalinis, nervis stigmatique testaceis ♀.

Long: 12; terebra 2 mm.

Hab. Borneo (Shelford).

Antennæ longer than the body, darker towards the apex. Head smooth and shining; the vertex and upper part of the occiput largely black; the front is broadly dark rufous in the middle. Face and clypeus yellowish, smooth and shining, sparsely covered with long black hair. Mandibles broadly black at the apex, yellow at the base, rufous in the middle. Thorax smooth and shining except the apex of the middle lobe and the furrows. Scutellum and post-scutellum yellow. The base of the median segment is black except in the middle depression; between the two keels are two large black marks, rounded and narrowed at the apex; the basal two-thirds of the mesopleuræ black; the middle of the metapleuræ broadly, and the greater part of the mesosternum black. Legs coloured like the body; the hinder femora are darker coloured at the base; the tarsi are minutely spinose. Wings yellowish-hyaline; the stigma testaceous; the nervures are of a darker testaceous colour. The petiole i

lighter in colour than the other segments; its central region is broadly black except narrowly down the middle; the second segment is black at the base to near the middle; the third has the basal third black; the fourth and fifth are more narrowly black at the base. On the metapleuræ are, in the middle, four short stout keels; the middle two are longer than the others.

*Acleasa*, gen. nov.

Median segment reticulated all over, without transverse keels; the apex with two large conical teeth. Thorax about three times longer than broad; the mesonotum with indistinct parapsidal furrows, and coarsely reticulated. Metapleural keels absent. Areolet large, wider than long, of equal width throughout; the transverse cubital nervures slightly oblique; the apical one distinct; the transverse median nervure is received behind the transverse basal; the transverse median nervure in hind wing broken shortly below the middle. Legs of moderate length; the basal joint of the hinder tarsi is thickened; the claws are small. The petiole becomes gradually wider towards the apex; its sides near the middle on the lower side project into a stout triangular tooth, the part behind this being keeled; in front of it is a rounded tubercle. The head is rather narrow; the eyes are large and projecting; the front is stoutly striated in the middle; the front and vertex are depressed; the eyes project above the vertex; the sides of the pronotum are indistinctly toothed at the base; they project at the tegulæ; the scutellar depression is larger and deeper than usual and bears four longitudinal keels.

A distinct genus of *Mesostenini* easily known by the completely reticulated median segment without transverse keels, by the stoutly spined petiole, by the raised scutellum, by the coarsely reticulated thorax, and by the thickened base of the hinder tarsi.

*Acleasa albispina*, sp. nov.

Nigra, scutello spinisque albis; abdomine rufo-balteata; pedibus flavis, coxis, trochanteribus posticis apice femorum posticorum apiceque tibiæ posticarum nigris; alis hyalinis, nervis stigmatique nigro-fuscis ♀.



Long: 12; terebra 2 mm.

Hab. Borneo (Shelford).

Antennæ stout, longer than the body; the ten middle joints white. Head black; the face and clypeus thickly covered with long white hair; the middle is irregularly striated. Clypeus roundly convex, shining; its upper part closely and finely punctured. The middle of the front is stoutly irregularly striated, more closely below than above. Mandibles rufous, their teeth black. Thorax black; the scutellum and teeth yellow. The middle lobe of the mesonotum is closely transversely striated, the lateral are coarsely irregularly reticulated and hollowed down the centre. Scutellum yellow, black at the base; smooth and shining; the basal depression is large; it has two stout complete keels in the centre, and an indistinct one on either side. Post-scutellum smooth and shining; its apex is dilated. Median segment coarsely, closely reticulated; the spines are large, conical and lemon yellow. Pro- and upper half of the mesopleuræ coarsely, irregularly striated; the lower part of the meso- smooth and shining; the furrow is crenulated. Metapleuræ closely reticulated. Legs yellow; the hinder coxæ, except the basal two-thirds above, the trochanters, apical third of femora, the extreme base of the femora and their apex more broadly, black. Abdomen black, the base of the petiole, its apex somewhat more narrowly and the apex of the other segments, yellow; the post-petiole is punctured and striated down the middle; the second, third, and fourth segments are closely punctured.

*Fislistina*, gen. nov.

Post-petiole much widened and clearly separated; its spiracles wider from each other than from the apex. Median segment rugose and reticulated; its sides bearing short thick spines; the spiracles large, oblong. Areolet small, square, open at the apex; the transverse basal nervure is interstitial; the transverse median nervure in hind wings broken below the middle; the stigma narrow, linear; below it, is a wide cloud. Antennæ stout, longish, annulated with white; the third and fourth joints subequal in length. Head as wide as the thorax; almost transverse and not much developed behind the eyes, which are large and parallel; the malar space is small. Clypeus clearly

separated from the face, roundly convex; its apex depressed. Mandibles large, wide; their apex with two equal triangular teeth. Parapsidal furrows extending beyond the middle. Scutellum roundly convex; the basal depression wide and deep. The metapleural furrow is wide, deep and reaches to the apex; there is only a basal keel on the median segment; the legs are stout and of moderate length; the fore tarsi are longer, the four hinder shorter, than the tibiae; the fore tibiae are distinctly narrowed at the base; the claws longish, curved. There are distinct gastrocœli on the second abdominal segment; the apical segment is transverse, bluntly pointed and bears distinct cerci.

Belongs to the *Mesostenini* and is most nearly related to the American genera *Mesostenoides* and *Christolia*.

*Fislistina maculipennis*, sp. nov.

Nigra, abdomine late flavo-balteata; pedibus rufi; tibiis late apiceque femorum posticorum nigris; alis hyalinis, fascia substigmalis fusca ♀.

Long: 10; terebra 2 mm.

Hab. Borneo (Shelford).

Antennæ stout, longer than the body, black with two white bands, one on joints 6-10 and another on joints 13 to 16. Head entirely black; the face rugose, roundly projecting in the middle; the front smooth; the lower part of the vertex stoutly, longitudinally striated. Mesonotum smooth and shining, the furrows appear to bear a silvery pubescence. Scutellum smooth, yellow; the basal depression has four keels. Median segment coarsely reticulated; the basal region in the middle smooth; the teeth are yellow, short and broad. The upper part of the propleuræ closely obliquely striated; the middle less closely and more strongly longitudinally striated; the basal half of the mesopleuræ is closely longitudinally striated; the apical smooth and shining; the metapleuræ coarsely rugose. Wings hyaline, a broad fuscous cloud extends from the stigma to the opposite side. Legs rufous; the tibiae and tarsi paler, the hinder white; the greater part of the four front tarsi, the four front tibiae in front, the apex of the hinder femora, the tibiae, except at the base, and the apical joint of the hinder tarsi,



black. On the abdomen, the post petiole, the apex of the second segments, the band roundly widened backwards in the middle, a broad band on the third segment, widest in the middle, and the greater part of the penultimate segment, yellow. The gastro-cœli are rufous.

*Chrysocryptus*, gen. nov.

Head and thorax densely covered with longish hairs, the abdomen sparsely haired. Median segment areolated; the transverse and longitudinal keels distinct; the areola large, twice longer than wide. Stigma conspicuous, wide, obliquely narrowed towards the base and apex. Areolet large, wide, not much narrowed above, five-angled. Radial cellule wide; the basal abscissa of the radius shorter than the apical and more curved than it; there are no nervelets on the disco-cubital and the recurrent nervures; the transverse basal nervure is interstitial. In the hind wings the cubitus is broken shortly below the middle. Head, if anything, wider than the thorax; the occiput rounded; eyes large, distinctly projecting; the malar space small. The middle lobe of the mesonotum is distinctly separated; the parapsidal furrows are deep and reach near to the scutellum. The furrow at the bottom of the mesopleuræ is distinct. The spiracles are linear; the spiracular area is well defined, as is also the area at its apex. The base of the metanotum is obliquely depressed. Ovipositor projecting; the sheaths are covered with longish white hair. Legs slender; the hinder coxæ and trochanters longish; the fore tarsi are longer than the tibiæ.

The first three joints of the antennæ are much lengthened, being fully four times as long as wide at apex, or longer; the abdomen is twice the length of the head and thorax united; the disco-cubital nervure is roundly curved, not angularly broken; the clypeus is roundly convex; its apex broadly rounded. The face is densely covered with golden hair. The median segment is completely areolated; the areæ are all large and have stout keels; the areola is rounded at the base, transverse at the apex; the lower part of the metapleuræ is stoutly keeled; the radius is thickly pilose at the base; the apical nervures in the hind

wings are faint and incomplete; the second transverse cubital nervure is bullated largely above.

This genus does not fit well into any of the known tribes of the *Cryptina*. The areolated median segment would place it near the *Hemiteini* and the *Phygadenonini*. Characteristic is the densely haired head and thorax.

*Chrysocryptus aureopilosa*, sp. nov.

Niger, capite thoraceque dense aureopilosis; abdomine pedibusque posticis rufo-testaceis; pedibus pallide testaceis; alis hyalinis, apice fumatis; stigmatibus nervisque testaceis ♀.

Long: 12; terebra 4-5 mm.

Hab. Borneo (Shelford).

Antennae rufo-testaceous; the scape paler, and thickly covered with pale testaceous hair. Head black, smooth and shining densely covered with longish bright fulvous hair. Mandibles rufo-testaceous; the teeth black. Legs rufo-testaceous, the anterior paler; the hinder tarsi infuscated; they have the coxae, trochanters and femora covered with long pale hair; the tibiae and tarsi are closely covered with short pubescence. Wings hyaline; the apex infuscated; the basal nervures are dark; the apical, pale testaceous. The apex of the abdomen is pale testaceous, the basal three segments are sparsely covered with long pale hair.

*Latteva*, gen. nov.

Median segment not areolated; the base smooth; the rest striolated; the sides spined. First joint of the flagellum, if anything, shorter than the second. Antennae over twenty-jointed. Eyes large, parallel, reaching close to the eyes; the hinder ocelli are separated from each other by about the same distance they are from the eyes. Pterostigma elongated, narrow; areolet small, square, open at the apex; the transverse median nervure is received behind the transverse basal. Radial cellule elongate, narrow, sharply pointed at the apex, the apical abscissa of the radius is not twice the length of the first; the nervures in the hind wing are complete; the transverse median nervure in the hind wings is sharply angled below the middle where the cubital nervure issues from it. Metathoracic spiracles small, twice longer than wide. Belongs to the *Hemiteini*. The non-

areolated strongly striolated median segment affords a good mark of recognition. The head is distinctly wider than the thorax; the clypeus is clearly separated from the face: the mandibles are large, broad and bi-dentate at the apex; the parapsidal furrows only extend to shortly beyond the middle of the mesonotum; the scutellar depression is deep, wide and keeled; the base of the mesopleurae is keeled; the petiole is longer than the second segment; the post-petiole is distinctly separated.

*Latteva albobalteata*, sp. nov.

Nigra, abdomine albo balteato; pedibus testaceis; tibiis, tarsis trochanteribusque posticis nigris; tibiis posticis albo annulata; alis hyalinis, fusco-bifasciatis ♀.

Long: 8 mm.

Hab. Sarawak, Borneo (Shelford).

Antennae black, the eighth to sixteenth joints white beneath; the scape is brownish on the under side. Head entirely black; the front is keeled down the centre; the keel on either side is oblique. Face opaque, alutaceous; the clypeus bare, smooth and shining. Mandibles black, rufous in the middle. Maxillary palpi white. Thorax black, except the scutellum which is broadly yellow in the middle; the apex of the middle lobe of the mesonotum is rugose. The median segment behind the keel is smooth; the middle is obliquely-longitudinally striated; the apical slope is transversely striated, the striae running into reticulations; the spines are black, longer than broad. The propleurae obliquely striated in the middle; the middle have a plumbeous hue; they are finely striated below the tubercles, stoutly behind the keel, and closely on the upper part of the depression behind the middle; the lower curved keel has some stout keels on the basal half as has also the apical bordering one. Metapleurae striated indistinctly at the base and much more strongly towards the apex; the oblique furrow behind the middle is broad and deep. The anterior four legs are testaceous, paler, more yellowish at the base; their tarsi fuscous; the hinder pair are black; the femora rufous, black at the apex; the coxae are pale rufous; there is a narrow white band near the base of the tibiae; the apical joints

of the tarsi are testaceous at the base. Wings hyaline; there is a fuscous cloud at the stigma extending from the base of the cubitus to the apex of the areolet; there is another fuscous cloud at the apex. Abdomen black; the petiole is rufous, with a broad fuscous band near the base of the post-petiole; the apical third of the second segment and the apical two segments are white.

*Friona*, gen. nov.

Radial cellule elongate. Areolet moderately large (larger than in *Mesostenus*) wider than long; the cubital nervures parallel, straight, not oblique; the second faint; the transverse basal nervure interstitial or nearly so. The transverse median nervure in the hind wings is broken far below the middle. Head wider than the thorax; the front is stoutly striated and is depressed in the middle. Eyes large, parallel; the malar space is moderate. Face short, not extending below the level of the eyes. Clypeus roundly convex, clearly separated behind; its apex broadly rounded; labrum projecting. Mandibles with two large triangular teeth. Thorax more than three times longer than broad; pronotum dilated in front; the parapsidal furrows deep, extending beyond the middle. Median segment elongate, its base smooth; there is a transverse keel near the base, the part beyond it is closely transversely striated; the apex of the segment has a straight, steep slope and projects bluntly at the edges above; the spiracles are small, about three times longer than broad. Legs longish, slender; the fore tarsi are twice the length of the tibiae. Antennae longish, slightly, but distinctly, dilated at the middle; the third joint is longer than the fourth.

Has the usual form and colouration of the *Mesosteni*. The generic distinctions lie in the strongly striated depressed front and the transversely striated median segment, with its steeply sloped, clearly separated apex.

*Friona striolata*, sp. nov.

Nigra, late flavo maculata; mesopleuris fere immaculatis; pedibus fulvis, posticis nigro-maculatis; alis hyalinis nervis stigmatique nigris ♀.

Long: 13-14; terebra 4 mm.

Hab. Sarawak, Borneo (Shelford).

The sixth to seventeenth joints of the antennae are white. Head black; the face, clypeus, labrum, inner orbits to the end of the eyes, the outer from shortly above their middle, almost the basal two-thirds of the mandibles and the palpi, yellow; the front in the centre is strongly obliquely striolated; the face is rugosely punctured. Thorax black; the projecting middle of the pronotum, the tubercles, tegulae, the scutellums, a large raised mark, narrowed on the inner side behind the hind wings, and a small curved mark behind the mesopleural suture, pale yellow. Pro- and mesonotum smooth and shining; the pro- and mesopleurae closely longitudinally striated, the striae becoming weaker somewhat towards the apex. The part of the median segment immediately behind the transverse keel is coarsely aciculated; the rest is closely and distinctly transversely striated; shortly behind the transverse keel a broad yellow band originates, which becomes broadly dilated on the apical slope, where it extends to the middle, its sides being dilated, and the centre rounded. Legs fulvous; the four front coxae and trochanters are pale yellow; the fore femora are lined with black above; the hinder coxae are black, yellow above and at the apex below; the trochanters, apex of femora and of tibiae black; the tarsi white. Abdomen black above; all the segments with their apices yellow, the apical one very narrowly.

*Lactolus*, gen. nov.

Median segment elongate, with one transverse keel; its base smooth, the rest closely transversely striated; its apex has a gradually rounded slope; the keel on the lower part of the metapleurae is complete and is roundly and broadly dilated at the base. Front and vertex depressed, stoutly striated. Areolet of moderate size, longer than broad; the transverse cubital nervures have an oblique slope from the top to the bottom; the apical one is faint; the transverse basal nervure is almost interstitial.

There is only one transverse keel on the median segment; its spiracles are of an elongate oval slope; the clypeus is not separated from the face; the thorax is more than three times longer than



broad; the legs are long; the claws longish; the hinder coxæ are long and reach near to the apex of the petiole; the spiracles on the petiole are separated from each other by about half the distance they are from the apex; the scutellum is stoutly keeled laterally to near the middle, the parapsidal furrows extend to shortly beyond the middle. The ♂ is similar to the ♀; the antennæ are longer and more slender, they are broadly ringed with white in both sexes; the apical abdominal segments in both sexes are marked with white.

The species of the genus are very similarly coloured to *Buodius* with which genus it agrees in some other respects; the difference in the form of the median segment enables them to be separated; in *Buodius* it is shorter, is stoutly spined, and the apex has a straight oblique, not a gradually rounded slope; in *Buodius*, too, the recurrent nervure is received at the apex of the areolet, almost united to the second transverse cubital nervure. Also the median segment is not transversely striated.

*Lactolus albomaculatus*, sp. nov.

Niger, annulo flagello antennarum tarsisque posticis albis; coxis posticis rufis alis fumato-hyalinis, nervis stigmatique nigris ♀.

Long. 13; terebra 3 mm.

Hab. Sarawak, Borneo (Shelford).

Antennæ longer than the body; the sixth to twelfth joints, for the greater part, white. Face rugosely striated in the middle; at the sides the striæ are oblique and more distinctly separated. Clypeus stoutly keeled in the middle; the rest aciculated and irregularly and not very strongly striated. The base of the mandibles closely striated; the teeth are for the greater part rufous. The front and the vertex from the hinder ocelli stoutly striated. Thorax shining; the pro-mesonotum and the base of the median segment smooth, striated. The mesopleural furrow is wide and deep; its lower part is stoutly striated. Legs black; the apical half of the metatarsus, the second and the fourth joints except at the apex, white; all the coxæ and the four front trochanters bright red; the anterior tibiæ and, to a less extent, the femora are brownish. Wings hyaline, with a slight, but distinct, smoky tinge; the stigma and nervures are black; the second transverse cubital nervure

is largely bullated; as is also the cubital-disco, and the recurrent nervures. Abdomen black; the top of the sixth, seventh, and the eighth segment more narrowly above, white; the apex of the second segment is obscure testaceous.

*Lactolus ruficornis*, sp. nov.

Niger, apice metanoti apiceque abdominis albis; pedibus fulvis, trochanteribus tibiisque posticis nigris; tarsis posticis albis; alis hyalinis, nervis stigmatique nigris ♀.

Long: 9-10; terebra 4 mm.

Hab. Sarawak, Borneo (Shelford).

Antennæ as long as the body; the middle of the flagellum is broadly white. The face is rugosely punctured, almost reticulated; the clypeus is smooth and shining, roundly convex; the curved keels on the lower part of the vertex are few in number and stout. Mandibles black; the palpi white. Mesonotum smooth and shining, except on the apex of the middle lobe, which is transversely striated. The scutellar depression is large and is stoutly keeled in the middle; the top of the scutellum is obscure brownish; the post-scutellum is white. The median segment at the base is smooth and shining; the rest of it, from the keel, is closely transversely striated; its apical slope is white; this white band is directed broadly backwards in the middle. The upper half of the propleuræ is closely, longitudinally reticulated, the lower strongly longitudinally striated. Mesopleuræ, except in the middle behind, strongly longitudinally striated; the base is smooth below; the striæ in the middle are smaller and closer; the metapleuræ, from the oblique keel, longitudinally striated; the striæ are waved. Legs fulvous, the anterior paler in tint; the hinder trochanters, the apex of the femora, the tibiæ and the base of the tarsi are black; the rest of the tarsi white. Abdomen black; the apical three segments white; the basal three segments are aciculated.

*Lactolus flavipes*, sp. nov.

Niger, annulo flagello antennarum late, abdominis apice tarsisque posticis albis; pedibus anterioribus flavis; alis hyalinis, nervis stigmatique nigris ♀.

Long: 10 mm.

Hab. Sarawak, Borneo (Shelford).

Antennae black; the apex of the fifth, the sixth to eleventh entirely, and the twelfth and thirteenth partly, white. Head black, the inner orbits in the middle narrowly white; the face rugosely punctured, the punctures running into reticulation above. Clypeus roundly convex, smooth and shining and sparsely covered with longish hair. Mandibles rufous before the middle, smooth; the base coarsely aciculated. Front irregularly striated, coarsely in the centre, more finely in the middle. Pro- and mesonotum with the scutellum smooth and shining; the apex of the middle lobe irregularly longitudinally striated. Median segment behind the keel smooth and shining; the rest of it strongly, transversely, closely striated; on the apex is a curved white band, which is dilated backwards in the middle. Pleurae closely longitudinally striated; the striae on the mesopleurae are more irregular and more or less curved. The four front legs are pale yellow; their coxae black, rufous towards the apex; their tarsi infuscated; the hinder tarsi are white, except narrowly at the base. The basal two segments of the abdomen are aciculated; the others smooth and shining; the second and third segments are narrowly pale at the apex; the apical three are for the greater part white.

*Buodias*, gen. nov.

Thorax three times longer than wide; the median segment behind the keel obliquely rugosely striated; its sides with a broad spine; the apical keel is wanting. Front stoutly striated. Petiole not much longer than the second segment, stout, becoming gradually wider towards the apex from the base. Areolet of moderate size, wider than long, wider at the apex than at the base; the recurrent nervure is received close to the apex; the transverse basal nervure is received behind the transverse basal. The petiolar spiracles are nearer each other than they are to the apex. Scutellum flat, keeled on the basal half. Ptero-stigma elongate, narrow. The median segment is about one half the length of the meso-thorax; its apex has an oblique, straight slope; its spiracles are small, about three times longer than broad. The abdomen is stout, not longer than the

head and thorax united, its apex is blunt and marked with white. The legs are long and stout; the tarsal claws of moderate length; the tibiae are slightly bent at the base. The clypeus is roundly convex, not very clearly separated behind; its apex is transverse and has a distinct margin. The mandibles are broad, curved, bidentate at the apex; lower tooth, small; the base is broadly raised on the upper side, the raised part forming a tubercle-like mass. The head is much wider than the thorax; the metapleural keel is complete and is dilated at the base. On the median segment in the middle behind the keel is an incomplete area, open behind.

In Ashmead's arrangement this genus should come near *Mesostenoides* and *Christolia*.

*Buodius ruficornis*, sp. nov.

Niger; annulo flagello antennarum abdominisque apice albis; coxis trochanteribusque anterioribus rufis; alis fusco-hyalinis, nervis stigmatique nigris ♀.

Long: 21 mm. terebra 4-5 mm.

Hab. Sarawak (Shelford).

Antennae not quite so long as the body, if anything, thickened towards the apex; the sixth to thirteenth joints white beneath; the scape bare smooth and shining. Head entirely black; smooth and shining; the front obliquely stoutly striated below the ocelli; the face coarsely irregularly reticulated. Mandibles black, rufous at the base above. Palpi testaceous. Thorax black; the sides of the scutellum to near the apex white; the spines on the median segment dull white. Pro- and mesonotum shining, bare; the middle lobe aciculated. The scutellum is, if anything, more shining than the mesonotum; post-scutellum is dull white. The base of the median segment in the middle is stoutly keeled; the part behind the keel is aciculated; there is one curved keel on the outer side of the stigma and several on the inner side; the teeth are broad and bluntly rounded at the apex. The middle of the propleurae obliquely, and the upper two-thirds of the apex stoutly, striated. The base and the part in the middle is irregularly reticulated, this part being bounded by a keel in front and by an irregular furrow behind; the lower apical part is irregularly crenulated.

Metapleuræ coarsely obliquely striated; the striae are irregular and more or less wrinkled; the base is aciculated behind the furrow, which is wide and deep; the upper part is irregularly, the median segment from the keel finely and closely transversely striated; the apical slope is rufous. The middle of the propleuræ, the greater part of the meso- and the meta- below the keel are closely and finely longitudinally striated. Legs black; the hinder coxæ rufous; the fore femora and tibiæ more or less testaceous, especially in front, the apical two-thirds of the basal joint of the hinder tarsi, the second, third, fourth and base of the fifth joints, white. Wings fuscous-hyaline; the nervures and stigma black; the second transverse cubital nervure is almost obliterated. Abdomen black; the sixth and seventh segments broadly above and the eighth narrowly white.

*Mesostenus Shelfordi*, sp. nov.

Niger; labro abdominisque apice albis; coxis posterioribus rufis; alis hyalinis, stigmatibus nervisque fuscis ♂.

Long: 9 mm.

Hab. Sarawak, Borneo (Shelford).

Head black; the labrum and palpi white. Mandibles black, whitish-testaceous near the middle. Face opaque, closely but not very distinctly, punctured; the clypeus smooth and shining. front and vertex alutaceous, shagreened; the upper part of the front is furrowed in the centre. Pro- and meso-thorax smooth and shining and with a plumbeous hue. Median segment opaque; above closely, but not very strongly, transversely punctured. Wings clear hyaline; the stigma and nervures fuscous. Legs black; the four posterior coxæ orange red; the front coxæ black, pale at the apex; the four front legs are fuscous; the hinder tarsi are white, except narrowly at the base. The abdominal segments are narrowly lined with dull white at the apex; the apex of the fifth and the sixth and the seventh entirely, are clear white.

This is a *Mesostenus* as defined by Ashmead in his generic revision of the *Ichneumonidae* (Bull. U. S. Nat. Mus. XXIII. 44)—The keel on the mesopleuræ curves broadly and roundly upwards on the apical half; the basal keel on the median segment is complete; the apical one does not reach to the sides; the



keel on the lower edge of the metapleuræ is broad and plate-like; the transverse median nervure is received shortly behind the transverse basal; the areolet is moderately large, about one half longer than broad; the recurrent nervure is received shortly behind the middle.

LISTRODROMINI.

*Maraces*, gen. nov.

Claws pectinated, scutellum flat throughout, its sides and apex keeled. Areola obliquely narrowed behind, open in front, not separated from the posterior median area; it is separated at the base from the lateral area. Antennæ dilated beyond the middle. Labrum hidden. Areolet narrowed at the top, nervure uniting there; the transverse median nervure is widely distant from the basal. The apex of the hind femora reaches to the middle of the fourth segment; the abdominal segments are aciculated.

The pronotum projects above, broadly at the base, more narrowly at the apex; the apex of the scutellum has a perpendicular slope and is clearly raised above the post-scutellum; the abdominal segments do not project much at their apices. The stump of a nervure on the disco-cubital nervure is almost obsolete. Clypeus separated from the face, foveate at the base. Mandibles large; the teeth large, widely separated. Metathoracic spiracles linear, much longer than wide.

The eyes are large and parallel; the malar space is large. There are seven abdominal segments; the ventral keel is on the third and fourth segments. The occiput is roundly incised and is keeled above. Median segment, short, rounded gradually behind.

The pectinated claws refer this genus to the *Listrodromini*. The claws have long teeth and are toothed uniformly to near the apex. There are no spines on the median segment; the spiracles on the first abdominal segment are elongated; the segments are banded with yellow; the ovipositor projects largely.

Of the known genera of *Listrodromini*, *Maraces* comes nearest to *Neotypus* which, among other differences, is readily separated from it by the very small, rounded spiracles of the petiole. If it were not for the pectinated claws the genus might be placed with the *Joppini*.

*Maraces flavo-halteata*, sp. nov.

Niger, late flavo-maculato; pedibus flavis, coxis trochanteribus femoribusque posticis nigris; apice tibiæ posticarum late nigro; alis hyalinis, nervis stigmatique nigris ♀.

Long: 14 mm. ♀

Hab. Sarawak, Borneo (Shelford).

Antennæ black, the middle of the flagellum broadly banded with white. Head black; the face, except for a broad black line in the middle, the clypeus, the inner orbits narrowly to the top of the eyes on the inner side, and the outer, entirely below and broadly above, pale yellow. The face closely, the clypeus sparsely, punctured. Mandibles black. Front and vertex impunctate, bare, shining. Thorax black; the edge of the pronotum, two marks on the mesonotum, obliquely and sharply narrowed at the base, the apex of the scutellum broadly, the mark narrowed behind the apical part of the scutellar keels, the post-scutellum, two marks on the apex of the median segment, narrowed below as they follow the outline of the lateral areæ, the lower part of the propleuræ, of the mesopleuræ more broadly, the tubercles, the hinder edge of the mesopleuræ, and the apical half of the metapleuræ, yellow. Mesonotum closely rugosely punctured, reticulated in the middle behind; the scutellum is similarly punctured. The base of the median segment is smooth; the areola is coarsely sharpened; the posterior area coarsely irregularly reticulated; the lateral area smooth at the base, the rest coarsely punctured; the spiracular area rugose, the apex transversely, coarsely striated. The upper part of the propleuræ is closely punctured, the apex stoutly striated, the striæ in the centre extending to the centre. Meso- and metapleuræ distinctly and closely punctured; the middle of the former finely and closely longitudinally striated. Wings hyaline, the nervures and stigma black. The four front legs yellow; the femora, tibiæ and tarsi black behind; the hinder coxæ, except at the apex on the inner side, the basal joint of the trochanters, the femora and the apical third of the tibiæ, black; the rest yellow. Abdomen black; the base of the petiole broadly, its apex and the apex of all the other segments, yellow; the middle segments of the abdomen are closely punctured; the gastrocoeli are

yellow, the steep apex of a more rufous hue; the base of the segment between them is striated; the sides of the apical three segments are yellow, the yellow becoming gradually broader towards the apex.

*Maraces pectinata*, sp. nov.

(Niger, late flavo ornato; pedibus fulvis, coxis trochanteribusque anterioribus flavis, posticis nigris; alis fulvo-hyalinis, nervis, stigmatique nigris ♀.

Long: 17 mm.

Hab. Khasia Hills (Coll Rothney).

Antennæ black, the eighth to sixteenth joints white; the scape covered with white hair. Head black; the face, clypeus, the inner orbits,—narrowly below, more broadly above, the yellow not extending beyond the inner top of the eyes,—and the outer orbits entirely, from above the middle of the eyes to the base of the mandibles, pale yellow. The face and clypeus obscurely punctured and thickly covered with white hair; there is a black line down the face and an elongate mark on the apex of the clypeus. Front and vertex smooth, shining, and bare. Mandibles yellow, the teeth black. Thorax black; the edge of the pronotum, two lines on the mesonotum, obliquely narrowed on the inner side at the base, the apical half of the scutellum, the mark roundly narrowed at the base, the base of the pronotum, the lower side of the propleuræ from behind the middle to the apex, the tubercles, a small mark on the middle of the mesopleuræ, a smaller one behind it lower down, the lower third of the mesopleuræ, the apex of the mesopleuræ broadly below the keel, yellow. Legs fulvous; the four front coxæ, trochanters pale yellow; the hinder coxæ black on the outer side and on the outer half of the top; the basal joint of the trochanters black. wings hyaline, the base with a slight fulvous tinge; the stigma and nervures black; the areolet oblique; the second transverse cubital nervure longer and with a more oblique slope than the first; they almost touch above; the recurrent nervure is received shortly behind the middle. Abdomen black, the apices of all the segments yellow; the band on the third is interrupted in the middle; the petiole shining, the base of the post-petiole strongly punctured: the second, third, fourth segments closely

punctured; the gastrocoeli narrow, deep, smooth, and shining. Mesonotum rather strongly and closely punctured; the scutellum is as strongly, and more widely punctured; its sides, under the keels, strongly but not closely, punctured. The base of the median segment is smooth; the rest coarsely punctured, the apex especially in the middle, closely reticulated; the supramedian area large, about as wide as long; the sides at the base obliquely narrowed, the middle straight, the apex is not clearly separated from the posterior median by a keel. Pro- and mesopleuræ smooth; the depression on the former stoutly striated; the lower half of the meso- is depressed and separated from the raised upper half; the meta- closely and strongly punctured. The median segment is thickly covered with white hair.)

*Joppini Zonojoppa*, gen. nov.

Antennæ short, distinctly dilated and compressed between the middle and the apex; the dilated joints hollowed laterally. Wings violaceous throughout; the areolet is narrowed at the top, the transverse cubital nervures almost uniting there; the transverse basal nervure interstitial. Scutellum roundly convex; not raised above the level of the mesonotum, its sides stoutly keeled. Areola widely separated from the base of the segment, rounded and narrowed behind, the basal half deeply hollowed, the lateral basal areæ clearly separated. Clypeus broadly rounded at the apex, the labrum hidden. Legs short; the apex of the hinder femora not extending beyond the apex of the third segment. The abdominal segments do not project much laterally at the apex; the second and third segments are longitudinally striated at the base, the last (seventh) segment is well developed; its cerci are much longer than usual.

The thorax is shorter than the basal three segments of the abdomen; the middle of the mesonotum is raised and separated in front; the post-scutellum is shortly striated and depressed laterally; the apical three areæ on the median segment are closely defined, as is also the spiracular area; the sides of the mesonotum are bordered by a wide deep furrow; there is a short stump of a nervure on the disco-cubital nervure.

The characteristic features of this genus are the violaceous wings, the stoutly keeled scutellum, and the excavated areola.

*Zonojoppa violaceipennis*, sp. nov.

Nigra, capite thoraceque flavo maculatis; abdominis basi late rufo; pedibus nigris; coxis trochanteribusque anterioribus flavis, alis violaciis, nervis stigmatique nigris ♀.

Long. 15 mm.

Hab. Sarawak (Shelford).

Antennæ black, the scape yellow beneath. Head black; the face, clypeus, base of mandibles, the inner orbits to shortly beyond the ocelli, the outer more broadly from near the top, the line becoming gradually wider from top to bottom, pale yellow. Front and vertex smooth, bare and shining; the black on them has a plumbeous hue, and they are sparsely covered with pale hair. Thorax black; the upper edge of the pronotum from near the base (the yellow with a black band in the middle), the keels of the scutellum from near the base, the apex of the post-scutellum; the base of the prothorax from the keel on the pleuræ, the tubercles and a large mark on the lower part of the mesopleuræ at the base, yellow. Mesonotum in the middle stoutly punctured; the punctured space prolonged laterally at the base; the sides near the tegulæ are deeply furrowed. The scutellum, except at the base, is irregularly and rather strongly punctured; the sides are stoutly keeled; in the centre of the post scutellum are four stout keels. The central basal depression of the median segment is smooth; the sides are strongly punctured; the areola has a large, round depression at the base, which extends to shortly beyond the middle; the apical central area is smooth and depressed at the base; the rest of it stoutly transversely striated; the lateral stoutly, irregularly striated, almost reticulated; the spiracular, beyond the spiracles, irregularly obliquely striated. The lower part of the propleuræ is aciculated and irregularly striated; the meso-, except behind, finely and closely punctured; the meta- closely and coarsely striated; the meso- and metapleuræ are thickly covered with white hair. The four anterior coxæ entirely, the trochanters, the femora, except at the base, and the tibiæ and tarsi in front, pale yellow, black behind; the legs black; the coxæ above, except at the base, the apical half of the trochanters, a line on the femora above and on the base of the femora, pale yellow; all the legs are thickly covered with pale



pubescence; all the calcaria are pale yellow. Wings uniformly violaceous; the transverse cubital and the recurrent nervures are largely bullated. The basal three abdominal segments and the base of the fourth broadly, ferruginous; the basal three segments are narrowly lined with yellow at the apex. The post-petiole is finely longitudinally striated, the sides punctured; the second and third segments have a narrow keel in the centre, bordered by some longitudinal striations; the gastraceli are large, smooth, and have two oblique stout keels on the outer side.

## MUTILLIDÆ.

*Mutilla herpa*, sp. nov.

Nigra, pro-mesothorace scutelloque ferrugineis; abdomine nitido, dense nigro piloso; segment secundodo dense albo piloso; alis violaceis; tegulis rufis ♂.

Long: 12 mm.

Hab. Sarawak, Borneo (Shelford).

Head as wide as the base of the mesonotum; coarsely rugosely punctured, running into reticulations on the front, which, at the apex, broadly projects; its apex and side are sharply keeled; the middle is obliquely incised; the sides are broadly rounded; the face is rugose and bears, on the middle at the apex, three irregular punctures. Antennæ black, the scape shining, pilose; the flagellum opaque, bare. Pro- and mesonotum, with the scutellum, closely rugosely punctured. The scutellum, is broad; its sides are smooth and project; its apex has a rounded slope. Wings violaceous, lighter in tint at the base; the third transverse cubital nervure is only indicated by a stump on the top; there being thus only two complete cubital cellules; the second transverse cubital nervure is broadly rounded. The median segment is coarsely reticulated; the basal three are of equal length, but the central is much narrower and is acutely pointed at the apex. The pro- and the upper two-thirds of the mesopleuræ are ferruginous; the propleuræ and the base and apex of the mesopleuræ are smooth. Abdomen deep black, shining; the petiole is broad and becomes gradually

wider towards the apex; the apex is smooth; at the base of this smooth part is a row of large punctures; from this the petiole slopes obliquely to the base; the ventral keel is straight, rounded at the base and apex; the second segment is covered with short depressed clear white pubescence; and is smooth and shining in the middle at the apex; the pygidium bears large round punctures all over. Legs black, thickly covered with long white hair; the spurs white.

*Mutilla ira*, sp. nov.

Nigra, dense albo piloso; alis fusco-violaceis, basi fere hyalinis ♂.

Long: 17 mm.

Hab. Sarawak, Borneo (Shelford).

Head distinctly narrower than the thorax; closely rugosely punctured and thickly covered with long white pubescence; the vertex and front with the hair sparser and shorter. The clypeus is smooth and shining and is keeled in the middle; the mandibles, at the base, are thickly covered with long white hair. Thorax densely covered with longish grey pubescence; the mesonotum is strongly, distinctly and uniformly punctured; the furrows are distinct on the apical half. Scutellum strongly, deeply and uniformly punctured and roundly convex; the post-scutellum opaque, coarsely aciculated. Median segment coarsely reticulated; the basal median reticulation is twice longer than broad and has the apical half abruptly narrowed. Mesopleuræ coarsely punctured in the middle and thickly covered with grey hair; the lower part of the metapleuræ is alutaceous, the upper punctured. Legs thickly covered with long white hair; the calcaria pale. Wings dark violaceous, paler at the base; the cubital cellules complete; the middle one, is, above and below, longer than the following. Abdomen black, the basal two segments, the basal half of the third rufous; the base of the petiole is broad, more than half the width of the apex; the ventral keel is broadly rounded; the hair is white on the basal segments, shorter and black on the apical two; the hypopygium is punctured, smooth and shining in the middle; there are no keels or furrows on the epipygium.

*Mutilla olbia*, sp. nov.

Black, densely covered with longish pale hair; the first, second and the base of the third abdominal segment red; wings fuscous-hyaline with a violaceous tinge; the stigma and nervures testaceous, ♂.

Long: 15 mm.

Hab. Penrissen, 4500 feet, Sarawak.

Antennae densely covered with a pale pile; the second and third joints together are equal in length to the fourth. Head distinctly narrower than the thorax; roundly, obliquely narrowed behind the eyes; the vertex strongly but not very closely, punctured; the front more closely rugosely punctured; the vertex sparsely, the front more thickly covered with long fulvous hair. Clypeus depressed in the middle; the edges rounded and forming a semicircle; the apical tooth of the mandibles is long, rounded at the apex; the subapical one is short and blunt. Thorax thickly covered with long fulvous hair; the pro- and mesonotum closely rugosely punctured; the scutellum is more closely punctured. Median segment closely reticulated; the basal three central areas larger than the others; the central is longer than the others. Propleurae at the base rugosely punctured; the lower part of the apex with five stout keels; the central, raised part of the mesopleurae is punctured but not deeply or strongly; the base of the metapleurae smooth, the apex reticulated. The third transverse cubital nervure and the second recurrent are faint; the first transverse cubital nervure is oblique and rounded; the second is roundly curved and not obliquely sloped; the second cubital cellule at the top is shorter, at the bottom longer than the third: the recurrent nervures are received near the base of the apical third of the cellules. Legs thickly covered with white hair; the spines and calcaria white. On the abdomen the first, second, and the base of the third segments are rufous; the basal five segments are covered with long pale, the apical with black, hair; the ventral keel is slightly dilated at the base, roundly narrowed at the apex; the last segment above has the apical two-thirds broadly smooth in the middle; below it is strongly punctured, except at the apex, which is smooth and rufous.

*Mutilla bagrada*, sp. nov.

Long: 16-17 mm. ♂.

Hab. Kuching, Sarawak.

This species comes very near to *M. casiphia*; the differences between the two may be expressed thus:—

Scutellum not furrowed down the middle; the keel on the petiole straight; the face not tuberculate; the propleuræ smooth except above. *bagrada*.

Scutellum furrowed down the middle; the keel on the petiole curved; the face tuberculate below; the propleuræ rugose. *casiphia*.

Flagellum of antennæ brownish beneath, the third and fourth joints equal in length. Front and vertex coarsely rugosely punctured; the punctures running into reticulations on the front; the apex of the tubercles rufous. Clypeus slightly depressed in the middle; smooth, shining; the apex transverse. The face thickly covered with long grey hair. Mandibles at the base thickly covered with grey hair; the subapical tooth is indistinct; the apex of the projection behind the middle is oblique, rounded on the lower part. Pro- and mesonotum closely and rugosely punctured; there is a smooth keel in the middle; there is a furrow on either side of it on the apical half. Scutellum roundly convex, coarsely rugosely punctured. Median segment reticulated; the basal area short, not reaching beyond the middle; its basal third widened. Propleuræ smooth; the upper part at the base bordered by an oblique keel. Mesopleuræ thickly covered with silvery pubescence, the base smooth; there is a wide oblique depression. Legs thickly covered with white hair; the calcaria pale. Wings fuscous-violaceous, paler at the base; the third cubital cellule at the top and bottom distinctly shorter than the second; the first transverse cubital nervure is almost straight, and oblique; the second is roundly bent outwardly in the middle; both the recurrent nervures are received shortly beyond the middle. Abdomen thickly covered with white hair; the hair on the apical two segments is black; the basal two segments and the base of the third are rufous; the ventral keel is roundly curved and narrowed at the base; the last dorsal segment has no area; its lower surface is flat.

*Mutilla agapeta*, sp. nov.

Black; the abdomen red, with the apical two segments black; the clypeus broadly convex in the middle; the centre of the scutellum smooth and shining, its base depressed; wings fusco-violaceous, hyaline at the base. ♂.

Long: 15 mm.

Hab. Kuching, Sarawak.

The third joint of the antennæ is shortly, but distinctly, longer than the fourth. Front and vertex closely punctured; the vertex sparsely covered with longish pale hair; the front, especially in the centre, thickly covered with silvery pubescence. Face roundly convex, smooth and shining; the clypeus short, depressed, clearly separated, slightly and broadly waved; the edges depressed. Mandibles densely pilose at the base; the lower basal tooth stout, obliquely directed downwards; there is no distinct subapical tooth; the palpi are black. Pro- and mesonotum closely punctured; the pronotum above thickly covered with griseous pubescence; the lower and hinder part of the propleuræ bear stout, clearly separated keels. The raised central part of the mesopleuræ is thickly covered with silvery pubescence and punctured but not very deeply or closely. Metapleuræ smooth, irregularly reticulated at the base above. Mesonotum closely punctured. Scutellum strongly convex; the basal and apical slopes oblique; the base in the centre is flat, smooth and shining. Median segment reticulated, thickly covered with silvery pubescence. Legs thickly covered with white pubescence; the calcaria and spines pale. Wings fusco-violaceous, hyaline behind the transverse basal nervure; the third cubital cellule at the top and bottom distinctly shorter than the second; the recurrent nervures are received beyond the middle of the cellules; the second transverse cubital nervure is roundly curved outwardly. Abdomen ferruginous; the basal half of the petiole below and the apical two segments black. The ventral keel is almost straight; the last segment is broadly smooth and bare in the centre; below it has the sides broadly, obliquely depressed and clearly separated from the centre which is depressed, especially at the apex, where it is bounded by keels.



*Mutilla ilerda*, sp. nov.

Black; the prothorax, mesonotum, scutellum and base of median segment rufous; abdomen black, with violet and purple tints; the apex of the second and of the third segments banded with clear white pubescence; wings fuscous-violaceous, lighter in tint at the base. ♂.

Long: 12 mm.

Hab. Kuching, Sarawak.

Antennae stout, covered with a microscopic pale pile; the third joint is about twice the length of the pedicle and not quite one-half the length of the fourth. Head distinctly narrower than the thorax, thickly covered with long white hair, except on the front and vertex where the hair is sparser, shorter and black. The ocellar region is bounded laterally and below by two stout keels; the space between is depressed; a keel runs into the front ocellus. The clypeus is broadly keeled above; its apex is transverse, with the sides oblique. Mandibles irregularly bidentate at the apex; their base sparsely covered with longish golden hair. The pro- and mesonotum are closely and strongly punctured and covered with golden pubescence. Median segment closely reticulated; the central basal area is twice longer than broad and has the apical half narrowed. Propleurae closely and strongly punctured except behind; the lower part is bounded by a stout keel: above this, on the apex, are four short keels, which become gradually shorter from the bottom to the top. Mesopleurae closely and strongly punctured. Metapleurae reticulated, except behind; on the upper part, at the base, is a narrow keel and above the middle a wide longitudinal furrow. Wings fuscous-violaceous, paler at the base; the third transverse cubital nervure is faint, as is also the cubitus from the second transverse cubital and the second recurrent nervures; the first transverse cubital nervure is oblique, and roundly curved on the lower part; the second is roundly curved outwardly in the middle; the second cubital nervure above is slightly more than one half of the length of the first. Abdomen shining, black; the third and following segments with blue and violet tints; the basal segments sparsely covered with white, the apical more thickly with black, hair; the apex of the second

and of the third with a broad band of depressed clear white pubescence; the basal ventral segment has a straight, rounded keel in the centre; its sides are stoutly punctured; its apex has an oblique slope; the pygidium is closely punctured and covered with black hair; its apex is depressed; there is no defined area on it; the epipygium is flat, closely and strongly punctured and has its sides margined.

Comes close to *M. gracillima*, Sm.

*Mutilla mamblia*, sp. nov.

Black, the scape of the antennæ, the thorax and the femora rufous, two oval spots of silvery pubescence on the base of the second abdominal segment; the whole of the third segment covered with depressed silvery pubescence; the sides of the pygidium fringed with silvery hair. ♀.

Long: 11 mm.

Hab. Kuching, Sarawak.

This species comes near to *M. prosperpina* Sm. which differs from it in having the legs ferruginous except that the knees and tarsi are slightly fuscous; the pubescence on the thorax is reddish-brown. The present species comes close to *M. gispa* Cam. but, apart from the difference in colouration, it may be known from it by the perfectly smooth pygidium.

Scape of antennæ rufous, covered with pale fulvous hair; the flagellum black, stout; the third joint twice the length of the fourth which is shorter than the fifth. Head nearly as wide as the thorax; closely rugosely punctured: the punctures longer than broad; the antennal tubercles black. Face and clypeus smooth and shining, sparsely covered with long pale fulvous hair. Mandibles rufous, black at the apex; the palpi blackish fuscous, darkest at the base; the subapical tooth straight and oblique at the apex. Thorax twice longer than broad, slightly narrowed in the middle; the base rounded with the edge irregular; the apex transverse, the sides above rounded; the sides of the median segment sharply denticulate; the outer edge of the pronotum is stoutly keeled above; the pleuræ smooth, impunctate; the upper part of the thorax is covered with longish black hairs. Legs black; all the coxæ, trochan-

ters and femora, except at the apex, rufous; they are covered with longish white hair; the spines on the four front tibiae are rufous, on the posterior black; on the tarsi they are rufous, and their basal joints are thickly covered with rufous, stiff pubescence. The basal segment of the abdomen is short and is much narrower than the second; underneath it is rufous, smooth below; the base of the keel obtusely dentate. On the base of the second segment are two irregular, broader than long, marks of silvery pubescence; the third segment is almost entirely covered with silvery pubescence; the pygidium is smooth and is fringed laterally with long silvery pubescence; the ventral segments are thickly covered with silvery hair.

*Mutilla paluca*, sp. nov.

Antennae and head black; the thorax red; the abdomen blue, thickly covered with long white hair, without any bands of depressed pubescence; wings uniformly fuscous, with a slight violaceous tinge; the third transverse cubital completely, and the second recurrent nervure almost completely obliterated ♂.

Long: 9 mm.

Hab. Kuching, Sarawak.

Antennae stout, black, covered with a pale pile; the third joint is slightly, but distinctly, shorter in length than the fourth. Head black, nearly as wide as the thorax; behind transverse, the edge of the occiput sharp and slightly raised above. Front and vertex shining closely punctured all over and covered with longish white hair. Clypeus largely depressed in the centre; the depression largest below, narrowed above; the apex raised and closely punctured. Mandibles bidentate; the apical tooth long and curved at the apex. Thorax entirely rufous, thickly pilose; the hair on the mesonotum fulvous, on the median segment longer and white. Pro- and mesonotum, with the scutellum closely rugosely punctured; the post-scutellum is bordered laterally by a keel and there is a short, less distinct, keel in the centre. Median segment reticulated; the central basal one is very broad at the base, the apex much narrowed; the areas surrounding it are large. The upper part of the propleurae is irregularly, indistinctly, punctured; at the apex is an elongated area, rounded below, which reaches to shortly below the middle.

Mesopleuræ closely punctured; the metapleuræ reticulated, smooth at the base. Legs black; the calcaria and spines pale; the hair dense, long and white. The second recurrent nervure is narrowed at the top, being there as wide as the space bounded by the first recurrent and the second transverse cubital nervures; the first transverse cubital nervure is straight and oblique; the second is curved and only slightly oblique. The first abdominal segment is broad at the base; below it is flat; its central keel does not project much and the part bordering it is irregularly punctured on either side of it. The last segment above is closely punctured, except for a smooth space in the middle at the apex; below, the apical half is stoutly keeled along the sides.

Agrees closely in colouration with *M. ilerda*, which may easily be separated from it by the keels on the front and vertex.

## SCOLIIDÆ.

*Scolia (Triscolia) aglana*, sp. nov.

Long: 12 mm. ♂

Hab. Sarawak (Shelford).

This species is not unlike what I take to be *S. opalina* Sm., which has also been taken in Borneo by Mr. Shelford. The difference between the two may be expressed thus:—

The frontal area clearly defined bounded by a ridge behind, the hair on the head and thorax black; the median segment punctured throughout. *opalina*, Sm.

The frontal area not clearly defined, not bounded by a ridge behind; the hair on the head and thorax white; the median segment not punctured throughout, there being a wide smooth space on the inner side of the lateral lobe.

*aglana*, sp. nov.

Antennæ opaque, bare. Head strongly and closely punctured and thickly covered with long white hair; the middle of the clypeus smooth impunctate; there is no defined frontal area. Mesonotum closely and rather strongly punctured, less closely in the middle. The scutellum and post-scutellum are similarly punctured. The central region of the median segment is bounded by a distinct deep furrow and is punctured, if anything, more strongly than the mesonotum; the inner half of the outer lobe

is smooth and impunctate, the outer punctured. The pro- and basal part of the mesopleuræ, are closely punctured; the apical part is smooth in the middle; with a punctured band above and a wider one below. The metapleuræ smooth, with a punctured band round the top; the punctures are smaller than on the mesopleuræ. Wings fuscous, with a distinct, violaceous tinge and highly iridescent. Abdomen black, with a distinct violet iridescence; the hair is black above, white below; the punctuation is distinct. Legs black; the hair is long and white; the fore calcaria are pale; the spines on the fore tarsi rufous.

*Scolia (Discolia) ergenna*, sp. nov.

Black; the greater part of the clypeus, the pronotum broadly, the scutellum, except at the apex, the post-scutellum, the sides of the metanotum and the apex of the metapleuræ broadly, lemon-yellow, as is also the upper part of the mesopleuræ at the base; abdomen broadly banded with yellow; legs black; the four anterior tibiæ lined with yellow; wings hyaline; the radial cellules infuscated, the stigma and nervures dark rufous ♂.

Long: 13 mm.

Hab. Pankalan Ampat, Sarawak.

Antennæ black; the scape covered with white hair. Head thickly covered with long soft white hair. Except immediately below the ocelli, the vertex is closely punctured; the front ocellus is larger than the hinder pair and is placed in a deep pit; except above, the front is closely and strongly punctured, and has an oblique slope. The face has a smooth, flat keel in the middle and is sparsely punctured; the clypeus is roundly convex, sparsely, and distinctly, punctured; it is yellow, except at the apex, where there is a black line, which is roundly dilated above. The yellow bands on the thorax are broad and of equal breadth throughout; they are united above by a narrow yellow line on the hinder edge of the pronotum. Mesonotum thickly covered with short fuscous hair; the scutellum with longer paler hair. The median segment is thickly covered with long soft white hair; the black central part is depressed; the sides are broadly rounded and project slightly. Mesopleuræ thickly covered with long pale hair; the pro- and metapleuræ shortly pilose. Legs thickly covered with white soft hair; the calcaria black. Wings



hyaline; the radial cellules infuscated; the nervures dark rufous; the second transverse cubital nervure is broadly rounded above. Abdomen thickly covered with white hair; the apices of the basal three segments are broadly yellow; the black on the basal segment is triangularly produced in the middle; on the second it is squarely produced, the dilated part being more broadly and more distinctly separated; on the third the black band becomes gradually narrowed towards the apex; on the fourth and fifth the black bands are not dilated and extend to the middle; the apical three segments are entirely black.

*Scolia (Discolia) patara*, sp. nov.

Long: 17-19 mm.

Hab. Santubong, Sarawak.

This species comes very near to *D. thyatira* Cam. but the two are, I consider, distinct. *D. patara* is smaller, it wants the curved yellow marks on the top of the clypeus, there is no yellow mark below the antennæ, and the lower part of the radius is broadly rounded outwardly and does not form an angle with the upper absissa.

Head black, the front, vertex and the upper half of the outer orbits narrowly orange-yellow; the front and vertex thickly covered with long pale fulvous hair; the face more sparsely with long black hair. The front and vertex strongly, the face, if anything, more strongly punctured, but not quite so closely; the clypeus is almost impunctate; the occiput is thickly covered with black hair. The orange band on the pronotum is narrowed behind, is broad, and covered with fulvous hair; the mesonotum is sparsely punctured and is thickly covered with short black hair; the scutellum is covered with long black hair except at the apex; the post scutellum is much more sparsely haired. The median segment is thickly covered with long black hair; as are also the pleuræ; the metapleuræ have also a pale pubescence. Wings uniformly fuscous-violet and moderately iridescent. Abdomen, except in the middle, thickly covered with black hair, smooth, shining, and, especially on the middle segments, bearing brilliant blue and violet tints, this being also the case with the ventral surface.

*Scolia (Discolia) acutinerva*, sp. nov.

Black; the apices of the basal four abdominal segments lined with yellow; the wings yellowish-hyaline, the cubital cellules with a more decided yellow tinge than the rest; the head and thorax covered with a pale golden pile and thickly with pale fulvous hair; the basal three segments of the abdomen have blue and violet tints and are fringed with pale fulvous hair; the hair on the apical segment is long, dense and black ♀.

Long: 23 mm.

Hab. Borneo.

Antennæ black, the scape shining and sparsely covered with long pale fulvous hair. The head, except on the ocellar region, is thickly covered with long pale fulvous hair; the vertex is more sparsely covered than the front; the vertex is shining and is strongly, but not closely, punctured; the front is impunctate and is furrowed down the middle. The clypeus is fringed above with long fulvous hair, is smooth above, the apex is irregularly, stoutly, longitudinally, striated; the extreme apex is depressed, smooth, and more or less piceous. The apices of the mandibles are piceous. The mesonotum is strongly punctured, except in the middle behind; the scutellum is, if anything, more strongly and closely punctured, except at the apex, which is smooth; the post-scutellum is more closely punctured. The golden pile on the median segment is dense, except laterally at the base, it is closely punctured. The golden pile on the pleuræ is very dense. Legs black, covered with fulvous hair. The long spines on the front tarsi are bright rufous; on the four hinder they are of a paler rufous colour; the tibial spines are pale yellowish; the calcaria are of a still paler yellow colour. The malar nervures are rufous; the transverse cubital nervure is sharply bent outwardly in the middle and projects there in a short branch. The abdominal segments are smooth, impunctate and are sparsely covered with long pale fulvous hair; the micaceous tints on the basal three segments are very distinct in certain lights; the hairs on the hypopygium are stout, stiff and black.

The clypeus is subtriangular and is broadly, roundly convex; its apex in the middle is transverse, its sides broadly rounded.

Comes near to *S. indica* Sauss. sec. Bingh. Characteristic is the peculiar form of the transverse cubital nervure.

*Dielis borneana*, sp. nov.

Black; the second and third segments broadly, and the others narrowly on the sides, red; the wings fuscous-violaceous, the apex without a violaceous tinge; the pile on the pygidium golden or rufous; the middle and apical segments of the abdomen fringed with rufous hair.

Long: 45 mm.

Hab. Bajong and Santubong, Sarawak.

Head; the vertex sparsely punctured; the ocellar region more sparsely punctured than the rest; the front is much more closely and strongly punctured and there is a smooth line down the middle. The face and clypeus closely punctured, except for a somewhat triangular large smooth space on the centre of the latter. The occiput is thickly covered with long black stiff hair; the vertex is almost bare; the front is covered with black hair, which has a rufous tinge; the face and clypeus are covered with shorter hair of a darker colour; the sides of the face are thickly covered above with silvery pubescence; the hinder orbits are covered with black hair and with silvery pubescence. Mesonotum strongly and closely punctured except for a smooth impunctate space behind the middle. The scutellum has a punctured, irregular band on the base and an irregular row of punctures before the apex. The post-scutellum is punctured at the base and there is an irregular row of punctures at the apex. The basal region of the median segment is closely and distinctly punctured, except broadly laterally at the base, and more narrowly down the centre; the apical slope is smooth, closely, minutely punctured above and at the sides. Propleuræ closely and rather strongly punctured, except behind; the meso-smooth; the middle thickly covered with black hair; the metapleuræ smooth and almost bare. Legs thickly covered with black hair; the hair on the hinder tarsi bright rufous. Wings fuscous-violaceous; the violaceous tinge absent from the apical portions, which are also lighter in tint; the stigma and nervures black, abdomen black; there are two large rufous marks on the second segment which are narrowed and rounded on the inner side;

the third segment is almost entirely rufous; the fourth and fifth segments are more or less rufous laterally; the apical fringe on the second and following segments is bright rufous; the pygidium is thickly covered with pale golden pubescence, which probably varies in tint.

Comes near to *E. luctuosa* Sm. and *E. 4 guttulata* Burm., but has the abdominal markings red, not yellow. *E. luctuosa* differs from it further in having the wings darker, more uniformly blue-violaceous in colour, in the scutellum and post-scutellum being much more strongly and broadly punctured, the punctuation on the former extending to near the apex, while the latter is strongly punctured at the base and apex.

POMPIDÆ.

*Salix sostratus*, sp. nov.

Black; the antennæ, head, pro- and mesonotum, with the scutellum ferruginous, and except the antennæ, thickly covered with golden pubescence; the legs entirely ferruginous; wings entirely flavo-hyaline, the stigma and nervures fulvous ♀.

Long: 22 mm.

Hab. Sarawak, Borneo (Shelford).

Antennæ bare, uniformly ferruginous. Head ferruginous, densely covered with golden pubescence, the lower outer orbits black. The apex of the clypeus is depressed, clearly separated, smooth, bare and broadly rounded; the sides straight and oblique. The apex of the mandibles black, the rest ferruginous; the palpi ferruginous. The eyes distinctly converge above where they are separated by not much more than the length of the fourth antennal joint; the hinder ocelli are separate from the eyes by a slightly greater distance than they are from each other. Thorax black, the hinder half of the pronotum, the mesonotum and the scutellum ferruginous; and the whole is covered with a golden pile. The pronotum is furrowed in the middle: the mesonotum is broadly rounded at the base; it is alutaceous. The scutellum is flat, smooth and is not much raised above the top of the post-scutellum; the latter is broadly rounded from the top to the bottom; the sides of both have a distinct oblique slope (and more particularly the post-scutellum) so that both are narrowed on the top. The part at the sides of

the post-scutellum is strongly, but not closely, transversely striated. The median segment has a gradually rounded slope to the apex; the base and apex are smooth; the rest coarsely, irregularly transversely striated. Wings uniformly yellowish-hyaline, the apex not infuscated; if anything paler than the rest of the wing; the stigma and nervures yellowish; the first transverse cubital nervure is obliquely curved; the upper (longer) part has a more sharply oblique slope than the lower; the second is straight and oblique; the third is broadly rounded; the first recurrent nervure is received near the base of the apical third of the cellule, not close to the second transverse cubital nervure as in *Mygnumia*; the first transverse cubital cellule is distinctly longer than the second above, but slightly shorter below; the second recurrent nervure is received at the apex of the apical fourth of the cellule. Legs uniformly ferruginous. Abdomen black; the last segment rufous all round and thickly covered with long rufous hair; the penultimate segment is covered with a golden pile.

This is a much more slenderly built species than *S. flavus*; and may be readily separated from it by the cubital cellules being more equal in length, by the pronotum not bulging broadly outwardly in the middle, not narrowed at the apex, by the median segment having a more gradually rounded slope and the head is shorter and more obliquely narrowed behind the eyes. Characteristic, as compared with most of the species of the *flavus* group, is the fact that the coxæ and trochanters are not black.

*Salius iobates*, sp. nov.

Claws with one tooth. Black, the abdomen with a bluish tinge; the antennæ rufous yellow, the scape and the apical four joints black; the basal half of the wings fuscous-violaceous; the apical yellowish-hyaline; the hinder wings entirely smoky violaceous. ♀.

Long: 24 mm.

Hab. Kuching, Sarawak.

Head black; the part between and below the antennæ testaceous; the clypeus with a brownish tinge; its apex rufo-testaceous. Front and vertex alutaceous; the front distinctly



furrowed down the centre. Eyes distinctly converging above; at the top separated from each other by the length of the fourth antennal joint; the hinder ocelli are separated from each other by a less distance than they are from the eyes. Thorax velvety black, sparsely covered with long black hair; the scutellum not projecting much over the mesonotum; the post-scutellum has a lower elevation than it; its apical two-thirds have an oblique, straight slope; its centre is keeled. Median segment obscurely transversely striated. The dark part of the wings extends close to the first transverse cubital nervure and on the lower edge near to the apex; the base of the cubital cellule is blackish also; the second cubital cellule is distinctly shorter than the third above and below; the third transverse cubital nervure is obliquely narrowed towards the second on the upper half; the accessory nervure in the hind wings is interstitial. Legs black; the calcaria and spines black; the tooth on the base of the claws short and bluntly pointed. Abdomen black, with a distinct plumbeous-sericeous tinge; the anal segment thickly covered with long black hairs.

*Macromeris aureopilosa*, sp. nov.

Nigra, antennis subtus brunneis; capite thoraceque dense aureopilosis; alis flavo-hyalinis, apice fusco-violaceo, ♀.

Long: 13 mm.

Hab. Borneo.

Antennae slender, black above, brown below. Head densely covered with depressed golden pubescence and more sparsely with long silvery hair. Apex of clypeus broadly rounded. Mandibles black, broadly rufous near the middle; the base covered with silvery pubescence. Palpi testaceous. Thorax densely covered with depressed golden pubescence; the apex of the median segment transversely striated; the pleural tubercle nipple-like. Wings yellowish-hyaline; the apex from the second transverse cubital to the middle of the second recurrent nervure bright fuscous-violaceous. Legs long; the fore knees and tibiae testaceous; the tarsi are minutely spined; the hind spurs are not much more than one fourth of the length of the metatarsus.

Allied to *M. castanea* (Bingh.)

*Pompilus citherus*, sp. nov.

Black, marked with yellow; a mark on the apex of the mesonotum, two spots on the scutellum and the tegulae, yellow; wings yellowish-hyaline, the apices of both smoky; the second and third cubital cellules equal in length; legs black, marked with red and yellow; the four front tarsi annulated with yellow. ♂

Long: 11-12 mm.

Hab. Kuching, Sarawak.

Antennae reddish-brown, black above; the fifth and following joints dilated on the underside. Head, if anything, wider than the thorax, black; the face, the inner orbits broadly to near the ocelli, the clypeus, except for a broad black line in the centre above, the base of the mandibles broadly, and the outer orbits to the outer edge, yellow; the hinder part of the vertex and the occiput and cheeks thickly covered with long soft pale hair. The apex of the clypeus is broadly rounded; the labrum is two-thirds of its length and is black; the eyes are parallel; the ocelli are in a curve, the hinder are separated from each other by a slightly greater distance than they are from the eyes. The temples are narrow; the occiput transverse. Thorax thickly covered with silvery pubescence, black; a broad line on the hinder edge of the pronotum, a mark, broader than long, on the apex of the mesonotum, two oval marks behind the middle of the scutellum, tegulae and a small oblique mark over the middle coxae, yellow. The apex of the pronotum is broadly rounded; the scutellum is roundly convex, but not much raised above the level of the mesonotum. Median segment alutaceous, and thickly covered with longish pubescence. Legs black; the coxae and trochanters black, except at the apex of the anterior; the fore femora, except at the base, the middle and four posterior to near the middle, and the anterior tibiae entirely, red; the four anterior tibiae are yellow behind; the anterior tarsi yellow, black towards the apex; the middle black, with the four basal joints annulated with yellow; the hinder black; the spurs yellow. Wings yellowish-hyaline, the apices of both fuscous; the third cubital cellule is slightly shorter than the second; the first recurrent nervure is received near the base of the apical

fourth; the second shortly beyond the middle; the transverse basal nervure is interstitial; the accessory nervure in the hind wing is received shortly beyond the cubital. Abdomen densely pruinose; there are two large marks, wider than long on the base of the second segment, two marks on the base of the fourth, two larger marks on the base of the sixth and the whole of the seventh, yellow. The tibial and tarsal spines are yellow.

Allied to *P. vagabundus* Sm., which, *inter alia*, may be known from it by the second cubital cellule being twice the width of the third.

*Pompilus iliacus*, sp. nov.

Black, pruinose; the wings fuscous-violaceous; the first recurrent nervure is almost interstitial; the third cubital cellule at the top shorter than, at the bottom longer than, the second. ♀

Long: 13-14 mm.

Head wider than the thorax, the temples very narrow, the occiput transverse. Eyes large, distinctly converging above; the ocelli in a triangle; the hinder separated from each other by about the same distance they are from the eyes. Clypeus transverse at the apex in the middle; the sides broadly rounded. Prothorax large; the basal part distinctly separated all round; at the sides it projects broadly. Median segment broadly rounded from the base to the apex. Wings fuscous-violaceous, the posterior lighter in tint; the second cubital cellule at the top is distinctly longer, at the bottom distinctly shorter, than the third; the transverse basal nervure is almost interstitial, as is also the first recurrent (as in the *Salix-Mygminia* section); the second is received almost in the middle of the cellule; the accessory nervure in the hind wings is interstitial. Legs black; the tibial and tarsal spines black; the long spur of the hinder tibiae does not reach to the middle of the metatarsus. Abdomen smooth; the last segment thickly covered with long black hair.

There is no transverse furrow on the second ventral segment; the meta-thoracic spiracles are large, raised and bordered behind by a furrow; the tibial and tarsal spines are long; the underside of the tarsi are thickly spined; the tarsal claws have a stout, sharp subapical tooth.

Comes near to *P. perplexus* Sm.

*Pompilus curiniscutis* sp. nov.

Niger, facie, clypeo, orbitis oculorum, linea pronoti, scutello post-scutelloque flavis; pedibus rufo-fulvis; coxis, trochanteribus, femoribus apiceque tibiæ late nigris; alis flavo-hyalinis, apice violaceis. ♂

Long: 13-14 mm.

Hab. Borneo (Shelford).

Antennæ black; the scape yellowish beneath. Head black, sharpened, sparsely pilose; the face, clypeus, the inner orbits broad below, narrowed above and the outer more narrowly and uniformly, bright orange-yellow. The hinder ocelli are separated from each other by a slightly less distance than they are from the eyes, which converge slightly below. Mandibles orange-yellow, black at the apex. On the thorax there is a broad, interrupted line on the pronotum not extending to the tegulæ; a mark, wider than long, and with the sides at the base slightly projecting, on the hinder part of the mesonotum, a large mark on the scutellum roundly narrowed towards the base, where there is in the middle, a rounded point: at its apex and touching it, is a transverse line, which does not extend to its outer edge; and the top of the scutellum, all orange-yellow. The scutellum is broadly rounded above; the post scutellum is slightly higher than it; it is more distinctly raised and separated: its top is keeled; the sides have an oblique slope. The median segment is opaque, and thickly covered with pale pubescence; it is, except in the middle at the apex, closely irregularly reticulated. Legs rufo-fulvous, the coxæ, trochanters, the femora to near the apex and the apical third of the hinder tibiæ, black. Wings, yellowish-hyaline; the apex of both wings violaceous, the first cubital cellule at the top is fully one-fourth longer than the second; the third transverse cubital nervure in the hind wings is interstitial.

The median segment has a gradually rounded slope; the inner spur on the hinder tibiæ is not half the length of the metatarsus; the basal segment of the abdomen is narrow at the base, becoming gradually wider towards the apex; the pronotum is rather short. The transverse median nervure is received in front of the transverse basal.

Belongs to the group of *P. multipictus* Sm., and the European *P. 4 punctatus*, Fab. Characteristic is the prominent, raised, keeled post-scutellum.

*P. 4 punctatus*, I may add, is found in Japan also.

*Pompilus parmenus*, sp. nov.

Niger, vertice, fronte, linea pronoti, scutello, post-scutello, macula mesonoto, linea abdominis segmento 2', maculaque segment 7', flavis; alis flavo-hyalinis, apice fusco-fumato. ♂

Long: 12 mm.

Hab. Borneo (Shelford).

Antennae black, the scape for the greater part yellow. Head black, the face, front and the vertex, except behind, lemon-yellow; the ocellar region black; smooth, shining, almost bare. The eyes distinctly converge above, where they are separated by slightly less than the length of the third antennal joint. The apex of the clypeus is broadly rounded. Thorax black; a broad band on the centre of the pronotum behind, a large mark on the apical half of the mesonotum, its sides straight, its base irregular; and it is broader than long; the greater part of the scutellum (the mark obliquely narrowed laterally at the base), the post-scutellum and a line on the base of the second abdominal segment, lemon-yellow. The median segment is thickly covered with greyish hair; wings yellowish-hyaline; the apex is smoky, broader at the top, where the cloud extends to the second transverse cubital nervure; the second cubital cellule at the top is distinctly shorter than the first; the two transverse cubital nervures converging there; the transverse basal nervure is interstitial. Legs black; the tibiae and tarsi rufo-fulvous. The base of the second abdominal segment is lined with orange-yellow; the last segment above is broadly pale yellow.

Has the general colouration of *P. cariniscutis* here described; but is readily known from it by the flat post-scutellum and by the interstitial transverse basal nervure.

*Pseudajenia reticulata*, sp. nov.

Nigra, abdominis basi late femoribusque posticis rufis: alis fusco-violaceis, basi hyalinis ♀.



Long: 11-12 mm.

Front, face and clypeus covered with a silvery pile. Eyes distinctly converging above; at the top they are separated by twice the length of the second joint of the antennæ; the hinder ocelli are separated from the eyes by the same distance they are from each other. Clypeus rather short, broader than long. Palpi black. Thorax thickly covered with silvery pubescence; the central part of the mesonotum punctured and clearly separated from the lateral by a narrow furrow; the lateral parts are smooth, and the central part is more strongly punctured on the sides. Scutellum sparsely punctured; the post-scutellum shagreened. Median segment thickly covered with white pubescence; irregularly, closely reticulated. Meso- and metapleuræ closely, irregularly reticulated. Legs black; the hinder femora clear red; the tibiæ obscurely rufo-testaceous, the calcaria black. Wings fusco-violaceous, narrowly hyaline at the base; the third cubital cellule at the top shorter than the second, below about equal in length to it; the first recurrent nervure is received shortly behind the middle; the second at the apex of the basal third. Abdomen smooth and shining; the basal three segments ferruginous, the apical black and thickly pruinose.

*Pseudagenia borneana*, sp. nov.

Nigra, dense argenteo-pilosa; femoribus posticis rufis; alis fere hyalinis, nervis stigmatique nigris; flagello antennarum late rufo ♀.

Long: fere 12 mm.

Hab. Sarawak, Borneo (R. Shelford).

Antennæ black, the fourth and following joints rufous beneath. Head alutaceous; the lower part of the front, the face, clypeus and base of mandibles densely covered with silvery pubescence. The hinder ocelli are separated from the eyes by a slightly greater distance than they are from each other. The apex of the clypeus in the middle is smooth and shining; mandibles piceous near the middle; the apical joints of the palpi pale testaceous. Thorax densely pruinose; the pile has a fulvous tinge; the pronotum is broadly rounded; the propleuræ behind have two rounded, clearly separated, tubercles, the basal being

the larger. The median segment has a short, rounded slope; it is irregularly transversely wrinkled. The wings are hyaline, with a slight, but distinct, fulvous tinge; the first and second cubital cellules are equal in length above; the first recurrent nervure is received in the middle; the second at the apex of the basal third of the cellule; the accessory nervure in the hind wings is interstitial. Legs black; the hinder femora entirely and the four anterior tibiae and base of tarsi rufous in front. Abdomen pruinose; the basal segment is triangular, and becomes gradually wider from the base to the apex, and without a neck at the base.

This species comes near to *P. tincta*. Sm., *sec.* Cam., Manch. Mem. 1891, p. 441. That species may be known from it by its head and thorax being densely pilose, by the first cubital cellule being distinctly shorter than the second above; by the apex of the propleurae not being so distinctly bituberculate, and by the first transverse cubital nervure being roundly curved, not straight, as in the present species.

*Agenia balteata*, sp. nov.

Nigra, scapo antennarum, clypeo, mandibulis, pedibusque pallide testaceis; femoribus, tibiis tarsisque posterioribus nigromaculatis; alis hyalinis; stigmate nigro, nervis fuscis ♀.

Long: 11 mm.

Hab. Kuching, Sarawak, and Singapore.

The basal two joints of the antennae rufo-testaceous; the third joint dark testaceous; the fourth, fifth and sixth joints dark testaceous beneath. Head black; the clypeus, mandibles, and lower inner orbits yellowish testaceous; the palpi at the base testaceous, the apical joints pale yellow; the hair bundle long and dark testaceous. The front is thickly covered with a golden pile and has a narrow furrow down the middle. The eyes distinctly converge above and are separated there by the length of the third antennal joint. Thorax thickly covered with golden pubescence; the prothorax yellowish-testaceous. Legs yellowish-testaceous; the four hinder trochanters, the middle femora with an irregular line above, the apex of the hinder femora, the apex of the middle tibiae, the apical two-thirds of the hinder tibiae, the apex of the fore tarsi and four posterior, except at the

base, black. Wings clear hyaline; the stigma black, the nervures paler. Abdomen black; the apices of all the segments testaceous; the last segment almost entirely testaceous.

This is an *Agenia* as now limited. The species recorded by Smith from the Malay and Indian regions are probably mostly referable to *Pseudagenia*, Kohl. The distinction between the two consists in *Agenia* having a bundle of stiff bristles at the base of the maxilla in the ♀.

## SPHEGIDÆ.

*Ampulx striatifrons*, sp. nov.

Dark green, largely tinged with blue; the flagellum of the antennæ black; the wings hyaline; the radial cellule and the space bounded by the first and third transverse cubital nervures and the discoidal cellule smoky; the front with three stout, longitudinal keels; the space bounded by them is transversely striated ♂.

Long: 12 mm.

Hab. Kuching, Sarawak.

Antennæ black, covered with a pale pile; the scape with hardly any metallic tint. Head blue, the ocellar region largely tinged with purple; the three keels in the front are stout and all reach to the base of the mandibles; the part between them, from near their top, bears stout, oblique striæ; the part on their outer side at the top bears some large punctures; the vertex is sparsely and strongly punctured. The front ocellus is separated from the hinder by a greater distance than these are from each other, and the latter are separated from the eyes by a distinctly greater distance than they are from each other. Clypeus distinctly keeled in the middle, green, smooth and thickly covered with white pubescence. Mandibles brownish-black; their middle, on the lower side, with a row of large punctures. Prothorax elongate, the base distinctly narrowed; it is sparsely punctured and the middle of the pleuræ bears a longitudinal furrow. The central part of the mesonotum is strongly and deeply punctured; the sides are more sparsely punctured, are coppery in colour and are depressed behind. Scutellum and post-scutellum sparsely punctured. Median segment irregularly transversely striated;

the striae are more widely separated in the centre; the second keel does not reach to the apex. The apical slope is thickly covered with white hair; the striation is close and obliquely transverse; the upper lateral teeth are small. The mesopleurae are distinctly, but not closely, punctured; the upper part of the metapleuræ is stoutly striated. Legs for the greater part blue; the femora more greenish in tint than the tibiae; the inner tooth of the claw is shorter and stouter than the outer. Wings hyaline, the radial cellule, the space bounded by the first and third transverse cubital nervures and the upper part of the discoidal are smoky; the three transverse cubital nervures are distinct. Abdomen largely marked with blue and purple tints; the third segment is largely marked with rosy and brassy tints; it is strongly punctured; its apical half is distinctly depressed, is more fiery in tint than the base and is more closely and not so strongly, punctured.

Comes near to *A. sodalicia*, Kohl. from Malacca.

*Tachytes borneana*, sp. nov.

Black; the head and thorax densely covered with pale silvery pubescence; the abdominal segments banded with silvery pile; the pygidium covered with golden pubescence; wings hyaline, with a faint yellowish tinge; the nervures pale testaceous; the second and third cubital cellules at the top equal in length ♀.

Long: 13 mm.

Hab. Kuching, Sarawak.

Antennæ black, covered with a pale microscopic pile. Front, face and clypeus densely covered with silvery pubescence and more sparsely with long pale hair; the vertex sparsely with long pale hair; alutaceous; the lower part of the vertex has a narrow furrow in the middle, which ends, above the ocelli, in a smooth depression. Mandibles black; the palpi testaceous. The basal portion of the median segment has a thin furrow down the middle, which becomes conically dilated at the apex; the furrow on the apical slope is wide and deep. The second and third cubital cellules are equal in length above; they are as wide there as the space bounded by the two recurrent nervures. Legs black; the front tarsi testaceous at the apex; the calcaria

testaceous; the tibial and tarsal spines white. The abdominal segments are banded with depressed silvery pubescence; the pygidium is covered with bright golden pubescence.

The radial cellule has the apex rounded, not acute; the eyes above are separated by not quite the length of the second and third antennal joints united; and there is no appendicular cellule in the fore wings. Comes near to *T. nitidula* F. and *T. rothneyi* Cam., from both of which it may be known by the golden pile on the pygidium.

*Notogonia umbripennis*, sp. nov.

Black, covered with silvery pubescence; the pygidium with a stiff golden pile; the wings fuscous-violaceous ♀.

Length 14-15 mm.

Hab. Kuching, Sarawak.

The lower part of the front and the clypeus thickly covered with silvery pubescence; the front and vertex closely and minutely punctured; the centre of the face has an impressed line; the clypeus is smooth, shining and bare. Eyes large, distinctly converging above, where they are separated by about the length of the fourth antennal joint. The base of the mandibles is thickly covered with silvery pubescence; the palpi are black and covered with a grey pile. The mesonotum is depressed in the middle at the base; and there is a short longitudinal furrow opposite the tegulae. Median segment alutaceous; there is a narrow keel down the centre of the basal two-thirds; above the middle of the mesopleurae is a distinct striated longitudinal furrow, which does not reach to the apex. Wings fuscous-violaceous; the second and third cubital cellules above are equal in length; the recurrent nervures are received close to each other near the apex of the basal third of the cellule. Legs stout; the apex of the hinder tibiae and the metatarsus covered with a golden pile; the spines on the tibiae and tarsi are black, as are also the calcaria. Abdomen pruinose; the segments banded with silvery pile; the pygidium densely covered with bright golden pile and thickly with long stiff fulvous hair. The sides of the median segment are obscurely obliquely striated.

Comes nearest perhaps to *N. jaculatrix* Sm. from which it may be known by the dark violaceous wings.



*Notogonia tegularis*, sp. nov.

Black, densely covered with silvery pubescence; the wings hyaline, highly iridescent; the apex slightly smoky; the mesonotum and scutellum closely minutely punctured; the median segment obscurely transversely striated; the base with a longitudinal keel ♂.

Long: 9 mm.

Hab. Kuching, Sarawak.

Front, face and clypeus densely covered with silvery pubescence, this being also the case with the outer orbits and the base of the mandibles; the eyes distinctly converge above, where they are separated by almost the length of the second and third joints united. Thorax covered with a silvery pile; the mesonotum and scutellum are closely, minutely punctured, the latter more strongly than the former. The post-scutellum is depressed in the middle. Median segment alutaceous; the basal half transversely striated, but not strongly or closely and keeled down the middle. The mesopleuræ closely and distinctly punctured; on the basal half, in the centre, is a distinct longitudinal furrow; the basal half of the mesopleuræ longitudinally striated in the middle, the striae longest in the middle. Legs black, pruinose; the spines on the tibiae are pale, on the tarsi rufous. Wings hyaline, highly iridescent, somewhat infuscated at the apex; the second and third cubital cellules at the top are about equal in length and are of the length of the space rounded by the two recurrent nervures, the outer of which is received in the middle of the cellule; the appendicular is longer than usual. Abdomen shining; the segments banded with silvery pubescence; the last segment entirely covered with silvery pubescence. The first transverse cubital nervure is broadly and roundly curved and is not angled. The tegulae are black at the base, pale testaceous in front.

*Cerceris crassidens*, sp. nov.

Black, the basal four or five abdominal segments rufous; the antennal keel, a mark on the apex of the clypeus and the base of the mandibles, pale yellow; the wings smoky violaceous, paler at the base; the lower part of the mesopleuræ projecting into a stout tooth ♀.

Long : 18-19 mm.

Hab. Pankalan Ampat, Sarawak.

Antennae stout, black, the apex rufous. Head black; the outer part of the antennal keel pale yellow; it is longish, stout and has a narrower keel on its apex; the face, cheeks and clypeus are thickly covered with silvery pubescence; except behind the ocelli it is closely and somewhat strongly punctured. Mandibles black; the basal half broadly yellow; behind the middle above they project upwards into a large, smooth shining, bluntly pointed tooth. Thorax densely covered with silvery pubescence; above closely and distinctly punctured, the punctures in the middle of the mesonotum running into longitudinal striations. The scutellum is slightly depressed in the middle; the area on the median segment is longitudinally striated. Legs black, covered with a silvery pile; the four hinder tibiae are broadly lined with pale yellow behind. The wings are dark smoky, the base and the discoidal cellule paler; the hinder wings are almost hyaline except at the apex. Abdomen red, the apical two segments for the greater part black above; it is smooth, with the petiole and the penultimate segment sparsely punctured. The pygidium is closely punctured, reticulated; the sides of it are fringed with stout stiff hairs; the oblique sides of the segment are sparsely punctured; the apical half of the epipygium is incised in the middle; the incision is distinctly bordered, is rounded and obliquely narrowed behind; the segment at the base is depressed on either side.

In colouration this species is not unlike *C. viligans* Sm. and *C. sepulcralis* Sm., but may be easily separated from them by the stout projecting tooth on the mesopleuræ. The head is large and is well developed behind the eyes; the apex of the clypeus is depressed and is bluntly and shortly tuberculated in the middle and at the sides, the basal half of the petiole is keeled in the middle; the second cubital cellule on the lower side is distinctly shorter than the third.

*Cerceris latidens*, sp. nov.

Black; the inner lower orbits, the base of the mandibles, the sides of the scutellum, the post-scutellum, the apex of the petiole and an interrupted line on the apex of the third segment,

yellow; the apex of the clypeus bidentate; the area on the median segment obliquely striated; the wings hyaline, with a smoky fascia on the apex ♀.

Long: 7 mm.

Hab. Kuching, Borneo.

Antennæ black, the scape yellow, the flagellum brownish beneath. Front and vertex closely punctured, except over each antennæ; the antennal keel is stout, yellow, black above; the face is sparsely punctured, as is also the clypeus, except at the apex, which is bidentate; the teeth are broad and slightly oblique at the apex. Mandibles broadly yellow at the base. Mesonotum punctured, but not closely or deeply, as is also the median segment; the scutellum is more closely punctured. The area on the median segment is obliquely striated, except in the centre. Mesopleuræ reticulated, more strongly and distinctly below than above; the centre is deeply furrowed. Legs black; the anterior and middle tibiæ in front, the anterior tarsi, and the base of the middle tarsi, pale yellow. Wings clear hyaline, the radial cellule at the apex and the upper part of cubital below it dark smoky; the petiolated cellule is distinctly shorter than the following and receives the recurrent nervure at the apex of the basal third. The apex of the petiole, the base of the second segment and an interrupted line on the third segment are yellow; the pygidium is brownish, smooth at the base, punctured at the apex; the sides are strongly punctured; the epipygium is broadly depressed.

There is a stout, curved keel on the lower part of the metapleuræ in the middle; the lower part of the clypeus, under the projecting toothed part, is bluntly bidentate; the apex of the mandibles is bluntly rounded.

#### VESPIDÆ.

*Icaria lateballeata*, sp. nov.

Dark ferruginous, variegated with black and yellow; the petiole short, wide, narrowed distinctly at the base; rufous, its apex broadly yellow; the apex of the second segment broadly yellow, much broader in the middle than at the sides; wings hyaline, the radial cellule dark smoky, except along the lower

edge; the stigma dark, the nervures of a lighter fuscous colour (worker).

Long: 9-10 mm.

Hab. Kuching, Sarawak.

Scape of antennæ yellow, dark testaceous above, the flagellum blackish, brownish at the base and apex beneath. Head dark ferruginous, thickly covered with silvery pubescence, the vertex sparsely with fuscous hair; the lower inner orbits to near the inner part of the incision; the sides of the clypeus broadly, its apex more narrowly, a line on the outer orbits near the top, another one below and the base of the mandibles broadly above, pale yellow. The front and the vertex to the end of the ocelli are distinctly, regularly, but not very closely, punctured; the clypeus is sparsely punctured, more especially noticeable on the dark central part. The meso- and metapleuræ are black, the metanotum dark rufous; the rest of the thorax rufous, with the following parts yellow: the base of the prothorax all round and broadest on the top of the pleuræ, a large mark on either side of the base of the scutellum, a broad band, incised in the middle, on the base of the post-scutellum, two large marks on the apical slope of the median segment, and a longish mark on the mesopleuræ below the tegulæ. The pro- and mesothorax with the scutellum are closely and distinctly punctured; the median segment is almost impunctate; its central furrow is wide, with oblique sides; its upper two-thirds irregularly transversely striated. Legs dark rufous; the anterior coxæ broadly, the apex of the femora (the anterior broadly) and the base of the tibiæ broadly, yellow. Wings clear hyaline, the apex of the costal and the radial cellules, except on its lower edge, smoky; the costa and stigma blackish; the nervures pale. The petiole is not quite so long as the second segment; its basal third is narrowed; the second segment is not narrowed at the base, is bell-shaped, its length greater than its width at the apex and it is closely and distinctly punctured, more closely and rugosely at the base than at the apex; the following segments are lined with yellow at the apex.

Comes near to *I. ferruginea*, but is smaller, and darker coloured; the clypeus is broadly black in the middle, the radial cellule entirely black above, not broadly hyaline at the base; the

stigma black, not clear testaceous, and the band on the second segment is broadly dilated backwards in the middle.

*Icaria flavo-bilineata*, sp. nov.

Black, the post-scutellum and the apex of the petiole yellow; the apex of the clypeus broadly pale yellow; wings hyaline, a fuscous spot in the radial and apical cellules; the stigma yellowish.

Long: 13 mm.

(worker).

Hab. Kuching, Sarawak.

This species comes near to *I. lugubris* Sm. *Sec.* Saussure, S. E. Z. XXIII, p. 134, which is also from Borneo. The two may be separated as follows:

The cloud occupying all the radial cellule, the second transverse cubital nervure, almost interstitial, the post-scutellum and apex of petiole not yellow, the stigma black.

*lugubris* Sm.

The cloud in the radial cellule commencing at the end of the stigma, the stigma yellow; the second recurrent nervure not interstitial; the post-scutellum and apex of petiole lined with yellow.

*flavobilineata*.

Flagellum of antennæ brownish beneath. Front and vertex alutaceous, sparsely punctured, there is a narrow keel between the antennæ; clypeus covered with a sparse pale down, sparsely haired, and roundly convex; its middle at the apex not distinctly toothed; it has the narrowed apical part pale yellow. Mandibles black, the teeth dark piceous. Thorax opaque; the mesopleura and scutellum closely and distinctly, but not strongly, punctured; the mesonotum is thickly covered with a fuscous down; the base of the prothorax is sharply keeled. The scutellum has a shallow furrow down the middle. The striation on the median segment is obscure. Legs black, pruinose; the calcaria and claws white. Wings hyaline; the costal cellule is slightly smoky; the cloud in the radial cellule is at the end of the stigma and at the second transverse cubital nervure and extends to the apex; in the cubital cellule it does not extend beyond the end of the radius; the recurrent nervures are received shortly behind and beyond the middle of the cellule. Abdomen black, densely pruinose, more thickly towards the apex; the apex of the petiole is yellow.



The middle of the median segment has only a shallow indistinct furrow, not a deep one, with oblique sides as in *lugubris*; the apex of the median segment is yellow, the yellow band extending sideways over the coxæ; the petiole is short, becomes gradually wider from the base to the apex; the second segment is not much, nor abruptly narrowed at the base; in length it is, if anything, shorter than its greatest width; the clypeus at the end of the eyes is as broad as its length.

*Icaria xanthopoda*, sp. nov.

Black, largely marked with yellow; two small marks on the apex of the petiole and two large ones, extending on to the ventral surface, on the base of the second segment; the legs yellow, the posterior trochanters and the base of the femora, black; wings hyaline, the stigma and nervures brownish (worker).

Long 11 mm.

Hab. Borneo (Shelford).

Antennæ brownish, marked above with black. Head black, the clypeus, the mandibles, except their teeth, the eye incisions entirely (the yellow mark is straight and oblique on the outer-side), a large mark, narrowed below, and ending in a sharp narrowed point above, and the outer orbits, bright sulphur-yellow; behind the ocelli are two small yellow marks. The clypeus is wider than long; its sides above are roundly curved; its apex does not end in a sharp tooth. Thorax black; the edge of the prothorax all round and broadest on the pronotum, two lines on the centre of the mesonotum, two large squarish marks on the base of the scutellum, two broad ones, narrowed and rounded on the inner side, on the post scutellum, the sides of the median segment largely, and a large mark, obliquely narrowed below on the mesopleuræ, orange-yellow. Abdomen black; a mark on the sides of the post-petiole, two large marks on the base of the second segment, continued on to the ventral surface, which has the basal half yellow; a narrow line on the apex on the second segment all round and the apical segment, orange yellow.

The petiole is not quite so long as the second segment; its basal fourth is greatly and distinctly narrowed compared to the enlarged apical part; the second segment is bell-shaped; its apex about two-thirds of the total length; the base of the pro-

thorax is sharply keeled; the third cubital cellule is of the same width above as below; the third transverse cubital nervure is parallel with the second, and both are roundly curved inwardly.

*Ischnogaster flaviplagiatus*, sp. nov.

Ferruginous brown, the clypeus, eye orbits, two marks, obliquely narrowed, on the base of the scutellum, the pleuræ and the apical half of the median segment, pallid yellow; the wings clear hyaline, the stigma testaceous, the third cubital cellule not half the length of the second, the fourth at the top as long as the third. ♀

Long: 13-14 mm.

Hab. Kuching, Sarawak.

Antennæ ferruginous, the flagellum darker in the middle above. Head smooth; the front and vertex covered with silvery pubescence; the clypeus with longer fuscous hair. The lower part of the clypeus is ferruginous, the upper yellow. Mandibles yellow, their apex black. The inner orbits and the eye incisions yellow. Thorax smooth and shining, thickly covered with glistening white hair. The base of the median segment is darker coloured than the mesonotum; it is smooth and is distinctly keeled down the centre. Wings clear hyaline, highly iridescent; the stigma clear testaceous; the nervures darker; the second cubital cellule is more than twice the length of the third, which, at the top, is as long as the fourth; the second and third transverse cubital nervures are straight and converge above; the first recurrent nervure is received quite close to the first transverse cubital; the second at fully twice the distance from the second; the second recurrent nervure is slightly and roundly bent outwardly in the middle. Legs paler in tint than the body, and thickly covered with pale hair. Abdomen coloured like the thorax, the segments mottled with pallid yellow; there is a distinct pale yellow mark on the base of the second segment at the sides and a large one on the side of its ventral surface; the extreme base of the narrowed neck is also yellow. On the mesopleuræ under the tegulæ is a mark which reaches to the middle; below the middle is a large curved yellow mark.

Comes near to *I. nitidipennis* Sauss. Sec. Bingham.

*Ischnogaster nigricans*, sp. nov.

Black; a line on the pronotum, a broad one on the post-scutellum, one below the tegulae and two small ones on the apex of the median segment, yellow; the four front tibiae yellow behind; the wings clear hyaline, iridescent, the stigma and nervures black. ♀.

Long. 12 mm.

Hab. Kuching, Sarawak.

Head entirely black; the face and clypeus thickly covered with silvery pubescence; opaque, closely, but not very strongly, punctured, the clypeus less strongly punctured; the apical tooth bluntly pointed. Thorax black, covered with silvery pubescence; the scutellum with long pale hair. Mesonotum, scutellum and median segment closely and distinctly punctured; the median segment with a distinct, narrow, deep furrow down the centre, which is widened and smooth at the apex. A line on the pronotum, a mark under the tegulae, the post-scutellum broadly and two marks on the apex of the median segment, yellow. The second and third transverse cubital nervures are straight, slightly oblique and converge slightly above; the fourth cellule at the top is two-thirds of the length of the third. All the knees are yellow, the four front ones broadly; there is a black line on the yellow, near the base of the middle tibiae. Abdomen entirely black; the petiole twice the length of the thorax; the base of the second segment is widely narrowed.

*Ischnogaster ornatifrons*, sp. nov.

Black, largely marked with yellow, the antennae brownish, black above, except at the sides of the median segment with a large yellow mark dilated at the apex; the wings hyaline, the radial and two apical cubital cellules infuscated. ♀.

Long: 22 mm.

Hab. Santubong, Kuching, Sarawak.

Head black, the front, except for a small black mark in the middle above, the clypeus, the mandibles, a small oblique mark on the outer side of the antennae, one in the eye incisions and a small one on the hinder edge of the vertex, yellow. The front is distinctly, but not closely, punctured and is furrowed in the

middle, deeply and distinctly above. Thorax shining and smooth; the mesonotum more opaque and closely punctured; the scutellum is more sparsely punctured, and has a narrow keel on the basal half. The following are yellow:— a broad line on the pronotum, two large marks, rounded behind, on the base of the scutellum, a broad band, almost interrupted in the middle, on the post-scutellum, a large broad band, widened below, on the sides of the median segment, half on the metanotum, half on the pleuræ, a large oval mark below the tegulæ and a large curved mark on the lower side of the mesopleuræ. Wings hyaline, the radial and the apical two cubital cellules smoky; the second transverse cubital nervure is slightly, roundly bent outwardly, the fourth cubital cellule is half the length of the third. The front legs are yellow, lined with black in front; the apical joints of the tarsi are brownish; the middle legs are brownish-black; the base of the tibiae, their apex more broadly and the base of the tarsi more narrowly, yellow; the four hinder coxæ are broadly yellow behind. Abdomen black; a large oval mark near the middle of the second segment below; a short line on the sides of the ventral surface, a band near the base of the third segment broadest on the sides, a mark on its ventral surface, rounded on the outer side, a smaller one on the fourth and a narrow line on the fifth dorsal segment, yellow.

*Ischnogaster fulvipennis*, sp. nov.

Black, with small yellow marks; the clypeus with two marks above and one in the centre below and two small marks on the apex of the median segment; the legs and petiole dark rufous; wings fulvous; the stigma testaceous; the second transverse cubital nervure is roundly curved; the fourth cubital cellule is fully half the length of the third. ♀

Long: 23 mm.

Hab. Mt. Penrissen, Sarawak.

Antennæ black, the scape and apical joints brownish beneath, the apical joints entirely so. Head black; an irregular mark roundly narrowed below and ending in a joint, on either side of the front, a longish mark on the upper half of the face, narrowed and curved above, and there is a longish broad mark on the centre near the apex, extending to the base of the tooth and

yellow. The lower part of the vertex is sparsely and distinctly punctured; the front is more closely and not so strongly punctured, except in the middle where it is smooth; almost bare and impunctate. Mandibles black, sparsely punctured and shining. Thorax smooth and shining, except on the mesonotum which is closely and distinctly punctured; the scutellum is less strongly punctured and has a narrow keel in the middle; both are thickly covered with fulvous hair. Median segment and pleuræ smooth and shining; the pleuræ have a plumbeous hue. On the thorax the following are yellow; a line on either side of the base of the pronotum, an irregular spot on either side of the base of the scutellum, two smaller spots on the base of the post-scutellum in the centre, two small marks on the apex of the median segment, a spot about three times longer than broad on the mesopleuræ in the middle below the tegulæ, and a curved mark below the furrow, this spot having the apex narrower and more oblique than the base. Legs dark rufous, probably varying in tint; the coxæ, tibiæ and tarsi are darker coloured than the femora; the hair is long and fuscous. Wings fulvous-hyaline, darker at the apex; the stigma is testaceous, the nervures fuscous; the second transverse cubital nervure is roundly curved outwardly; the fourth cellule is fully half the length of the third. The petiole is brownish, the node black above except at the base; there is an oval, small yellow mark on either side of the second segment below, and two elongate marks on the base of the third, with a small spot on either side. There are two obscure yellow marks on the base of the median segment.

The ♂ is more richly coloured than the ♀ the yellow markings being larger and the rufous colour of the legs and petiole much brighter in tint. The front is yellow, except for a black line in the centre, the clypeus entirely yellow; the mandibles are dark testaceous; the marks on the thorax are larger, especially the upper mark on the mesopleuræ and on the base and apex of the median segment. The petiole is almost twice the length of the rest of the abdomen; the rufous colour extends to the narrowed part of the second segment, the lower half of the clypeus is keeled in the middle, the tubercle on the propleuræ is large; there is a narrow striated band on its apex, and a broader, oblique one below its middle.



One of the largest of the Oriental species.

*Ischnogaster flavolineata*, sp. nov.

Black, largely marked with yellow; two small marks on the lower part of the front, one on either side of the ocelli behind, two lines on the mesonotum, the basal half of the scutellum, the post-scutellum and the median segment, except a squarish black mark on the base, yellow; legs pale yellow, the apical half of the tibiae and of the tarsi black; wings clear hyaline; the fourth cubital cellule not much more than half the length of the third. ♀.

Long: 17 mm.

Hab. Lingga, Sarawak.

Antennae black, the apical joints of the flagellum brownish beneath. Head black; a curved mark on the vertex behind the ocelli and touching the eyes and obliquely narrowed towards the apex, the eye incisions, an ovate transverse mark over each antennae, the lower orbits broadly, the sides and apex of the clypeus, the outer orbits and the mandibles, pallid yellow. The vertex is obscurely, the face somewhat more strongly, punctured. Clypeus is smooth; its sides are covered with long silvery pubescence; the apical tooth is clearly separated, twice longer than broad, and its apex is slightly incised; the black mark has its sides at the apex prolonged, the part between them at the base being also separated. The upper edge of the pronotum is yellow, as is also the lower half of the propleurae. Mesonotum black, except for two lines on the basal half, these being dilated on the outer side at the base. The yellow mark on the scutellum is dilated laterally. Post-scutellum yellow, its apex black. Pleurae yellow, slightly streaked with fuscous; the median segment yellow, except for an irregularly squarish black mark at the base. Legs yellow; the hinder trochanters, the under side and base of the hinder femora, the basal two-thirds of the hinder tibiae and the four apical joints of the hinder tarsi, black. Wings clear hyaline, the stigma testaceous, the nervures fuscous; the fourth cubital cellule is half the length of the third; the third transverse cubital nervure is straight and slightly oblique; the second is slightly, but distinctly, roundly curved on the lower half. Abdomen black, thickly covered with longish pale

hair; there is a clear yellow band at the base of the dilated part of the petiole, a narrower one at the base of the second segment, a large oblique mark on either side of its middle, a narrow longitudinal line in its centre, and the apices of the other segments narrowly, yellow. The black on the abdomen has a brownish tinge.

## EUMENIDÆ.

*Montezumia? forticeps*, sp. nov.

Black; the clypeus, the underside of the scape, two oblique lines on vertex, a large mark, narrowed below on the outer orbits; the pronotum broadly, two lines on the mesonotum, two marks on the scutellum, two lines on the post-scutellum, the sides of the median segment broadly, a line on the side and apex of petiole, two lateral marks on the second abdominal segment and the apices of the second, third and fourth segments, yellow; the wings fuscous-hyaline, with a fulvous tinge. ♀.

Long: 21 mm.

Hab. Mt. Matang, Sarawak.

Antennæ black; the scape largely yellow below. Head largely developed behind the eyes; closely punctured, the front more closely and strongly than the vertex; the eye incision less closely punctured than the vertex; above the antennæ is a small, somewhat conical mark, which is smooth and furrowed in the middle, except above. Clypeus distinctly broader than long; sparsely but distinctly, punctured; its apex is narrowly black; the sides of the incision are oblique and project at the apex. The marks on the pronotum become roundly dilated on the outer side and do not quite reach to the middle; the two lines on the mesonotum are in the middle following the parapsidal furrows and are about equal distance from the base and apex. The two marks on the scutellum do not quite reach to the middle and are broader than long. The two marks on the median segment extend on the inner side to the edge of the furrow and are roundly narrowed on the inner side above. There is a yellow mark on the mesopleuræ, longer than broad, below the base of the front wings. Pro- mesonotum and scutellum closely punctured; the mesonotum less closely and strongly at the sides. The parapsidal furrows commence shortly beyond the middle. The

median segment is closely and strongly punctured at the base; the apical furrow is wide, becomes gradually wider towards the apex and is keeled down the middle; its apical slope is oblique. The second cubital cellule is narrowed at the top, the nervures almost touching there; both are straight and oblique; the first recurrent nervure is received distinctly behind the middle; the second close to the second transverse cubital, almost interstitial. Legs black; the forefemora are yellow at the apex; the four hinder are rufous below and probably in some examples above. Abdomen black; the apices of the basal four segments yellow; there is an oval oblique mark on either side of the second segment at the base. The petiole is nearly as long as the second segment; it is stout, with the basal third distinctly narrowed; it is distinctly, but not very closely, punctured; the second segment is alutaceous.

The generic location of this species is doubtful. It has 3- and 6-jointed palpi as in *Zethus*, and it has further the head largely developed behind the eyes as in that genus and thereby differs from *Eumenes*. The petiole is shorter and stouter than it is in the typical *Zethus* and also the second segment is not contracted at the base into a neck. The form of the cubital cellules is different from what they are in *Eumenes* and more like what they are in *Zethus*. It is not a typical *Montezumia* either, although it has certainly some affinity to that genus, which has five jointed maxillary palpi. I leave it, in the meantime, in *Montezumia*, which is, strictly speaking, an American genus.

*Zethus varipunctatus*, sp. nov.

Black; the upper side of the mandibles and a large mark on the apex of the clypeus, yellow; the scape of the antennae, the tegulae and the legs rufous; the hinder tibiae and tarsi blackish; wings fuscous-violaceous towards the apex; the apex of the clypeus broadly rounded, not dentate ♀.

Long: 17 mm.

Hab. Kuching, Sarawak.

Head thickly covered with short pale pubescence, rugosely punctured, the punctures running into reticulations on the front. Clypeus roundly convex; its greatest width greater than its greatest length; closely and distinctly punctured, but not so

coarsely as the front; its apex broadly rounded. The scape of the antennæ is broadly rufous below; the apical joints are brownish beneath. Thorax entirely black covered with a pale pile; the mesonotum and scutellum more thickly with longer pale pubescence. Mesonotum closely rugosely punctured; the punctuation sparser towards the apex; the two furrows are indistinct at the base, being confounded with the punctuation. Scutellum strongly punctured and with a narrow furrow in the middle; the post-scutellum is, if anything, more rugosely punctured; its apex is opaque, alutaceous. The median segment is opaque, alutaceous, keeled down the centre and at the sides; above it is obscurely striated. Propleuræ smooth, the upper part at the base striated. Mesopleuræ closely, rugosely punctured; the basal and apical slopes smooth. Metapleuræ strongly punctured, except for a large oblique space on the base and apex. Petiole closely and uniformly punctured; the base rufous, smooth. Legs bright rufous; the hinder tibiæ and tarsi infuscated.

This does not appear to me to be the ♂ of *4.-dentatus*, as apart from the difference in colouration, there are structural differences between them not of a sexual nature. The two may be separated thus:—

The furrow at the base of the scutellum with five stout keels; the metapleuræ coarsely punctured and striated throughout.

*4.-dentatus*.

The furrow at the base of the scutellum with eight short keels; the metapleuræ sparsely punctured above, smooth below.

*varipunctatus*.

*Odynerus cilicius*, sp. nov.

Black, largely marked with yellow; the median segment yellow, except in the middle, a dagger-shaped line on the front, a mark on either side of the ocellar region and two oblique large marks on the mesopleuræ, yellow; wings hyaline; the radial cellule and the cubital nervures in front dark smoky ♀.

Long: 12-13 mm.

Hab. Kuching, Sarawak.

Front distinctly, but not very closely, punctured, the vertex almost impunctate. The mark on the front is narrowed in the

middle and obliquely narrowed above; the eye incision, the inner orbits narrowly, and the outer orbits more broadly, yellow. Clypeus yellow, smooth, obscurely punctured at the apex; its greatest length is slightly greater than its greatest breadth; the apical incision is wide and shallow. The scape is yellow, the flagellum brownish beneath. The basal two-thirds of the pronotum broadly yellow; there are two short, narrow lines in the centre of the mesonotum; almost the basal half of the scutellum is yellow; the apex is more strongly punctured than the base, and on the sides at the apex are three oblique keels. Post-scutellum strongly and closely punctured; it has a rounded slope from the base to the apex, and is on a level with the top of the median segment, which has a rather steep straight slope, with rounded sides and a deep furrow in the middle. Pleuræ punctured, but not strongly or closely; the mesopleuræ yellow, except at the apex and extreme base; the yellow is divided in two by an oblique furrow. Wings almost hyaline, the costal cellule infuscated; the stigma yellowish, the nervures black. Legs clear yellow, the hinder femora slightly lined with black above. Abdomen yellow; a large black mark, narrowed and rounded laterally, on the apical half of the first segment; the base of the second narrowly, a large mark, narrowed laterally, and extending from near the base to near the apex, the greater part of the third, fourth and the whole of the sixth, black. The ventral surface, except the last segment, is black.

Comes close to *O. maculipennis* Smith.

*Odynerus hyades*, sp. nov.

Black, largely marked with yellow; two short lines on the mesonotum, the sides of the scutellum, the mesopleuræ largely, the sides of the petiole and two large irregularly oval marks on the second abdominal segment, yellow; legs yellow, the femora slightly lined with black; the wings hyaline, with a slight fulvous tinge; the apex smoky; the stigma and nervures black ♀.

Long: 15 mm.

Hab. Sarawak (Shelford).



Antennæ black, the scape yellowish, the flagellum brownish beneath. Head black; the clypeus, the eye incision, a large mark, narrowed in the centre above, and the outer orbits to near the top, yellow. Front and vertex rugosely punctured, the punctures running into reticulations on the sides; the space between the antennæ is yellow and smooth. Clypeus long, pyriform, its width at the base half the length; the basal part roundly convex and irregularly marked with elongate punctures; the apex transverse. Thorax black, a large mark, obliquely narrowed on the hinder part, two short narrow lines on the mesonotum, two irregular marks on the base of the scutellum on the sides, the post-scutellum broadly in the middle and the sides of the median segment, broadly, yellow. Mesonotum strongly and closely punctured; the punctures large, deeper and closer on the base than on the apex. Scutellum flat, on the same level as the mesonotum; its apex rounded; it is not quite so strongly punctured as the mesonotum, especially at the base; the post-scutellum is more coarsely punctured. The median segment is keeled down the middle; above in the middle it is stoutly irregularly transversely striated; the sides, broadly above, narrowly below, are stoutly punctured; below the middle the sides distinctly project into stout blunt teeth. The upper part of the propleuræ is irregularly, stoutly, obliquely striated; the lower part bears stout, longitudinal keels. Mesopleuræ coarsely reticulated. Metapleuræ on the upper half irregularly, closely striated. Legs clear yellow, the tarsi darker; the femora irregularly lined with black. The second cubital cellule at the top is not quite half the length of the third; the first recurrent nervure is received shortly behind the middle; the second is interstitial. All the abdominal segments have a narrow yellow line before the apex; that on the second is broader than on the others and is largely dilated backwards at the sides; on the petiole there is a large semicircular mark behind and united to it; on the base of the second laterally is a large irregularly oval mark which is incised at the base on the lower side; the petiole is coarsely, the second and third segments are finely and closely punctured; the last segment is smooth; its apex is narrowly yellow. On the second ventral segment, on the sides, is a large yellow mark, which is rounded on the inner side.

*Odynerus lybas*, sp. nov.

Black; a band, greatly widened laterally, on the pronotum, the basal two-thirds of the scutellum, a large mark on the mesopleuræ under the tegulæ, two oval marks on the base of the second abdominal segment, the apex of the first and second segments and a transverse line on the middle of the fourth, pale yellow; the legs for the greater part black; wings hyaline, the radial cellule, except along the cubitus at the base, and the apex of the apical cubital cellule dark smoky; the costa and nervures black. ♀.

Long: 10 mm.

Hab. Sarawak (Shelford).

Antennæ black; the scape yellow, the apical joints brownish beneath. Head black; the lower part of the eye incision, the basal half of the clypeus, two oblique marks near its apex, a mark immediately over the antennæ and a band, narrowed below, on the upper half of the outer orbits, yellow. Front and vertex closely and strongly punctured; the front thickly covered with pale pubescence. Clypeus punctured, but not closely; the top almost smooth; its apex is depressed slightly in the middle; the teeth are short, broad and short. Mandibles black; the lower half yellow, tinged with rufous towards the apex. Prothorax and mesonotum closely, rugosely and distinctly punctured; the mesopleuræ obscurely punctured; the upper half of the metapleuræ is punctured, but not strongly; both are thickly covered with pale pubescence. Legs black; the apical half of the four front femora in front, the four anterior tibiae except behind, and a broad band on the outer side of the hinder tibiae, yellow. Abdomen pruinose; the petiole on the dilated apical part punctured, distinctly, but not strongly, or closely; the other segments smooth; the basal spots on the second segment are large, oval and oblique.

Comes near to *O. 2-pustulatus* Sauss. There is no suture on the base of the petiole; the basal slope of the petiole is long, straight and oblique and is distinctly longer than the apical.

## ANTHOPHILA.

*Nomia robusta*, sp. nov.

Nigra, fulvo-pilosa; alis hyalinis, apice fusco-violaceo. ♀.

Long: 14 mm.

Hab. Borneo (Shelford).

One of the larger species. Head covered with deep fulvous pubescence, the vertex distinctly punctured, less closely and more deeply at the sides than in the middle; the front is rugose in the middle, with the sides punctured as in the vertex. Face roundly projecting in the middle and strongly, but not closely, punctured above. Clypeus clearly separated from the face; its middle depressed; it is strongly, but not very closely, punctured; its apex is transverse, with the sides rounded. The pubescence on the thorax is deep fulvous and dense, especially on the pleuræ; the mesonotum and scutellum are closely and somewhat strongly punctured, the base of the scutellum less strongly than the rest. The basal area on the median segment is smooth and shining, punctured round the edges; the furrow at its base is irregularly striated, especially laterally. Legs densely covered with long bright fulvous pubescence; the apices of the tarsi rufous. Wings hyaline, with a slight fulvous tinge; the apex is smoky, with a distinct violaceous tinge, the nervures, except at the apex, are dark testaceous. Abdomen shining; the apices of the segments fringed, but not very thickly, with fulvous pubescence; the ventral segments are more thickly fringed with similarly coloured hair. The tegulæ are for the greater part rufo-testaceous, the second transverse cubital nervure has a more oblique slope than the first; the recurrent nervure is received very close to it.

*Nomia borneana*, sp. nov.

Black; the basal four segments of the abdomen with smooth, shining blue bands; the legs fulvous and covered with fulvous hair; the clypeus smooth, not keeled, the face distinctly tuberculated in the middle; wings hyaline, the stigma fuscous—black; the nervures paler. ♀.

Long: 11 mm.

Hab. Borneo (Shelford).

Head black; the front, the face and sides of the clypeus thickly covered with fulvous pubescence, smooth, shining and impunctate; the tubercle on the face is more prominent than usual; the labrum is fringed with long golden hair. Mandibles

ferruginous, black at the apex. Thorax closely covered with fulvous hair; the post-scutellum region densely covered with fulvous hair; the mesonotum and scutellum are smooth and shining. The area on the median segment is shining, distinctly bordered behind and irregularly striated, the striae more widely separated in the middle than laterally; the rest of the segment is opaque and densely covered with fulvous pubescence. The transverse cubital nervures are paler than usual, this being especially the case with the second. Legs uniformly yellowish-fulvous and thickly haired; the hair is paler in colour; the calcaria pallid yellow. Abdomen black; the basal four segments with smooth, shining, bare, bluish bands; the back is smooth and shining; the basal segment at the base is thickly covered with fulvous hair; the others are sparsely haired; the ventral segments are closely punctured; their apices thickly fringed with fulvous hair. The blue belts on the abdomen are slightly tinged with yellow; the furrow on the median segment is indistinct; the scape is testaceous at the base; the second cubital cellule is about one-third of the length of the top of the third; the third transverse cubital nervure is roundly curved; the first recurrent nervure is interstitial.

Comes close to *N. elegans* Sm. which may be known from it by the clypeus being coarsely punctured, subtuberculate on each side, and with a "central longitudinal depression."

*Nomia leucozonata*, sp. nov.

Black, the basal half of the abdomen above and the ventral surface rufous, the second, third, and fourth segments banded with white on the apex; the apex of the clypeus rufous; the legs black, densely covered with long white hair; the wings hyaline, the costa and stigma rufo-testaceous, the nervures paler. ♀.

Long: 8 mm.

Hab. Bidi, Sarawak.

Head thickly covered with pale fulvous pubescence, black, the apex of the clypeus broadly rufous; the clypeus and face strongly, but not very closely, punctured; the front is more sparsely punctured and has a narrow longitudinal keel in the middle. Mandibles rufous, black at the apex. Mesonotum and scutellum minutely and closely punctured; the post-scutellar

region is thickly covered with pale fulvous pubescence. Median segment smooth, shining and bare; its sides thickly covered with long pale fulvous hair; the basal depression is not clearly bordered behind and bears narrow longitudinal keels. Legs black, thickly covered with long pale fulvous hair; the hinder tibiae become gradually thickened towards the apex; the metatarsus is thickened. Wings hyaline; the costa and stigma testaceous, the nervures paler; the second cubital cellule is narrow; the second transverse cubital nervure is faint; the first transverse cubital nervure is interstitial. Abdomen rufous; the apical three segments (the basal two broadly) marked with black in the centre; the apices of the second, third and fourth are banded with cream-white.

*Coelioxys eriocephala*, sp. nov.

Black, the head and thorax rugosely punctured; the scutellum coarsely reticulated, its sides at the apex projecting into sharp spines; the basal area of the median segment aciculated, the rest of the segment closely punctured, the upper half deeply furrowed in the middle; wings hyaline, the apex from the base of the radial cellule fuscous-violaceous. ♀.

Long: 11 mm.

Hab. Kuching, Sarawak.

Head rugosely punctured, the clypeus more closely and less strongly than the rest; the cheeks, the face and apex of the clypeus covered with pale pubescence; the outer orbits and occiput are thickly covered with white pubescence; the vertex and front are sparsely haired. Mesonotum closely rugosely punctured; the scutellum is coarsely reticulated-punctured; the apex is rounded, with the sides projecting into teeth, which are twice longer than broad; its apex projects over the post-scutellum. The basal area of the median segment is coarsely aciculated; the rest is closely punctured and thickly covered with white pubescence. Mesopleurae closely rugosely punctured and covered with white pubescence; the sides and apex are aciculated. The second transverse cubital nervure has the lower half oblique; the upper distinctly oblique and straight. Legs thickly covered with woolly hair; the basal joint of the hinder tarsi is thickly covered with fulvous pubescence. The five



basal segments of the abdomen are shining, closely, distinctly, but not strongly, punctured; their basal furrows are covered with white pubescence; the basal part of the last segment is minutely, but not very closely, punctured; the narrowed apical part is closely rugosely punctured; in its centre is a smooth narrow keel. The last ventral segment is long, narrow, acutely pointed and projects over the dorsal.

There is a distinct furrow on the front; the lateral teeth on the scutellum distinctly project beyond the middle of the apex; the ventral surface of the abdomen is more strongly and closely punctured than the dorsal; the transverse median nervure is received shortly, but distinctly, behind the transverse basal. The last dorsal segment is distinctly depressed laterally at the base of the narrowed part. The second transverse cubital nervure is received shortly behind the middle of the radius; the nervures and stigma deep black.

*Megachile alticola*, sp. nov.

Nigra, albo-pilosa; scopa abdominisque apice supra dense albo pilosis; alis hyalinis; nervis stigmatumque nigris. ♀ et ♂.  
Long: 9 mm.

Hab. Matang, Sarawak, 3,000 feet.

♀ Head black; the face and clypeus covered closely with dark fulvous pubescence; the front and vertex less closely with longer black, intermixed, on the vertex, with shorter fulvous pubescence. The front is closely rugosely punctured; the vertex is more distinctly, more strongly and less closely, punctured. The apex of the clypeus is almost transverse, with its sides rounded; its sides above bear a belt of dense white pubescence. Mandibles strongly punctured; their apex broad, slightly oblique; the lower tooth is smooth, shining and sharply pointed; the subapical is shorter and blunter. Thorax closely and distinctly punctured; the mesonotum is densely covered with short black pubescence; the pronotum behind is covered densely with white pubescence; the scutellum has posteriorly long black hair. The basal area of the median segment is smooth, bare, aciculated and almost shining; and it bears a shallow furrow in the centre; the sides of the segment are thickly covered with white hair. There is a curved band of

white pubescence round the tubercles; the sternum is covered with white pubescence. The hair on the legs is thick, stiff and black. Wings hyaline, their apex slightly infuscated. Abdomen black; the segments are probably edged with white pubescence; the apical two above are thickly covered with grey depressed pubescence; the scape is white.

The ♂ is similarly coloured; the hair on the face is longer and denser, and has a slight fulvous tinge; the hair on the median segment is longer and denser; the basal four abdominal segments have, on the sides at the apex, broad bands of white pubescence; the apical segment is round and has a broad projecting border on the lower side; the ventral segments are fringed with soft white pubescence and there is also a band of similar pubescence on their middle; the pubescence on the four hinder tarsi is fulvous.

Comes into Bingham's Section *F*, but cannot well be confounded with anything there described.

*Megachile viriplaca*, sp. nov.

Black; the head and thorax covered with fuscous-black pubescence; a tuft of long white hair below the antennæ; and the labrum is fringed with long white hair; the wings yellowish-hyaline, the nervures and stigma yellowish-fulvous. ♀.

Long: 18 mm.

Hab. Kuching, Sarawak.

Antennæ black, bare, longer than the thorax. Front and vertex rugosely, closely punctured. The hinder ocelli are separated from each other by a slightly greater distance than they are from the eyes. Between the antennæ and the top of the clypeus is a dense tuft of longish white hair; and the labrum is fringed with longer white hair. The apical tooth of the mandibles is long, curved and sharply pointed; the subapical is bluntly rounded. Mesonotum and scutellum closely punctured and covered thickly with short black hair; the median segment is thickly covered with longish sooty-black hair, as are also the pleuræ and breast. Wings yellowish-hyaline; the stigma and nervures are yellowish; the two recurrent nervures are almost interstitial; the second cubital cellule at the top is about half the length of the bottom; the second transverse cubital nervure is broadly

rounded in the lower half and middle; the first is roundly curved. The hair on the legs is short and black; the base of the fore tarsi is rounded and deeply incised; the fore coxæ are stoutly toothed. Abdomen black and covered with short black hair; the middle segments are depressed at the base; the apical segment is, at the apex, roundly incised; its middle on the apical half is depressed, below the base it is roundly raised.

*Megachile moera*, sp. nov.

Black; the four posterior femora red; the head and thorax covered with long white hair; the abdominal segments narrowly edged with pale fulvous pubescence; the ventral scope white; the base of the hinder tarsi densely covered with long fulvous hair; the wings hyaline, the stigma and nervures black.

Long: 8-9 mm.

Hab. Kuching, Sarawak.

Head closely punctured; the face and clypeus more coarsely than the vertex; the apex of the clypeus in the middle projecting, smooth and shining; the sides of the face and clypeus thickly covered with white pubescence; the front less thickly covered with long white hair, the vertex more sparsely with shorter back hair. Mandibles closely, rugosely punctured and covered with white pubescence; towards the apex they are coarsely irregularly striated; the apical two teeth are of about the same size, are opaque and moderately acutely pointed; the rest of the outer edge is smooth and shining. Pro- and mesothorax, with the scutellum, closely and uniformly punctured; the pronotum and base of the mesopleuræ above thickly covered with woolly hair; the mesonotum has a short pubescence; the median segment is covered with long, soft hair; its basal area is closely aciculated, the rest punctured, but not strongly or closely; its middle is deeply furrowed. Wings hyaline; the first recurrent nervure is received nearer the transverse cubital than is the second. The four hinder femora and the hinder trochanters and coxæ are bright rufous; the hair on the basal four joints of the hinder tarsi is larger than usual; and it becomes gradually shorter towards the fourth joint; it is bright fulvous; the hair on the middle tarsi is shorter and paler. Abdomen closely punctured; the transverse furrows on the basal three

segments are distinct; the segments are banded with pale fulvous bands of pubescence; the ventral scopa is white; the basal ventral segments are more or less rufous; the apical dorsal segment is closely rugose and sparsely haired.

The scutellum is broadly rounded behind and has a rounded slope on the apex; the depression at its base is covered with white pubescence.

Comes near to *femorata* Sm.

*Megachile aola*, sp. nov.

Black; the head and thorax covered with pale fulvous hair; the face and clypeus more densely with long pale hair; the abdomen densely with ferruginous pubescence, except on the base of the basal three segments; the wings almost hyaline, the stigma and nervures black; the apical abdominal segment entire. ♂.

Long: 12 mm.

Hab. Matang, 3600 feet. Sarawak.

Antennæ black, base shining. Head black, closely and strongly punctured; the lower part of the front, the face and clypeus are densely covered with long pale fulvous hair; the apex of the clypeus is transverse. Mandibles closely, irregularly longitudinally striated to near the apex; the lower part at the apex smooth and shining; the apical tooth is long and sharply pointed at the apex. Pro- and mesothorax closely and distinctly punctured and covered with long pale fulvous hair; the median segment is more thickly haired and the hair is longer; the basal area is closely punctured and has a wide and distinct furrow down the centre. Pleuræ thickly covered with long pale fulvous hair. Legs black; the four anterior tarsi are fringed with very long fulvous hair; the hair on the hinder tarsi is short and fulvous; on the rest of the legs it is short and paler. The second recurrent nervure is almost interstitial; the first is received close to the transverse cubital. The basal three segments of the abdomen are black, with their apices covered with bright ferruginous pubescence, the other segments are entirely covered with similar pubescence; the apical segment is entire and broadly rounded; the apical ventral segments are covered with fulvous pubescence.

*Megachile osea*, sp. nov.

Black; the face and apex of the clypeus fringed with long pale hair; the pleurae and median segment covered densely with pale fulvous hair; the abdomen, except at its basal slope, with ferruginous pubescence; wings hyaline with a slight fuscous tinge, the stigma and nervures black; the apical abdominal segment roundly incised in the middle. ♂.

Long: 14 mm.

Hab. Matang, 3600 feet. Sarawak.

Head closely punctured; the front thickly covered with long black hair, the vertex more sparsely with shorter black hair; between the antennae is a clump of long white hair and the apex and the clypeus is fringed with similar hair. The clypeus is shining and covered with short black hair; it is irregularly, somewhat strongly, but not very closely, punctured, and is clearly separated from the face, which is closely rugosely punctured—there is a smooth narrow shining line down the middle. The basal half of the mandibles is opaque and irregularly punctured; the apex is smooth and shining and with an acutely pointed apical tooth. Mesonotum and scutellum closely rugosely punctured and thickly covered with short black hair; the pleurae and median segments are thickly covered with long pale fulvous hair. The apex of the scutellum is broadly rounded. The area on the median segment is opaque and shagreened; the rest of the segment is shining and closely punctured. Wings hyaline, with fuscous-fulvous tinge, and slightly clouded at the apex; the stigma and nervures black; both the recurrent nervures are received close to the transverse cubitals. Legs black; the anterior tarsi testaceous at the base; the femora and tibiae are sparsely covered with short pale hair; the tarsi thickly with short fulvous pubescence on the under side; the anterior coxae are toothed at the apex. Abdomen, except on the basal slope, and on the basal two ventral segments, thickly covered with bright ferruginous pubescence; the apical segment is widely but not very deeply, incised in the middle; it is broadly rounded and is not keeled down the middle.



*Megachile amstela*, sp. nov.

Long: 12 mm.

Hab. Borneo.

This species may be separated from *turea*, with which it agrees closely in colouration, as follows:—

The clypeus distinctly narrowed, smooth and rounded above; the base of the mandibles broadly rounded, without a projection in the middle, the outer side not strongly, uniformly punctured. *turea* Cam.

The clypeus not distinctly narrowed above, where it is strongly punctured, and where there is a smooth, transverse keel; the outer side strongly punctured; the inner side on the basal part projecting in the middle. *amstela* Cam.

Head strongly, closely, rugosely punctured; the top and apex of the clypeus with a smooth, impunctate band; its hair black; long and dense on the front and vertex, shorter on the clypeus. The clypeus is distinctly separated from the face by a smooth, shining band, the sides being also bounded by similar bands; its apex projects slightly and roundly at the sides. Mandibles rugosely punctured, irregularly, strongly, longitudinally striated towards the apex; the apical edge and the lower on the apical half, smooth and shining; the apical tooth is large; the subapical is shorter and broader. Thorax closely punctured; the hair on the pleuræ, mesonotum and scutellum is black; on the median segment it is long and soot-coloured; on the sternum pale fulvous. Wings hyaline, with a slight, but distinct fulvous tinge; the nervures are blackish, the stigma dark rufous. Legs black; the hair pale, mixed with black; on the base of the tarsi it has a slightly more rufous tint. The abdomen is black; its basal three segments are banded with bright ferruginous pile; the other segments are thickly covered with bright ferruginous hair, mixed with black in the middle; the scopa is ferruginous.

On the outer side of the apex of the tibiæ, in the middle, is a longish, sharp-pointed spine.

*Megachile tarea*, sp. nov.

Black; the head covered with black, the pleuræ and metanotum with pale, pubescence; the abdominal segments banded with ferruginous pubescence; the scopa fulvous, rufous towards the apex; the legs covered with fulvous, the metatarsus on the inner side with rufous, pubescence; wings hyaline, the apex slightly infuscated, the costa and nervures dark fuscous. ♀.

Long: 12 mm.

Hab. Borneo.

The hair on the upper part of the head is deep black; below long and white; on the front and cheeks it is longer and denser than on the face and vertex. The clypeus is narrowed and rounded above, where it is smooth and impunctate, or only sparsely, and indistinctly punctured; the lower two-thirds punctured, but not closely or strongly; there is a narrow, impunctate band in the centre; the centre, in the apex, is not quite transverse. The mandibles are broadly, roundly dilated in the middle; there are two apical teeth; the apical is the longer; the subapical is broader and more rounded; their upper side is irregularly punctured; on the inner side is a row of punctures, on the outer, on the apical half, a curved deep furrow; the basal half on the outer side is strongly punctured, the punctures becoming larger and more elongated towards the middle. Thorax closely and distinctly punctured, the pleuræ somewhat more strongly than the mesonotum. Legs black; the hair on the femora and tibiæ is long and pale; on the tarsi it is dense and rufous; the anterior calcaria rufous; the outer joint is straight, transverse, and not dilated at the apex; the subapical is curved and surrounded by a hyaline horny process; the claws are rufous at the base.

On the apex of the fore tibiæ in the middle is a large, platelike projection, which becomes gradually narrowed towards the apex, is rufous in colour and has the sides raised; on the outer side of it is a stout tubercle.

*Megachile Shelfordi*, sp. nov.

Nigra, opaca, nigro pilosa; alis flavo-hyalinis, apice fumato; nervis stigmatique fulvis. ♀.

Long: 17 mm.

Hab. Borneo (Shelford).

Front and vertex closely and uniformly punctured, thickly covered with black hair, which is much longer on the front. The clypeus is rather strongly, closely and uniformly punctured, except above in the middle, where it is smooth and shining. Mandibles below smooth and shining, above closely punctured; there are four teeth, all bluntly rounded; the inner two project more than the apical. The upper part of the thorax is thickly covered with short, stiff hair; the hair on the pleurae is longer, is thick and sooty-black in colour. The hair on the legs is long and black; on the under side of the middle tarsi it is bright rufous. The hair on the abdomen, above and below, is deep black; the second segment at the base is deeply depressed, at the apex obliquely raised; the last segment has the apex depressed and broadly rounded. Wings yellowish-hyaline, the stigma and nervures rufo-fulvous; the apex, outside the radius, the second transverse cubital, the second recurrent and the discoidal nervures, smoky; the apex of the hind wings is likewise smoky.

Comes near apparently to *M. tuberculata* Sm.

*Protoanthidium rufobalteatum*, sp. nov.

Black; the head and thorax densely covered with stiff, moderately long black hair; the apical two segments of the abdomen entirely, the middle segments banded with rufous-yellow; the ventral fringe bright ferruginous.

Long: 12 mm. ♀.

Hab. Matang, 3600 feet. Sarawak.

Antennae black, short, smooth and shining. Head closely rugosely punctured, the clypeus more finely than the vertex; in front and above closely covered with short, black hair; below with longer, soft, pale hair. The clypeus is distinctly, narrowly keeled in the middle; the apex of the clypeus is transverse, its sides are rounded. Mandibles opaque, closely rugose above, below covered with a pale fulvous pile and below also with some long pale hair; the apical tooth is bluntly rounded and projects; behind it are three short, bluntly rounded teeth. Thorax closely and somewhat strongly and uniformly punctured;

above the hair is black; on the sides it is longer and whitish; the scutellum distinctly projects over the median segment; its apical incision is rounded and not very deep. Legs black; the hair on the tibiae and tarsi is thick and stiff; on the inner side it is rufous, the apical three joints of the tarsi are rufous. Wings hyaline with a faint, fulvous tinge; the stigma and nervures are black. Abdomen black; there is a narrow rufous line on the sides of the second segment; an almost entire one on the apex of the third, a broader one, narrowed at the sides, on the fourth and the whole of the apical two segments are rufous. The ventral scopa is bright ferruginous.

The ♂ has the antennae much longer; there is a dense mass of white pubescence over them; the clypeus, the cheeks, on either side of it, and the mandibles, except at the apex, are rufous-yellow; the apical abdominal segment is roundly, but not deeply, incised. The mandibles are bidentate at the apex; the apical is more narrowed at the apex than the subapical, which is shorter, broader and more broadly rounded; the clypeus is slightly depressed in the middle at the apex.

*Protoanthidium oratum*, sp. nov.

Black: the hair on the thorax and abdomen black; there is a tuft of fulvous hair on the front; the face and clypeus are covered with short rufous pubescence; the scopa rufous; the wings to the stigma smoky, the apex milk-white, the stigma and nervures black. ♀.

Long: 14 mm.

Hab. Matang, 3600 feet. Sarawak.

Antennae black, the under side of the scape rufous. Front and vertex closely, uniformly and strongly punctured; the face is less closely and less strongly punctured; its apex and a line in the centre, smooth. Clypeus closely and uniformly punctured; its middle keeled, but not very strongly; its apex slightly bent inwardly. The lower part and the apex of the mandibles are smooth and shining; the rest punctured and pilose; besides the apical tooth, there are three short, round ones. Mesonotum and scutellum closely and uniformly punctured; the mesonotum thickly covered with fuscous-black pubescence; the apex of the clypeus is roundly incised and projects over the median segment

which is closely, but not strongly, punctured, and is shining in the middle. Legs black, and thickly covered with black hair; the calcaria black. The wings, to the stigma, smoky-fuscous; the rest milky-hyaline; the stigma and nervures black; the first transverse cubital nervure is straight and has an abrupt oblique slope; the second is two-angled, the upper is the longer and is more oblique; both the recurrent nervures are received distinctly beyond the transverse cubitals. Abdomen black; closely but not strongly, punctured above and covered, but not densely, with short black hair; the apical segment is roundly incised; the ventral scopa is fulvous.

*Protoanthidium rufomaculatum*, sp. nov.

Black; the clypeus and the antennæ rufo-testaceous; the top of the head and the mesonotum with the scutellum thickly covered with rufo-fulvous pubescence; the scopa rufo-fulvous; the apical-dorsal segment covered with whitish pubescence; the wings dark smoky-fuscous to the base of the stigma, lacteous beyond it. ♀.

Long: 8 mm.

Hab. Kuching, Sarawak.

Antennæ shining, base rufous, the second joint black. Head closely, rugosely punctured; above thickly covered with short rufous pubescence; the face and clypeus with soft, paler hair. The apex of the clypeus on the lower side is flat, smooth and shining. Mandibles rufo-testaceous, the lower edge of the apex black; it is sparsely punctured; its apical tooth is large, is sharply pointed and clearly separated from the subapical, which is short, does not project and is not defined behind. Mesonotum closely, uniformly and somewhat strongly punctured; there is an impressed line down its centre. The scutellum is flat, is not raised above the level of the mesonotum, and its apex projects largely over the median segment; it is more closely punctured than the mesonotum; its sides and apex are broadly rufous and the apex has a shallow, rounded incision in the middle. The median segment has a vertical slope, is closely punctured and covered with a short pubescence. Mesopleuræ closely punctured like the mesonotum; the base of which is smooth, projecting. Legs black; the hair black; the greater part



of the front femora, the apex of the front tibiae and the base of the tarsi, rufous, as are also the apical joints of the hinder tarsi; the hair on the hinder tibiae and metatarsus long, black and thick; the fore tibia at the apex on the outer side is armed with a short narrow tooth. The radial, cubital and recurrent nervures are pale; the first transverse cubital nervure is straight and oblique; the second is roundly curved; both the recurrent nervures are received shortly, but distinctly, beyond the transverse cubitals; the second recurrent nervure has the upper part roundly bent outwardly. The basal five dorsal segments of the abdomen are smooth, shining; the basal ones minutely and closed punctured and almost bare; the large apical segment is thickly covered with glistening grey hair and is rounded at the apex; the scopa is rufous.

*Xylocopa Shelfordi*, sp. nov.

♀ Black, the thorax above, the upper parts of the pleurae and the basal two segments of the abdomen clothed with bright yellow pubescence; the wings hyaline, iridescent, the apex infuscated. The ♂ with the sides of the thorax and a broad band on the basal three segments of the abdomen clothed with bright yellow pubescence, otherwise coloured as in the ♀.

Long: 16 mm.

Hab. Matang, 3,600 feet. Sarawak (Shelford).

♀ Head densely covered with black hair; the black hair on the face mixed with white. Front and vertex closely and distinctly punctured; the furrow on the front is distinct; its lower half is bordered by distinct, flat, smooth keels; the clypeus is more strongly punctured than the face and has a smooth flat furrow in the centre. Mandibles smooth and shining; the basal half in the centre punctured; the apical teeth bluntly pointed and of almost equal length. Thorax above smooth and shining; the centre of the mesonotum and metanotum bare and impunctate; the apex of the latter sharply margined. The pubescence on the upper part of the mesopleurae is yellow; on the hinder edge it is paler; on the rest and on the sternum, black; legs black and covered with black hair; abdomen black; the upper surface of the basal two segments covered with yellowish hair; the other segments clothed more sparsely with shorter black

hair. Wings hyaline, with a slightly fuscous-coppery iridescence; the apex much darker coloured.

The ♂ has the upper part of the thorax covered with yellowish pubescence except in the centre, where there is a broad band of black pubescence of the same width as the lateral bands; and on the apex, where there is a thin band of pale yellow pubescence; the upper part of the pleuræ is covered with pale yellow pubescence; the tarsi are thickly covered with long black hair; the black hair on the tarsi is mixed with rufous beneath, on the tibiæ with pale, hair. The hair on the sides of the inner and outer orbits is pale.

I believe I have correctly united the sexes of this species. Both are in the Sarawak collection from Matang, where the ♀ has also been taken by Mr. Shelford at an elevation of 3000 feet. The ♂ agrees, in the arrangement of the hair bands, with that of *X. perversa* Weid. from Java, but the females are different.

*Trigona erythrogastra*, sp. nov.

Black; the basal three segments of the abdomen rufo-testaceous, the others black, suffused with rufo-testaceous, especially at the sides; the wings yellowish-hyaline to the stigma, the rest hyaline; the stigma and nervures rufo-testaceous. ♀.

Long: 7 mm.

Hab. Sarawak (R. Shelford).

Antennæ black, the flagellum brownish beneath, more broadly and distinctly on the apical than on the basal half. The occiput and the hinder part of the vertex are thickly covered with long black, stiff hair; the front is covered with a dark fuscous, thick pubescence and above sparsely with black hair. Clypeus thickly covered with fuscous down and smooth and shining. Mandibles black, smooth and shining. Thorax black, smooth and shining; the base of the mesonotum and the hinder part of the scutellum covered with long stiff black hair; the propleuræ covered sparsely with long black hair; the metapleuræ thickly with a fulvous down. The first transverse cubital nervure is faint above and almost obliterated in the middle; the second is very faint. Legs black; their hair is also black. The ventral surface is rufous.

Comes near to *T. lactefasciata* Cam. but that has the second cubital cellule narrower at the top; the thorax rufous and the femora rufous.

*Trigona flavistigma*, sp. nov.

Rufo-testaceous, the hinder tibiae and the basal joints of the four hinder tarsi black; wings hyaline, the basal half with a distinct yellowish tint; the stigma fulvous yellow, the nervures slightly darker in tint; antennae rufo-testaceous, the apical four joints black. ♂.

Long: 8 mm.

Hab. Kuching, Sarawak.

Head smooth and shining; the front and vertex sparsely covered with long black hair; the hairs on the vertex longer than those on the front. Cheeks and clypeus covered with golden pubescence; the clypeus also sparsely with black hair. Mandibles rufous, blackish at the apex. The thorax is narrower than the head and is similarly coloured; the mesonotum and scutellum are sparsely covered with blackish hair; the sides and apex of the former have a yellowish down; the pronotum is glabrous in front, sparsely haired behind; in the centre, at the base, is a wide depression. The apical slope of the scutellum is thickly covered with long pale fulvous hair. The centre of the metanotum is very smooth, shining and glabrous; the mesopleuræ thickly covered with long, pale fulvous hair. The wings have a distinct yellowish tinge to the base of the stigma; the stigma and nervures are bright rufo-fulvous. Legs coloured like the body: the hinder tibiae and the basal joints of the four hinder tarsi black; the hair on the black part of the legs is black. Abdomen coloured like the thorax; its base lighter in tint; the ventral surface is darker and is thickly covered with long blackish hair.

*Trigona lateballeata*, sp. nov.

Black; the base of the scape, the basal and the apical two segments of the abdomen rufo-testaceous; the anterior legs, the middle coxæ, trochanters, femora and the tibiae in front; the hinder coxæ and trochanters beneath, rufous, the wings clear hyaline, the stigma and nervures testaceous. Worker.

Long: 5 mm.

Hab. Kuching. Sarawak.

Head black, the apex of the clypeus pale testaceous; the front, face and clypeus covered thickly with pale pubescence; the front has a narrow furrow down the centre; the labrum is testaceous; mandibles pale rufous, blackish towards the apex; the occiput is fringed with long fusco-rufous hair. The mesonotum is bordered all round by a distinct belt of fulvous pubescence; there is a broader belt on the sides and apex of the scutellum; the post-scutellum is covered with short fulvous pubescence. Median segment closely, uniformly and distinctly punctured. Pleuræ covered with fulvous pile. Wings clear hyaline; the stigma pale, the nervures of a deeper testaceous colour; the two transverse cubital nervures are faintly indicated, the first more distinctly than the second. Antennæ black, the base of the scape broadly testaceous. Abdomen smooth and shining; rufo-testaceous; the second, third and fourth segments deep black.

*Trigona lacteifasciata*, sp. nov.

Dark luteous, the head, except the centre of the clypeus, the flagellum of the antennæ and the tibiæ and tarsi, black; wings hyaline, a milky cloud at the end of the stigma; the stigma and nervures luteous; there are two transverse cubital nervures, which are straight, oblique and approach close to each other near the top. ♀.

Long: 8.9 mm.

Hab. Borneo.

Antennæ black; the basal two-thirds of the scape rufous. Head black, the clypeus broadly, in the centre rufous; the front, face and clypeus thickly covered with a pale down; the hinder part of the occiput thickly covered with stiff blackish hair. Mandibles black. Thorax dark rufous, thickly covered on the mesonotum and scutellum with short, stiff, dark fulvous hair, which is thickest and longest on the base of the mesonotum. Median segment smooth and shining and is bare in the middle. The coxæ, trochanters and femora are coloured like the thorax; the tibiæ and tarsi black, except the apical joint of the tarsi; and they are covered with black hair. The basal two joints of the tarsi are mahogany coloured; the others are darker in tint; the ventral segments are similarly coloured and are sparsely covered with longish fuscous hair.