

Utah State University

DigitalCommons@USU

---

Bu

Bee Lab

---

1-1-1904

## Descriptions of New Species of Aculeate and Parasitic Hymenoptera from Northern India

Peter Cameron

Follow this and additional works at: [https://digitalcommons.usu.edu/bee\\_lab\\_bu](https://digitalcommons.usu.edu/bee_lab_bu)



Part of the [Entomology Commons](#)

---

### Recommended Citation

Cameron, Peter, "Descriptions of New Species of Aculeate and Parasitic Hymenoptera from Northern India" (1904). *Bu*. Paper 193.

[https://digitalcommons.usu.edu/bee\\_lab\\_bu/193](https://digitalcommons.usu.edu/bee_lab_bu/193)

This Article is brought to you for free and open access by the Bee Lab at DigitalCommons@USU. It has been accepted for inclusion in Bu by an authorized administrator of DigitalCommons@USU. For more information, please contact [digitalcommons@usu.edu](mailto:digitalcommons@usu.edu).



indebted for so many interesting

dily distinguishable from any of its  
oration, as it is far brighter in tone  
M. Goudoti, the species most similar

omus *Hindei*, sp. n.

member of the *N. pumilus* group.

f ears, tragus, skull, &c., apparently  
rs. A marked tuft of brown hairs  
ane of the ears.

ace chocolate-brown, finely flecked  
of the hairs (which attain about  
ther lighter. Under surface brown,  
perificially with whitish, especially  
of the belly; a creamy white line  
the wings with the flanks. Ears,  
interfemoral membrane dark brown.  
the body whitish brown, paling to  
rt of the wing, and darkening again

*Emini*, though with less marked pre-  
upper premolar outside the middle  
shed than in *limbatus*, more so than  
incisors deeply bifid.

(measured in skin):—

tail 35; thumb close to membrane  
pal 39, 1st phalanx 15.5; fifth

17.6; basal length 14.6; zygomatic  
nine to back of  $m^3$  6.7.

Kenya district, British East Africa.

M. no. 3.3.2.4. Original number  
1903, and presented by Mrs. Hinde.

most closely related to *N. Emini*,  
German E. Africa\*, but differs by  
re closely crushed upper premolars.

ntally printed in the original description,  
p. 41 (1901).

XXIII.—Descriptions of new Species of Aculeate and  
Parasitic Hymenoptera from Northern India. By P.  
CAMERON.

Apidæ.

*Nomia pilosella*, sp. n.

Black; the head, thorax, base of abdomen, and legs  
densely covered with longish white pubescence; the second,  
third, fourth, and fifth segments with emerald-coloured  
smooth bands; postscutellum with two stout longish black  
spines; metanotal area stoutly closely striated. Wings  
hyaline, the apex infuscated; tegulæ black, pale round the  
outer border. Face and clypeus not very strongly keeled  
down the middle. Hind femora roundly dilated above, the  
basal slope longer and not so long as the apical; the hind  
tibiæ become gradually wider towards the apex and rounded  
on the outer side; on the inner side the basal two thirds is  
straight, only slightly roundly dilated; the apex is broadly  
bluntly roundly dilated, somewhat as in *N. Westwoodi*, but  
the femora are much thicker, more dilated in the centre  
above, the basal and apical slopes are straight, oblique,  
whereas in *Westwoodi* the femora above have a gradually  
rounded curve from the base to the apex; in the present  
species they are shaped more as in *N. Elliotii* as figured by  
Smith (Trans. Ent. Soc. 1875, pl. i. fig. 7), but not by  
Bingham ('Fauna of Brit. India,' i. p. 449). The pubescence  
on the present species is much longer and denser than in  
*Elliotii*, which has the base of the metanotum "finely punc-  
tured," and there is a green band on the first abdominal  
segment; the present species has no band on it and the base  
of the metanotum is stoutly reticulated; also it is much  
more densely pilose.

*Hab.* Khasia Hills. Coll. Rothney.

*Habropoda fulvipes*, sp. n.

Black; the basal two segments of the abdomen rufo-  
fulvous, the lower inner orbits, a dagger-shaped mark (the  
"handle" above) on the centre of the face and clypeus, apex  
of clypeus, labrum, and basal half of mandibles pale yellow;  
the hair on the front of the head darker, behind paler coloured  
than it is on the thorax; antennæ dark brownish; legs

14\*

1906

fulvous, with paler hair. Wings hyaline, slightly tinged with fulvous; the nervures and stigma black. ♀.

Length 14 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Clypeus punctured; in its centre a stout keel which reaches near to the apex; the face tuberculate in the middle, its apex and that of the labrum margined; on the centre of the face at the apex is a broadly triangular yellow-fulvous mark; on either side of the top of the labrum is a brownish mark, with two projections, the inner of which is raised. There is a dark brownish plate on the outer side of the base of the hind tibiae, which is longer than broad, shield-shaped, roundly narrowed towards the apex, and with the outer edges raised. Basal four abdominal segments fringed with pale fulvous hair, the apical with longer black hair all over; the pygidium bare, with the sides broadly depressed on the apical half. Vertex smooth and shining, the front punctured, with a narrow keel down the centre.

What I suppose is the male has the head black, except the clypeus, which is pale yellowish testaceous; the lower part of the front and the sides of the clypeus thickly covered with depressed fulvous pubescence, as is also the greater part of the thorax; the abdomen above is black except the base and apices of the basal two segments broadly, and the ventral surface, which are honey-coloured; the legs are similarly coloured, except that the coxae, trochanters, femora, and hind tibiae are black above; the hind femora become gradually roundly dilated from the base to the apex; the hind tibiae curved, not swollen. Clypeus margined laterally, not very convex, its apex transverse, margined narrowly on the inner edge. Antennae entirely black. The wings are more clear hyaline than in the female, and the nervures and stigma are lighter coloured.

Compared with Bingham's figure of *H. Magrettii* the male of the present species has the thorax only very slightly haired and the hind femora are not at all so strongly dilated; its abdomen, too, is longer and narrower. It (the male) appears to be much more slenderly built than any species of *Habropoda* I have seen, is much less hairy, and has a much longer malar space, the eyes being widely distant from the base of the mandibles. The third joint of the antennae is swollen, not narrowed at the base, and is hardly so long as the fourth, whereas in what I take to be the male of *H. Radoszkowskii* it is clearly longer and distinctly narrowed at the base. In neither the female nor the male of my species is the second recurrent nervure interstitial. The Indian species

of *Habro-*  
genus.  
certainly  
it with  
females.

This sp  
agrees in  
pubescen  
clypeus t  
scutellum  
the later  
narrowed  
depressed  
plate, wh  
a triangle  
There are  
six apical  
behind th  
oblique s  
lower pai  
others; t  
are less c  
furrow o  
oblique d  
the ocella  
behind th  
bent dow  
fuscous vi

Length  
*Hab.* K

Black;  
the sides.  
minal seg  
rest of th  
wings brig  
nervures a  
Length  
*Hab.* K  
Vertex  
are much  
the raised

Wings hyaline, slightly tinged  
nerves and stigma black. ♀.

Coll. Rothney.

in its centre a stout keel which  
; the face tuberculate in the middle,  
labrum margined; on the centre of  
a broadly triangular yellow-fulvous  
the top of the labrum is a brownish  
tubercles, the inner of which is raised.  
a plate on the outer side of the base  
is longer than broad, shield-shaped,  
towards the apex, and with the outer edges  
abdominal segments fringed with pale  
with longer black hair all over; the  
sides broadly depressed on the apical  
and shining, the front punctured, with  
centre.

male has the head black, except the  
yellowish testaceous; the lower part of  
of the clypeus thickly covered with  
pubescence, as is also the greater part of  
above is black except the base and  
segments broadly, and the ventral  
grey-coloured; the legs are similarly  
coxae, trochanters, femora, and hind  
the hind femora become gradually  
at base to the apex; the hind tibiae  
clypeus margined laterally, not very  
narrowly, margined narrowly on the inner  
black. The wings are more clear  
black, and the nervures and stigma are

am's figure of *H. Magrettii* the male  
the thorax only very slightly haired  
not at all so strongly dilated; its  
narrower. It (the male) appears  
built than any species of *Habro-*  
less hairy, and has a much longer  
widely distant from the base of  
joint of the antennae is swollen,  
black, and is hardly so long as the  
I take to be the male of *H. Radosz-*  
ger and distinctly narrowed at the  
male nor the male of my species is  
are interstitial. The Indian species

of *Habropoda* can hardly be looked upon as typical of the  
genus. Possibly my male represents a distinct species. It  
certainly appears to be too slender for the female, comparing  
it with the males of other species of the genus and their  
females.

*Cælioxys cariniscutis*, sp. n.

This species is very similar to *C. khasiana*, with which it  
agrees in size, form, and coloration, including the fulvous  
pubescence on the underside of the tarsi, but it differs in the  
clypeus being keeled down the centre, in the sides of the  
scutellum being deeply furrowed, with the outer edge raised;  
the lateral teeth are stouter, depressed in the centre, become  
narrowed gradually to the apex, which is rounded and not  
depressed; the pronotum at the base projects into a large  
plate, which becomes gradually narrowed outwardly, forming  
a triangle of which the upper side is longer than the lower.  
There are eight teeth on the apical segment, two basal and  
six apical; the upper central pair are the shorter; the space  
behind them is depressed, the base of the depression with an  
oblique slope and shallower than it is at the apex; the apical  
lower pair of teeth are much longer and stouter than the  
others; the apices of the third, fourth, and fifth segments  
are less closely punctured than the rest; there is an oblique  
furrow on the sides of the second near the base and an  
oblique depression on the sides of the fifth near the base;  
the ocellar region has some smooth spaces at the sides and  
behind the ocelli; the lateral teeth of the scutellum are not  
bent downwards as in *khasiana* and *basalis*; the wings are  
fuscous violaceous to the transverse basal nervure. ♂.

Length 11-12 mm.

Hab. Khasia Hills. Coll. Rothney.

*Cælioxys khasiana*, sp. n.

Black; clypeus, face, lower part of the vertex broadly on  
the sides, pleurae, metanotum, and the apices of the abdo-  
minal segments narrowly covered with white pubescence, the  
rest of the head and thorax with short white pubescence;  
wings bright dark fuscous violaceous, highly iridescent, the  
nervures and stigma black. ♀.

Length 11-12 mm.

Hab. Khasia Hills. Coll. Rothney.

Vertex covered with large, deep, round punctures, which  
are much more widely separated at the sides of the ocelli;  
the raised central part of the front more coarsely punctured,



almost reticulated; its sides more closely and less coarsely punctured; face and clypeus closely irregularly punctured; the hair fringing the apex of the clypeus has a fulvous tinge. Mandibles to near the apex coarsely punctured, the punctures longish. Pro- and mesothorax closely covered with large deep punctures, which are small on the base of the mesonotum. Apex of scutellum broadly rounded; the lateral teeth large, almost smooth, curved down slightly at the apex. Median segment and apex of mesopleuræ smooth. Abdomen sparsely but distinctly punctured; the apex of the last segment closely rugosely punctured, the centre raised, smooth, the sides obliquely depressed; the apex becomes gradually narrowed to a point; the apical ventral segment becomes gradually narrowed and projects largely beyond the upper. Legs covered with white pubescence; the tarsi below with longish stiff fulvous hair.

Comes near to *C. basalis*, Sm.

*Nomia Rothneyi*, sp. n.

Black; an interrupted band of white pubescence on the apex of the first abdominal segment; a broad smooth white band on the apices of the second, third, and fourth segments; flagellum of antennæ brownish beneath; wings hyaline, the stigma testaceous, the nervures darker coloured. ♀.

Length 7 mm.

*Hab.* Mussoorie (*Rothney*).

Face broadly roundly raised in the middle; clypeus opaque, with clearly separated scattered punctures, its apex transverse; front and vertex closely punctured; an impressed line on the centre of the upper three fourths of the front; the sides of face, of the front, and the outer orbits thickly covered with white pubescence. Thorax closely punctured; the sides and apex of the mesonotum and the postscutellum thickly covered with grey pubescence. Basal area of metanotum large, clearly defined, strongly transversely striated; the striae distinctly separated; the narrowed basal inner edges obliquely striated. Pleuræ, sternum, and legs densely covered with long cinereous pubescence. Abdomen smooth and shining; there is a transverse, curved, impressed line behind the white band on the third and fourth segments; the apical two segments are brown.

This species cannot well be confounded with any of the described white-banded species of *Paranomia*.

sides more closely and less coarsely punctured; clypeus closely irregularly punctured; apex of the clypeus has a fulvous band near the apex coarsely punctured, Pro- and mesothorax closely covered with hairs, which are small on the base of the scutellum broadly rounded; the mesonotum smooth, curved down slightly at the front and apex of mesopleuræ smooth. Mesonotum distinctly punctured; the apex of the mesonotum closely punctured, the centre raised, the sides evenly depressed; the apex becomes a point; the apical ventral segment broad and projects largely beyond the second with white pubescence; the tarsi with fulvous hair.

*alis*, Sm.

*Rothneyi*, sp. n.

band of white pubescence on the second segment; a broad smooth white band on the second, third, and fourth segments; brownish beneath; wings hyaline, the nervures darker coloured. ♀.

*neyi*).

raised in the middle; clypeus with scattered punctures, its apex broadly punctured; an impression of the upper three fourths of the face of the front, and the outer orbits with white pubescence. Thorax closely covered with grey pubescence. Basal area of the mesonotum and the sides clearly defined, strongly transverse, distinctly separated; the narrowed mesonotum striated. Pleuræ, sternum, and mesopleuræ with cinereous pubescence. Abdomen; there is a transverse, curved, white band on the third and fourth segments are brown.

It be confounded with any of the species of *Paranomia*.

*Nomia interrupta*, sp. n.

Black; a narrow line of pale green on either side of the apex of the second abdominal segment, the tibiae rufous; the pleuræ densely covered with long fulvous pubescence, the mesonotum more sparsely with shorter black, the cheeks with pale fulvous, the face and clypeus with fulvous pubescence. Flagellum of antennæ for the greater part rufotestaceous. Wings hyaline, slightly tinged with fulvous. Face and clypeus stoutly keeled down the centre. ♀.

Length 13 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Face distinctly projecting in the middle, the projection with an oblique slope, almost smooth; the clypeus distinctly sparsely punctured; the central keel on the two is continuous. Clypeus roundly convex, its apex broadly roundly. Front and vertex sparsely indistinctly punctured. Mesonotum closely and distinctly punctured, the scutellum as closely but not so strongly. Basal area of metanotum closely, strongly, irregularly transversely striated. Tegulae rufotestaceous. Back of abdomen minutely punctured, except at the apex of the segments; the scopa bright rufous. Hair on legs long, fulvous, glistening on the tibiae and tarsi.

A distinct species, easily known by the single interrupted, smooth, greenish band on the second abdominal segment, strongly keeled face and clypeus, and the four rufous front tibiae.

*Nomia tuberculata*, sp. n.

Black; the pubescence on the head, thorax, and underside of the abdomen pale fulvous; the base of the first abdominal segment thickly covered with fulvous pubescence, the rest of the pubescence on the back short, sparse; the apices of the basal three segments broadly smooth and shining. Face roundly dilated in the centre, almost smooth; the clypeus broadly depressed in the middle, the sides roundly dilated, smooth and shining, bearing some large punctures; the top in the centre keeled, the apex is transverse, clearly separated. Wings hyaline, the apex with a fuscous cloud; stigma and nervures testaceous. ♀.

Length 13 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Face, front, and vertex sparsely punctured, the face in the centre smooth. Postscutellum thickly covered with fulvous pubescence. Median segment smooth and shining, its basal

area not defined, its sides with a few keels. Tegulae testaceous. The hair on the legs is long, dense, and fulvous, the spurs dark rufous.

Characteristic of this species is the fact that the raised centre of the face and the sides of the clypeus form three large tubercles. In Bingham's arrangement ('Fauna of Brit. India,' Hym. i. p. 459) it comes near *N. terminata*, Sm.

*Megachile khasiana*, sp. n.

The pubescence on the head, thorax, and base of abdomen dense, fulvous, on the rest of the back of abdomen and on the apex of ventral surface black; on the base the ventral scopa black; legs covered with cinereous pubescence; the pubescence on the underside of the base of four front tarsi rufous, on the hinder black; wings fuscous violaceous, the base more hyaline, paler. ♀.

Length 13 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Face and clypeus strongly but not closely punctured, the pubescence on them paler and sparser than on the front. Mandibles widely furrowed along the outer edge; the apical part bordered by a narrow curved furrow, the central with some irregular furrows, of which the apical is the wider and deeper; the apical tooth is long and stout, rounded at the apex, the second is broader and shorter and becomes gradually narrowed to the apex, which is rounded; the rest is broadly bluntly rounded and toothless. Abdomen opaque, closely punctured; the basal three segments have transverse furrows near the middle, the apex of the third is more widely depressed. Calcaria testaceous; metatarsus nearly as wide as the tibiae; apex of clypeus transverse.

Of the Indian species this comes nearest to *M. umbripennis*, Sm., recorded by Smith from Borneo and Nepal and by Bingham from Sikkim and Tenasserim. The number of mandibular teeth is not given by Bingham, but Smith states (Cat. Hym. Brit. Mus. i. p. 175) that they have four stout teeth, so his species is readily separated from *M. khasiana*.

FOSSORES.

*Trypoxylon placidum*, sp. n.

Black; the antennal scape, face, and clypeus thickly covered with silvery pubescence, the pleurae, sternum, and median segment with longish white hair, the pro- and

mesonotum with white nervures thickened, joints unite the centre; tured. Cly rounded, ra the base. the basal fr gradually w the furrow. V-shaped; coarsely tr with only t three segme Length 1 *Hab.* Kh

Black; t the scape tl shorter bla palpi pale t fulvous; t and the ba testaceous. and tarsi fi the middle the base of with a slig the latter l Face, eye-i of mesonot scutellum fulvous ha pubescence former ab half triang the median deeply fur petiole dila Comes n a small tu present sp



P. Cameron on new

sides with a few keels. Tegulae testaceous, the legs is long, dense, and fulvous, the

This species is the fact that the raised sides of the clypeus form three Bingham's arrangement ('Fauna of B. 459) it comes near *N. terminata*, Sm.

*achile khasiana*, sp. n.

the head, thorax, and base of abdomen rest of the back of abdomen and on face black; on the base the ventral covered with cinereous pubescence; the underside of the base of four front tarsi black; wings fuscous violaceous, the

er. ♀.

Coll. Rothney. strongly but not closely punctured, the paler and sparser than on the front. furrowed along the outer edge; the apical row curved furrow, the central with of which the apical is the wider and th is long and stout, rounded at the broader and shorter and becomes the apex, which is rounded; the rest ded and toothless. Abdomen opaque, basal three segments have transverse e, the apex of the third is more widely staccous; metatarsus nearly as wide typeus transverse.

this comes nearest to *M. umbripennis*, from Borneo and Nepaul and by and Tenasserim. The number of given by Bingham, but Smith states . p. 175) that they have four stout adily separated from *M. khasiana*.

FOSSORES.

*lon placidum*, sp. n.

scape, face, and clypeus thickly escence, the pleurae, sternum, and ngish white hair, the pro- and

*Hymenoptera from Northern India.*

217

mesonotum thickly with long fuscous hair, the legs sparsely with white pubescence. Wings hyaline, the apex smoky, nervures and stigma black. Apical joint of antennae thickened, nearly twice the length of the preceding two joints united. Front alutaceous, indistinctly furrowed in the centre; vertex opaque, finely, not very distinctly punctured. Clypeus not carinate in the middle, its apex broadly rounded, raised, smooth. Palpi pallid testaceous, black at the base. The apex of metanotum has an oblique slope; the basal furrow extends from the base to the apex, becomes gradually wider, is shallow and finely transversely striated; the furrow on apical slope wide, deep on the basal half, V-shaped; the apical third of the segment is somewhat coarsely transversely striated. Abdominal petiole narrow, with only the apex dilated; it is as long as the succeeding three segments united, its apex distinctly clavate. ♀ ♂.

Length 13 mm.

*Hab.* Khasia Hills. Coll. Rothney.

*Trypoxylon fulvocollare*, sp. n.

Black; the basal five or six joints of the antennae fulvous, the scape thickly covered with white hair, the flagellum with shorter blackish pubescence; clypeus and mandibles rufous, palpi pale testaceous; the apex of pronotum and tubercles fulvous; the base and sides of the first abdominal segment and the base of the second and third segments broadly rufotestaceous. Apex of fore coxae, trochanters, femora, tibiae, and tarsi fulvous, the femora of a deeper hue, the apex of the middle femora, the middle tibiae and base of tarsi, and the base of the hinder tibiae pale testaceous. Wings hyaline, with a slight fulvous tinge, the costa and stigma fulvous, the latter lighter in tint; the radius and cubitus testaceous. Face, eye-incision, outer orbits and the base, sides and apex of mesonotum thickly covered with golden pubescence; the scutellum with short fuscous, the postscutellum with longer fulvous hair; the pleurae and sternum with short pale fulvous pubescence. Front and vertex sparsely punctured, the former above with a wide and shallow furrow; the lower half triangularly keeled. On the apex of the basal half of the median segment is an elongated fovea; the apical half deeply furrowed in the middle. The apical third of the petiole dilated.

Comes nearest to *T. coloratum*, Sm.; that species has only a small tubercle above the base of the antennae, while in the present species there is not a tubercle, but a long stout



keel. There is no lateral furrow on the base of the median segment. ♀.

Length 17-18 mm.

*Hab.* Khasia Hills. Coll. Rothney.

*Trypoxylon khasia*, sp. n.

One of the larger species. In size comes near to *T. coloratum*, which differs in having the pubescence golden. In Bingham's table ('Fauna of Brit. India,' Hym. i. p. 224) it comes into B, except as regards the size and  $\delta^3$ . "Abdomen red, basal segment only black."

Black; apex of clypeus testaceous; mandibles yellow, their teeth black; palpi yellow; scape and base of flagellum of antennæ pale yellow, the rest black, brownish beneath; the base and sides of mesonotum with a distinct fulvous-yellow band; tubercles yellow, except at the base, and fringed with silvery hair; abdomen rufo-testaceous, the petiole black, except at the apex, the black there being triangularly incised in the middle. Four front legs yellowish, the femora of a more testaceous hue, the base of all the coxæ black; the hind femora black, running into testaceous towards the middle; the hinder tibiæ yellowish beneath, flavo-testaceous above, blackish towards the apex, the tarsi blackish, the apices of all the joints testaceous. Wings hyaline, the stigma testaceous. The clypeus, orbits, and eye-incision densely covered with silvery pubescence; the front obscurely punctured; a narrow furrow runs from the ocelli. Thorax densely pilose, the pile fuscous on the mesonotum, longer and more silvery on the sides; the sides and apex of median segment thickly covered with pale hair; at its base in the middle is an elongated somewhat pear-shaped depression; the apex is deeply and widely furrowed and densely covered with long white hair. The greater part of the pleuræ covered with silvery pubescence.

Length 20 mm.

*Hab.* Khasia Hills. Coll. Rothney.

*Trypoxylon orientale*, sp. n.

Antennæ black, distinctly thickened towards the apex, the scape thickly covered with long white hair; flagellum bare. Face, lower part of eye-incision, and clypeus thickly covered with silvery pubescence; front and vertex opaque, covered with long fuscous hair. Palpi pale testaceous. Thorax thickly covered with pale hair. Median segment short, its apex with an oblique slope; on the base is a striated depres-

sion, which is moderately deep. Furrow on the femora longer than the apex of the segment rufo-testaceous. Following the first transverse keel.

Length 22 mm.

*Hab.* Khasia Hills.

Black; the whole of the antennæ with white hair. The first cubital segment, the first transverse keel, the second transverse keel.

Length 5 mm.

*Hab.* Khasia Hills.

Antennæ black, beneath sparse. Clypeus thickly covered with white hair, and vertex also parallel to the furrow; antennæ sides distinct. The first transverse keel is a larger broad rounded. Petiole with white hair, punctured; on the base of the petiole with stout silvery pubescence becoming slightly a few stout silvery leaf-like expansions covered with white hair. Projection. Prothorax below the median oblique furrow is a smooth depression, the apex of the first transverse keels.

Comes from therefrom.

furrow on the base of the median

Coll. Rothney.

*Mylon khasia*, sp. n.

es. In size comes near to *T. colo-*  
aving the pubescence golden. In  
of Brit. India, Hym. i. p. 224) it  
regards the size and  $\delta^3$ . "Abdomen  
black."

testaceous; mandibles yellow, their  
; scape and base of flagellum of  
rest black, brownish beneath; the  
tum with a distinct fulvous-yellow  
except at the base, and fringed with  
rufo-testaceous, the petiole black,  
back there being triangularly incised  
nt legs yellowish, the femora of a  
e base of all the coxae black; the  
ning into testaceous towards the  
yellowish beneath, flavo-testaceous  
the apex, the tarsi blackish, the  
aceous. Wings hyaline, the stigma  
orbits, and eye-incision densely  
scent; the front obscurely punc-  
runs from the ocelli. Thorax  
fuscous on the mesonotum, longer  
sides; the sides and apex of median  
with pale hair; at its base in the  
somewhat pear-shaped depression;  
ely furrowed and densely covered  
e greater part of the pleurae covered

Coll. Rothney.

*on orientale*, sp. n.

ly thickened towards the apex, the  
long white hair; flagellum bare.  
ision, and clypeus thickly covered  
front and vertex opaque, covered  
Palpi pale testaceous. Thorax  
hair. Median segment short, its  
; on the base is a striated depres-

sion, which becomes gradually wider towards the apex and is moderately deep; there is a smooth, narrower, oblique furrow on the sides. Legs black, pilose, the hair on the femora longer. Wings clear hyaline, the sides of the dilated apex of the petiole and the base and sides of the second segment rufous; the petiole long and slender, as long as the following three segments united. ♀.

Length 22 mm.

Hab. Khasia Hills. Coll. Rothney.

*Psen rufo-balteata*, sp. n.

Black; the apex of the second abdominal segment and the whole of the third rufous; the fifth and following joints of the antennae testaceous beneath. Legs thickly covered with white hair; the spurs pale rufous. Wings clear hyaline; the first cubital cellule in front is half the length of the second, the first recurrent nervure is received very near the first transverse cubital, the second at twice the distance from the second transverse cubital. ♀.

Length 5 mm.

Hab. Khasia Hills. Coll. Rothney.

Antennae stout, thickened towards the apex; the scape beneath sparsely covered with long black hair. Face and clypeus thickly covered with silvery pubescence; the front and vertex almost bare, sparsely punctured; the eyes almost parallel; the ocelli in pits; the front with a shallow central furrow; antennal tubercle large, the apex triangular, its sides distinctly margined, the middle depressed; below this is a larger broader one, roundly incised at the apex, the sides rounded. Pro- and mesothorax shining, sparsely covered with white hair; the mesonotum distinctly but not closely punctured; the scutellums with a few fine punctures. Area on the base of metanotum narrow, elongate, and marked with stout striæ; the central furrow is wide and deep, becoming slightly wider towards the apex, and marked with a few stout striæ; on either side of it at the apex is a large leaf-like expansion, its apex transverse, on the outer side covered with long hair; at its base is a small rounded projection. Pro- and mesopleurae finely and sparsely punctured; below the mesopleural tubercles is a wide, deep, slightly oblique furrow, marked with some transverse keels; there is a smooth furrow near the base of the metapleurae; the apex of the latter is rugose and is marked with some transverse keels.

Comes near to *P. rufiventris*, but is quite distinct therefrom.

## Ichneumonidæ.

*Suvalla annulipes*, sp. n.

Agrees with *S. levifrons* in having the front and vertex smooth, but is smaller, has two yellow marks on the mesonotum, the scutellum yellow from base to apex, not yellow on the basal half only; the mark on the mesopleuræ is smaller, and there is none on the metapleuræ; the black on hind femora broader, reaching to the middle; the hind coxæ are yellow at the base, not broadly in the middle, and the hind trochanters yellow, not black.

The ninth to twenty-second joints of antennæ yellowish white below; the flagellum thickly covered with black short hair; scape shining, sparsely pilose. Face, clypeus (except at apex), labrum, mandibles, palpi, apical two thirds of scutellum, a large somewhat triangular mark on the sides of the first abdominal segment at the apex, the other segments broadly at the sides, and the apical almost entirely, yellow. Four front legs fulvous, yellow at the base, the end joint of tarsi black; hind coxæ black, with a large yellow band on the base above; the trochanters and basal half of femora fulvous, the apical half of femora and of the tibiæ black, the basal half of the tibiæ and of the tarsi yellowish. ♀.

Length 12 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Face strongly punctured above, in the middle transversely striated; apex of clypeus smooth, the base punctured; mandibles punctured at the base. Front and vertex smooth and shining, sparsely covered with long black hair; the space between the hind ocelli with large deep punctures. Mesonotum rugosely punctured, reticulated in parts. Scutellum covered with long fuscous hair, its yellow mark is rounded before and behind, its sides coarsely punctured, the depressions stoutly striated behind. In the centre of the metanotum at the base is an area wider than long and having inside a few stout oblique keels; the rest of the basal region reticulated, the reticulations wider on the inner side; the rest is strongly closely reticulated; the teeth large, broad, rounded at the top, looked at from behind. Propleuræ sharply margined at the base above, above strongly punctured; the rest stoutly striated. Mesopleuræ stoutly longitudinally striated, except in the middle behind; immediately under the tubercles the striae are vertical or oblique; near the base under the tubercles is a keel. Metapleuræ coarsely rugosely punctured, the punctures running into reticulations.

Areolet also  
the apical th  
punctured;  
punctures;  
tured, the  
pressed in t

Agrees cl  
known from  
not so much  
the head is  
pronotum is  
distinctly tu

Length to

*Hab.* Kha

Black; cl  
near the en  
the top, edg  
behind), scu  
the sides of  
a large mar  
narrowed to  
the lower si  
rowed above  
the hind wi  
and the si  
Four front  
yellow; the  
legs are of  
slightly mo  
fourth of ti  
tarsi are of  
hyaline, the  
areolet is  
broad, the

Mesonotu  
into striatic  
fuscous hair  
The depress  
at its sides  
punctured.  
punctured;  
the sides;  
side of the



meumonidæ.

*annulipes*, sp. n.

ms in having the front and vertex as two yellow marks on the meso- low from base to apex, not yellow the mark on the mesopleuræ is on the metapleuræ; the black on hing to the middle; the hind coxæ t broadly in the middle, and the ot black.

cond joints of antennæ yellowish n thickly covered with black short ely pilose. Face, clypeus (except les, palpi, apical two thirds of at triangular mark on the sides of at at the apex, the other segments he apical almost entirely, yellow. ellow at the base, the end joint of ck, with a large yellow band on anters and basal half of femora emora and of the tibiæ black, the of the tarsi yellowish. ♀.

H. Rothney.

above, in the middle transversely s smooth, the base punctured; e base. Front and vertex smooth er with long black hair; the celli with large deep punctures. red, reticulated in parts. Scu- fuscous hair, its yellow mark is its sides coarsely punctured, the behind. In the centre of the area wider than long and having eels; the rest of the basal region s wider on the inner side; the ulated; the teeth large, broad, l at from behind. Propleuræ se above, above strongly punct- ed. Mesopleuræ stoutly longi- the middle behind; immediately e are vertical or oblique; near is a keel. Metapleuræ coarsely tures running into reticulations.

Areolet almost square, the recurrent nervure received near the apical third. Abdomen short; petiole shining, strongly punctured; between the apex and the middle are scattered punctures; the second and third segments are closely punctured, the others smooth; gastrocoeli smooth, hardly depressed in the middle.

*Suvalta pallidinerva*, sp. n.

Agrees closely in coloration with *S. annulipes*, but may be known from it by the longer, more slender petiole, which is not so much dilated at the apex nor so strongly punctured; the head is wider compared with the mesothorax, and the pronotum is much more dilated at the base, it being there distinctly tuberculate.

Length to apex of petiole 7 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Black; clypeus (except at apex), palpi, the inner orbits to near the end of the vertex, the outer more broadly from near the top, edge of pronotum (broadly in front, more narrowly behind), scutellum broadly in the middle, a broad line on the sides of median segment from shortly above the spines, a large mark on the base of propleuræ (broad above, gradually narrowed towards the apex), the tubercles, a large mark on the lower side of the base of mesopleuræ (curved and narrowed above), the base and lower side straight, a mark under the hind wings, the greater part of the lateral scutellar keels, and the sides of the abdominal segments broadly, yellow. Four front legs fulvous, their coxæ and trochanters pallid yellow; the fore femora lined with black above; the hinder legs are of a deeper fulvous tint; the coxæ, trochanters, slightly more than the apical third of the femora, apical fourth of tibiæ, the spurs, and base of metatarsus black; the tarsi are of a more yellowish tint than the tibiæ. Wings hyaline, the nervures, stigma, and costa pale testaceous; the areolet is of equal width throughout, a little longer than broad, the recurrent nervure is received in its apical third.

Mesonotum rugosely punctured, the punctures running into striations in the middle. Scutellum covered with long fuscous hair, smooth; postscutellum very smooth and shining. The depression in the middle of median segment is smooth, at its sides it is finely punctured, the outer part coarsely punctured. The segment outside the keel is coarsely rugosely punctured; the teeth broadly rounded at the top, oblique at the sides; behind them are some curved keels. The upper-side of the pronotum is roundly incised in the centre, the



sides of the propleuræ above are strongly punctured, the middle strongly obliquely striated. Mesopleuræ longitudinally striated (except on the lower side at the base, which is punctured, and a small smooth space in the middle behind); the tubercles are punctured. Metapleuræ closely rugosely punctured, above more closely punctured. Mesosternum punctured, its furrow triangularly widened at the apex. Petiole shining; before its apex is a punctured band, surrounding a smooth space; the second and third segments are closely punctured; gastrocœli shallow, coarsely aciculated, the part behind them is raised on the outer side and is very smooth and shining.

The antennæ and the apical abdominal segments are broken off.

*Algathia rufopetiolata*, sp. n.

Black; the scape of antennæ rufous below, the base of flagellum brownish, joints 11-14 white below; a triangular yellowish mark in the centre of the face above; apex of mandibles testaceous; palpi pale yellow; scutellums, the sides of the apex of median segment, the apices of the second, third, fifth, and the whole of the apical abdominal segments pale yellow; the petiole rufous, its apex yellow. Legs red; the four front coxæ and trochanters pale yellow; the femora and tibiæ fulvous, the tibiæ paler, the fore tarsi fuscous except at the base, the middle blackish; the hinder coxæ, trochanters, and femora rufous; a large mark on the coxæ below at the apex, the apex of femora, and tibiæ (except a small dull rufous band at the base) black; the calcaria pale; tarsi black, with rufous spines. Wings hyaline, with a slight fuscous tinge, the nervures and stigma black; areolet narrowed in front; transverse median nervure received shortly in front of the transverse basal. ♂.

Length 8-9 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Face sparsely punctured laterally, more thickly in the centre; clypeus punctured, more sparsely below; labrum fringed with long fulvous hair; the face thickly covered with fuscous pubescence. Front sparsely punctured, the inner orbits above sharply margined. Mesonotum closely punctured, the apex in the middle broadly longitudinally striated, almost reticulated; middle lobe raised at the base. Scutellum roundly raised, smooth, covered with long pale hair. Postscutellum bifoveate at the base. Metanotum at the base with large deep punctures; the areola longer than

broad, round the rounded distinct one striated; the propleuræ a lower with closely and rugosely above the middle, base; the base of the striated, the shallow, broad

Black; a base of the lateral area first abdominal middle black the sides of seventh entire antennæ rufous above trochanters rufous, the and the tarsal tinge, the st

Length 8-

*Hab.* Khasia Hills.

Face closely punctured, the clypeus front and vertex indistinctly longitudinally striated, shining, the sides obliquely marked towards the base and a less distinct the sides, the posterior middle striated; the at the base stoutly on

above are strongly punctured, the  
ely striated. Mesopleuræ longitu-  
on the lower side at the base, which  
small smooth space in the middle  
are punctured. Metapleuræ closely  
ve more closely punctured. Meso-  
furrow triangularly widened at the  
before its apex is a punctured band,  
ace; the second and third segments  
gastrocoeli shallow, coarsely acicu-  
them is raised on the outer side and  
ing.

the apical abdominal segments are

*rufopetiolata*, sp. n.

antennæ rufous below, the base of  
ts 11-14 white below; a triangular  
centre of the face above; apex of  
palpi pale yellow; scutellums, the  
an segment, the apices of the second,  
e of the apical abdominal segments  
rufous, its apex yellow. Legs red;  
trochanters pale yellow; the femora  
tibiæ paler, the fore tarsi fuscous  
middle blackish; the hinder coxæ,  
rufous; a large mark on the coxæ  
apex of femora, and tibiæ (except a  
the base) black; the calcaria pale;  
pines. Wings hyaline, with a slight  
res and stigma black; areolet nar-  
se median nervure received shortly  
basal. ♂.

Coll. Rothney.

ed laterally, more thickly in the  
ed, more sparsely below; labrum  
us hair; the face thickly covered  
t. Front sparsely punctured, the  
ly margined. Mesonotum closely  
the middle broadly longitudinally  
d; middle lobe raised at the base.  
smooth, covered with long pale  
cate at the base. Metanotum at  
punctures; the areola longer than

broad, rounded at the base, gradually narrowed to the apex;  
the rounded basal part has a stout central keel and a less  
distinct one at the apex; the lateral area stoutly obliquely  
striated; the posterior median strongly, closely, transversely  
striated, the sides more sparsely and strongly. Base of  
propleuræ aciculated, the upper half strongly punctured, the  
lower with some stout striations; meso- and metapleuræ  
closely and strongly punctured, the latter more strongly and  
rugosely above the keel. The postpetiole has a depression in  
the middle, which is wide at the apex, narrowed towards the  
base; the second and third segments closely punctured, the  
base of the second closely and strongly longitudinally  
striated, the striæ going on to the gastrocoeli, which are  
shallow, brownish, and aciculated at the apex.

*Algathia tibialis*, sp. n.

Black; a mark in the middle of the face above and the  
base of the mandibles rufous; palpi lemon-yellow, the apical  
lateral area of the mesonotum except at the base yellow;  
first abdominal segment blood-coloured, the sides in the  
middle blackish, the apex yellow, the apex of the second,  
the sides of the third broadly at the apex, and the sixth and  
seventh entirely, pale yellow; the basal three joints of the  
antennæ rufous beneath, the tenth to sixteenth white,  
fuscous above; four front legs rufo-fulvous, the coxæ and  
trochanters yellow; the middle tarsi fuscous; the hind legs  
rufous, the apex of the femora, tibiæ (except at the base),  
and the tarsi black. Wings hyaline, with a slight fulvous  
tinge, the stigma fuscous. ♀.

Length 8-9 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Face closely punctured, thickly covered with fuscous  
hair, the clypeus more strongly and sparsely punctured, the  
front and vertex shining, sparsely punctured, the former  
indistinctly keeled. Mesonotum strongly punctured, longi-  
tudinally striated in the middle towards the apex. Scutellum  
shining, thickly covered with long fuscous hair. Arcola  
obliquely narrowed towards the base and to a less extent  
towards the apex, which is transverse; its base has a central  
and a less distinct and more irregular longitudinal keel on  
the sides, the middle irregularly transversely striated; the  
posterior median area is (except at the base) transversely  
striated; the outer basal area coarsely punctured, smooth  
at the base; the apical stoutly transversely striated, more  
stoutly on the apical than on the basal half; spiracular

area finely rugose at the base, more coarsely transversely rugose before the spiracles, the middle with some stout curved keels, the apex much more closely but not quite so sharply obliquely striated. Mesopleuræ closely punctured, strongly irregularly striated under the tubercles; the metapleuræ uniformly, somewhat strongly punctured and deeply depressed at the base above. Postpetiole obscurely punctured laterally and furrowed in the middle. Scutellum thickly covered with long fuscous hair; postpetiole striated at the base.

Agrees closely in coloration with *A. rufopetiolata*, including the rufous petiole; may be known from it by the narrower areola, which receives the keel in the middle, while in *rufopetiolata* it is received clearly above the middle.

*Algathia latiballeata*, sp. n.

Agrees closely in coloration with *A. zonata*; may be known from it by the black hind coxæ, by the base of metanotum being rugosely punctured, by the areola being rounded at the base and stoutly transversely striated, &c.

Black; a small mark on the face below the antennæ (broad at base, gradually narrowed to the apex, as long as it is wide at the base), palpi, tegulæ, scutellums (except the scutellum at the base), the apex of the first abdominal segment, of the second and third more broadly, the apical two thirds of the penultimate, and the whole of the last, pale yellow. Mandibles black, the apical third rufous. Palpi pale yellow. Four front legs fulvous, the tarsi infuscated, the coxæ and trochanters pale yellow; the hind coxæ black, the top and more or less of the inner side rufous; basal joint of trochanters rufous, apical yellowish; the femora with the apex broadly black, the black more extended above; tibiæ and tarsi black, the former broadly rufous at the base; calcaria pale. Wings hyaline, with a fulvous tinge; nervures fuscous, darker at the base; areolet much narrowed in front; the second transverse cubital nervure faint.

Length 11 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Scape of antennæ testaceous in the middle below; the flagellum at the base obscure brownish, thickly covered with black hair. Face and clypeus with widely separated punctures and covered with long fuscous hair. Front and vertex shining, sparsely haired and punctured; the front ocellus surrounded by a furrow. Mesonotum shining, punctured in the middle, at the base of the basal lobe almost striated.

The  
with  
than  
apex  
furrow  
versel  
a stout  
irregu  
apex;  
and d  
base, t  
with l  
obliqu  
tured,  
the ke  
domin  
and pu  
second  
second  
closely  
thickly

Black  
below;  
the top  
(except  
behind  
metano  
an irre  
part of  
small n  
scutellu  
abdomi  
segment  
trochan  
black;  
black.  
blackish  
touchin  
Leng  
*Hab.*  
Face  
cence;  
centre a  
Ann.



base, more coarsely transversely striated, the middle with some stout punctures much more closely but not quite striated. Mesopleuræ closely punctate, transversely striated under the tubercles; the propleuræ somewhat strongly punctured and the base above. Postpetiole obscurely furrowed in the middle. Scutellum with a fuscous hair; postpetiole striated

Comparison with *A. rufopetiolata*, included may be known from it by the postpetiole receiving the keel in the middle, the keel received clearly above the middle.

*latibalteata*, sp. n.

Comparison with *A. zonata*; may be known from it by the black hind coxæ, by the base of metanotum punctured, by the areola being rounded and transversely striated, &c.

On the face below the antennæ narrowed to the apex, as long as it is, tegulæ, scutellums (except the apex of the first abdominal segment and third more broadly, the apical segment, and the whole of the last, pale except the apical third rufous. Palpi black, legs fulvous, the tarsi infuscated, the hind coxæ black, the inner side rufous; basal joint yellowish; the femora with the black more extended above; tibiae black, broadly rufous at the base; areolet with a fulvous tinge; nervures black; areolet much narrowed in the cubital nervure faint.

Rothney.

Black in the middle below; the areola brownish, thickly covered with black punctures with widely separated punctures; fuscous hair. Front and vertex black, punctured; the front ocellus shining, punctured in the basal lobe almost striated.

The basal lateral area of metanotum strongly punctured and with a curved furrow on the inner side; areola twice longer than wide, bulging out obliquely in the middle and with the apex wider than the base and transverse; its basal half is furrowed deeply down the middle, the sides irregularly transversely striated; the base of the posterior median area with a stout longitudinal keel in the middle, the rest transversely irregularly striated, the striae weaker and closer towards the apex; the outer area strongly punctured, the punctures large and deep, the spiracular area irregularly reticulated at the base, the rest strongly transversely striated. Propleuræ above with large deep punctures, the depressed middle stoutly, obliquely, irregularly striated. Mesopleuræ strongly punctured, rugosely near the tubercles; the metapleuræ between the keels strongly and uniformly punctured. The first abdominal segment smooth, the sides of postpetiole depressed and punctured; the middle with a large deep depression; the second and third segments closely punctured, the base of the second closely longitudinally striated; the gastrocœli shallow, closely striated (except at the apex); the apical segments thickly covered with white hair.

*Algathia rufipes*, sp. n.

Black; the tenth to fifteenth joints of antennæ white below; face, clypeus, inner orbits and the outer from near the top, mandibles at the base, palpi, edge of pronotum (except at the base) narrowly, an irregular squarish mark behind the middle of the mesonotum, scutellums, sides of metanotum broadly, the lower edge of propleuræ, tubercles, an irregular mark (narrowed in the middle) on the lower part of mesopleuræ, a short line below the hind wings, a small mark over the hind coxæ, the centre and apex of scutellum, the apices of the first, second, third, and fourth abdominal segments, of the sixth, and the whole of the apical segment, yellow. Legs rufous, the four front coxæ and trochanters yellow; the hind coxæ and base of trochanters black; the hind tarsi white, the basal and apical joints black. Wings hyaline, the stigma testaceous, the nervures blackish; areolet narrowed above, the nervures almost touching there; transverse median postfurcal. ♀.

Length 10, terebra 1 mm.

Hab. Khasia Hills. Coll. Rothney.

Face closely punctured and covered with silvery pubescence; there is an irregular diamond-shaped mark in its centre and a less distinct black line above the clypeal foveæ.

Ann. & Mag. N. Hist. Ser. 7. Vol. xiii.

15



Clypeus punctured, but not strongly or closely, its apex slightly bent inwardly. Scutellum opaque, closely and uniformly punctured, more strongly than the mesonotum. Metanotum closely punctured (except in the centre at the base); areola longer than broad and of nearly equal width; the teeth broad, large, rounded, and narrowed at the apex. Propleuræ have a plumbeous hue and are irregularly striated at the apex; the meso- closely, the metapleuræ if anything still more closely, punctured, the punctuation running into striæ at the apex. The second to fifth abdominal segments are closely punctured; the gastrocœli striated at the base; the petiole is aciculated in the middle towards the apex.

*Algathia erythropoda*, sp. n.

Black; the face, clypeus in the centre above, the mark rounded at the apex, the basal half of the mandibles, palpi, the inner orbits, the line dilated in front of the ocelli, a line on the pronotum (broad at the base, gradually narrowed to the apex), basal half of tegulæ, tubercles, scutellum (except at the base), postscutellum, the fifth abdominal segment (narrowly in the middle), the apical two thirds of the sixth and seventh entirely, yellow. Scape of antennæ yellow below, thickly covered with white hair; the joints of flagellum dilated below (especially near the apex) and brownish. Legs rufous, the front coxæ and trochanters yellow, the middle coxæ yellow, broadly black at the base behind; the hinder black, with the apical half yellow above; the basal joint of the trochanters, apex of hinder tibiæ, and the hind tarsi black; the spurs pale fulvous. Wings not very clear hyaline, the stigma and nervures black; areolet narrowed in front, being there less in length than the space bounded by the recurrent and second transverse cubital nervures. ♂.

Length 11 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Face coarsely punctured, thickly covered with short white hair; the clypeus less strongly punctured, almost smooth at the apex; the vertex strongly punctured, the front smooth. Mesonotum closely strongly punctured. Scutellum shining, covered with long fuscous hair. Areola broader than long, the sides at the base obliquely truncated, the apex slightly bent inwardly, the sides distinctly depressed, with two or three keels at the apex; the centre raised, rugosely punctured; the posterior median area widened at the base, closely, strongly, transversely striated; the lateral basal areæ coarsely aciculated on the inner side, the rest strongly irregularly

striated;  
rest coarsely  
strongly  
at the base  
keels. U  
aciculated  
strongly  
striated;  
petiole stri  
and obliqu  
abdominal  
covered w  
the inner

Comes  
white band  
yellow, no  
is much s

Black;  
the middle  
what triangular  
ocelli, the  
narrow ab  
teeth), a l  
lums, the  
of the too  
the hind c  
segments  
slightly in  
yellow; th  
pale yellow  
middle ab  
apical joint  
femora, ab  
metatarsus  
dark rufous  
black. ♂

Length

*Hab.* K

Antennæ  
joints whit  
the flagellu  
and vertex  
with short

nt strongly or closely, its apex  
Scutellum opaque, closely and  
e strongly than the mesonotum.  
red (except in the centre at the  
road and of nearly equal width;  
ded, and narrowed at the apex.  
s hue and are irregularly striated  
sely, the metapleuræ if anything  
ed, the punctuation running into  
ond to fifth abdominal segments  
gastrocoeli striated at the base;  
the middle towards the apex.

*Pythopoda*, sp. n.

s in the centre above, the mark  
asal half of the mandibles, palpi,  
lated in front of the ocelli, a line  
the base, gradually narrowed to  
ake, tubercles, scutellum (except  
n, the fifth abdominal segment  
he apical two thirds of the sixth  
ow. Scape of antennæ yellow  
white hair; the joints of flagellum  
ar the apex) and brownish. Legs  
d trochanters yellow, the middle  
at the base behind; the hinder  
yellow above; the basal joint of  
hinder tibiae, and the hind tarsi  
s. Wings not very clear hyaline,  
black; areolet narrowed in front,  
than the space bounded by the  
verse cubital nervures. ♂.

H. Rothney.

thickly covered with short white  
ngly punctured, almost smooth at  
gly punctured, the front smooth.  
y punctured. Scutellum shining,  
hair. Areola broader than long,  
uely truncated, the apex slightly  
distinctly depressed, with two or  
the centre raised, rugosely punc-  
n area widened at the base, closely,  
ed; the lateral basal areae coarsely  
ide, the rest strongly irregularly

striated; the spiracular area behind strongly aciculated, the  
rest coarsely transversely striated; the other basal areae  
strongly but not closely punctured; the apical areae smooth  
at the base, the rest with stout, clearly separated, transverse  
keels. Upper part of propleuræ strongly punctured, the base  
aciculated, the middle below stoutly striated; mesopleuræ  
strongly punctured, the base above and the apex below  
striated; metapleuræ punctured strongly all over. Post-  
petiole strongly and closely punctured, raised in the middle  
and obliquely narrowed at the apex; the second and third  
abdominal segments closely punctured, the apical thickly  
covered with fulvous hair; the gastrocoeli large, the base on  
the inner side with a stout keel, the outer with two keels.

Comes near to *A. parvimaculata*; it is larger, wants the  
white bands on the postpetiole, the hinder trochanters are  
yellow, not black; the black mark on the face and clypeus  
is much smaller and the middle coxæ are yellow.

*Algathia varipes*, sp. n.

Black; the face, clypeus, labrum, the inner orbits from  
the middle of the ocelli to the white face, an oblique some-  
what triangular mark near the eyes opposite the hinder  
ocelli, the lower three fourths of the outer orbits (the mark  
narrow above, broader below), palpi, mandibles (except their  
teeth), a line on the pronotum (curved at the base), scutel-  
lums, the posterior intermedian, the apex of spiracular end  
of the tooth-bearing areae of median segment, a mark above  
the hind coxæ, the apices of the first and second abdominal  
segments broadly (the lines dilated in the middle), the sixth  
slightly in the middle, and the whole of the last segment pale  
yellow; the four front legs fulvous, the coxæ and trochanters  
pale yellow; the hinder coxæ black, broadly yellow in the  
middle above; the basal joint of the trochanters yellow, the  
apical joint and the base of femora rufous, the rest of the  
femora, almost the apical half of the tibiae, and the base of  
metatarsus black; the rest of tarsi white, the hinder spurs  
dark rufous. Wings hyaline, the costa, stigma, and nervures  
black. ♂.

Length 9 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Antennal scape beneath and the fourteenth to twenty-fifth  
joints white; scape thickly covered with long fuscous hair,  
the flagellum with short black pubescence. Face, clypeus,  
and vertex closely and strongly punctured, thickly covered  
with short fuscous hair. Mesonotum closely punctured,

opaque; scutellums shining, smooth, sparsely covered with short fuscous hair; the basal scutellar keel stout, sharp, white at the apex, and extending to the middle. Base of median segment depressed at the base, the furrow smooth, slightly curved, petiolar area open at the base, areola longer than broad, rounded at the base, rounded inwardly at the apex; posterior median area slightly widened at the base, almost smooth, only obscurely transversely striated; the supra-external area has some scattered punctures, the tooth-bearing coarsely punctured, the spiracular finely punctured at the base, the rest obliquely striated; the posterior intermedian coarsely obliquely striated; the teeth indistinct. Propleuræ punctured above, below smooth, the apex in the middle with some stout punctures; mesopleuræ closely punctured, the metapleuræ more strongly, with the punctures somewhat more widely separated.

*Algathia Rothneyi*, sp. n.

Agrees closely in coloration with *A. varipes*, except that it has a yellow mark on the mesopleura; otherwise differing in having the base of the metanotum opaque and aciculated, the keels bordering the petiolar area are longer, the areola is not smooth, the teeth are more distinct, the hinder femora have only the apical third black, and the scutellar keels reach to the apex.

Black; the head below the antennæ, mandibles (except the teeth), palpi, the inner orbits above, the line narrowed below, the apical three fourths of outer orbits, edge of pronotum (the basal half of the line dilated and acute at the base), the tubercles, a mark on the hinder edge of mesopleuræ (longer than wide and obliquely truncated at the apex), scutellums, the sides of the median segment (broadly at the apex), a narrow band on the first abdominal segment (roundly dilated in the middle), the apex of second broadly, and the apical two entirely, pale yellow. Four front legs rufous, their coxæ and trochanters pallid yellow; hinder legs of a deeper red; coxæ black, yellow above and within, the basal joint of trochanters yellow; the apical third of femora and of the tibiæ and the basal two thirds of metatarsus black; the rest of the tarsi pale yellow, except the apex of the last joint. Wings clear hyaline, the stigma testaceous, nervures fuscous. ♀.

Length 8 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Scape below and joints 9-16 of antennæ white, the

flagellum at the clypeus, labrum fringed, ocelli closely shining. Median segment distinctly depressed to the base of side; scutellum notum large, apex; areola obliquely narrowed; posterior median area rounded and opaque; the teeth strongly, irregularly above closely longitudinal, broader than metapleuræ aciculated, striated.

Is very similar, may be known strongly striated, distinctly defined and by the apical black.

Black; face, mandibles, palpi, ocelli, the line below, the petiolar area to the base, the line dilated at mesonotum, the median segment (the yellow dilated sternum, the apical tinged with yellow coxæ and basal tibiæ and the coxæ behind. Wings hyaline.



ing, smooth, sparsely covered with basal scutellar keel stout, sharp, extending to the middle. Base of d at the base, the furrow smooth, area open at the base, areola longer the base, rounded inwardly at the area slightly widened at the base, obscurely transversely striated; the some scattered punctures, the tooth-red, the spiracular finely punctured quely striated; the posterior interely striated; the teeth indistinct. ove, below smooth, the apex in the t punctures; mesopleuræ closely ræ more strongly, with the puncely separated.

*A. Rothneyi*, sp. n.

ation with *A. varipes*, except that the mesopleura; otherwise differing metanotum opaque and aciculated, etiolar area are longer, the areola is e more distinct, the hinder femora black, and the scutellar keels reach

w the antennæ, mandibles (except er orbits above, the line narrowed urther of outer orbits, edge of pro the line dilated and acute at the ark on the hinder edge of meso- e and obliquely truncated at the es of the median segment (broadly nd on the first abdominal segment ddle), the apex of second broadly, ly, pale yellow. Four front legs chanter pallid yellow; hinder legs æk, yellow above and within, the yellow; the apical third of femora e basal two thirds of metatarsus rsi pale yellow, except the apex of ear hyaline, the stigma testaceous,

*A. Rothneyi*.

ts 9-16 of antennæ white, the

flagellum at base and apex brownish beneath. Face closely, the clypeus more sparsely punctured; palpi pale yellow; labrum fringed with long fuscous hair. Vertex below the ocelli closely transversely punctured; front smooth and shining. Mesonotum closely punctured. Scutellum not distinctly depressed at base and apex, its keels sharp and extending to the base of the apical third and depressed on the inner side; scutellum bifoveate at the base. Basal area of metanotum large, aciculated, obliquely narrowed towards the apex; areola nearly as long as wide, the sides at the base obliquely narrowed, its apex slightly roundly turned inwardly; posterior median area smooth, of uniform width and rounded at the base; the other areae closely punctured, opaque; the spiracular and the tooth-bearing areae more strongly, irregularly, and not so closely striated. Propleuræ above closely punctured, the apex broadly, in the middle longitudinally striated; the base with a broad yellow band, broader than the upper one on the pronotum; mesopleuræ closely punctured, the middle behind coarsely aciculated, the metapleuræ at the base aciculated, the rest closely obliquely aciculated. Gastrocoeli shallow, striated; the base rufous, striated.

*Algathia robusta*, sp. n.

Is very similar in form and coloration to *A. maculiceps*; may be known from it by the yellow scutellum, by the strongly striated front and vertex, by the areola being distinctly defined and the posterior median area not at all, and by the apex of the hind tibiæ and the metatarsus being black.

Black; face (except a small black mark above), clypeus, mandibles, palpi, the inner orbits to beyond the hinder ocelli, the lower outer orbits (narrowly above, broadly below), palpi, a narrow line on the pronotum (not reaching to the base), tegulæ, a line on the base of the propleuræ, the line dilated at the base, two short marks in the centre of the mesonotum, scutellums, two broad lines on the sides of median segment, the tubercles, lower part of mesopleuræ (the yellow dilated upwards at the base), the base of mesosternum, the apices of the first to fourth abdominal segments, and the apical entirely, yellow. Four front legs fulvous, tinged with yellow, the apices of the tarsi black; hinder coxæ and basal joint of trochanters, apex of femora, of the tibiæ and the metatarsus, black; on the apex of the hind coxæ behind is a large yellow mark, narrowed behind. Wings hyaline, the nervures and stigma pale testaceous; the



areolet slightly oblique, narrowed in front, the nervures almost uniting there. ♂.

Length 11 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Face closely and strongly punctured, its middle above almost transversely striated; clypeus less strongly punctured, its apex smooth; it is thickly covered with short white, the labrum fringed with long fulvous, hair. Vertex near the eyes punctured, the centre with three keels, running from the hinder ocelli, which turn outwardly to the central furrow, the front finely transversely striated. Mesonotum closely and strongly punctured. Scutellum at base rugosely punctured, the rounded apex closely longitudinally striated; the keels large, black, and reaching to the apical third. Middle of metanotum at base with an oblique slope, the sides margined, very smooth and shining; areola somewhat wider than long, the base rounded, apex transverse, it is irregularly longitudinally rugose, the middle keeled; the lateral areæ rugose, in the centre almost reticulated; teeth large, wide. Propleuræ obscurely punctured above; the mesopleuræ strongly punctured, the middle slightly, the apex more strongly longitudinally striated; the metapleuræ with the spiracular area closely obliquely striated. Postpetiole closely, finely, irregularly striated, the other abdominal segments closely punctured; gastrocœli deep, finely striated at the base; apex aciculated.

*Algathia flavo-balteata*, sp. n.

Agrees in size and colour with *A. robusta*; may easily be separated from it by the form of the scutellum, which has a much sharper oblique slope, by the postscutellum being only obscurely, not strongly striated at the base, by the base of metanotum being aciculated, by the areola being longer compared with its breadth, by the black on the hind femora and tibiæ being less extended, by the hind coxæ being almost entirely yellow above, and by the less distinct curve on the apex of clypeus.

Joints 9-15 of antennæ below, the face and clypeus (except for a small oval mark below the antennæ), mandibles, palpi, inner eye-orbits, the outer from near the top more broadly, white. A narrow line on the pronotum, tegulæ, a large mark near the apex of mesonotum, scutellums, two broad curved lines on the sides of median segment at the apex, a broad line on the lower side of the propleuræ, a broader one (curved upwards at the base on the lower side of the mesopleuræ), a

on new

ed in front, the nervures

hney.

nctured, its middle above  
 eus less strongly punctured,  
 vered with short white, the  
 us, hair. Vertex near the  
 three keels, running from  
 outwardly to the central  
 sely striated. Mesonotum  
 Scutellum at base rugosely  
 sely longitudinally striated;  
 hing to the apical third.  
 h an oblique slope, the sides  
 g; areola somewhat wider  
 ex transverse, it is irregu-  
 middle keeled; the lateral  
 st reticulated; teeth large,  
 ctured above; the meso-  
 middle slightly, the apex  
 ated; the metapleuræ with  
 uely striated. Postpetiole  
 ted, the other abdominal  
 trocœli deep, finely striated

narrower one on the hinder edge, a mark below the hind wings, the basal four abdominal segments at the apex, and the apical two entirely, whitish yellow. Four front legs fulvous; the coxæ and trochanters yellow; the hind coxæ black, the middle above broadly yellow, and there is an elongated mark on the side at the base; the hind femora and tibiæ rufous, as are also the trochanters; the hinder knees, apex of tibiæ, and metatarsus black, the rest of the tarsi white. Wings hyaline, with a slight fuscous tint; stigma testaceous. ♀.

Length 11 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Face closely and strongly punctured; the clypeus sparsely punctured, its apex transverse; the black part of vertex strongly aciculated, the front strongly transversely striated and furrowed down the centre. Propleuræ with a plumbeous hue; the meso- closely, the metapleuræ if anything more strongly, punctured. Scutellum strongly but not closely punctured. Median segment closely punctured; the apex in the centre closely, at the sides much more strongly, transversely striated; the teeth large; areola longer than broad, rounded at the base, slightly narrowed at the apex, which is transverse. Postpetiole shagreened or finely striated in the middle; the gastrocœli large, wide, the base finely striated.

*Algathia femorata*, sp. n.

*teata*, sp. n.

*A. robusta*; may easily be  
 the scutellum, which has a  
 the postscutellum being only  
 at the base, by the base of  
 the areola being longer  
 black on the hind femora  
 the hind coxæ being almost  
 less distinct curve on the

the face and clypeus (except  
 antennæ), mandibles, palpi,  
 near the top more broadly,  
 notum, tegulæ, a large mark  
 tellums, two broad curved  
 ent at the apex, a broad  
 euræ, a broader one (curved  
 side of the mesopleuræ), a

Black; a mark (elongated and rounded at the apex) in the centre of the face above, palpi, scutellum, the outer areæ on the apex of the median segment, the middle of the tubercles, the apex of the first and second abdominal segments, a large triangular mark on the sides of the third, the apex of the penultimate, and the whole of the last segment yellow. Scape of antennæ rufous beneath, the middle of flagellum broadly white. Four front legs fulvous, their coxæ and trochanters yellowish white; the middle tarsi and the hind legs black, except the trochanters, which are yellow, and the extreme apex of the femora, which is rufous; the calcaria white. Wings hyaline, the nervures and stigma black. ♀.

Length 10 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Face strongly, clypeus sparsely punctured. Mesonotum closely and strongly punctured, thickly covered with short thick hair. Scutellum smooth, shining. Areola elongate, the base distinctly, the apex slightly narrowed; the lateral keels received in front of its middle; posterior median area

aciculated, shagreened towards the apex; the lateral area with some rough transverse keels, the spiracular closely punctured, the apex raised on the inner side, transversely striated; the outer two basal area are closely but not strongly punctured, the outer apical have three stout, curved, transverse keels on the apex. The apex of the median segment is thickly covered with long soft white hair. Propleuræ coarsely punctured more finely at the base, the middle below obscurely striated. Mesopleuræ coarsely punctured, below the tubercles obliquely striated; metapleuræ punctured like the mesopleuræ. First segment of abdomen smooth, the depressed sides of the apex punctured; the second and third segments closely punctured, the base of the second striated laterally; gastrocœli not depressed, rufous at the apex. The apical segments of the abdomen are thickly covered with soft white hair.

*Algathia cariniscutis*, sp. n.

Black; the scape of antennæ and joints 8-12 underneath, the face and clypeus (except for a line in the centre of the former, which gets gradually thicker until it reaches the clypeal foveæ, the clypeus being also black in the centre), and the labrum white; the inner orbits broadly to the end of the eyes, the outer more narrowly on the lower half, maxillary palpi, lower part of propleuræ, the edge of pronotum (except at the base), tegulae, tubercles, scutellums, the apical half of median segment, a large oblique mark on the mesopleuræ above the coxæ, a smaller one behind the posterior coxæ, one under the hind wings, the apices of the first and second abdominal segments, a mark on either side of the third, and the apical two segments, yellow. Legs rufous, the four front coxæ and trochanters bright lemon-yellow, the apices of the tarsi fuscous; the hinder coxæ black, broadly yellow above and at the sides above on the inner side, their middle behind next to the black part rufous; the trochanters black, the basal joint for the greater part yellow above; the apex of the hind femora and the base of the tibiae more narrowly black; the hinder tarsi paler, not so rufous in tint as the anterior, their apex black. Wings clear hyaline, the stigma and nervures black. ♀.

Length 8 mm.

*Hab.* Khasia Hills. Coll. Rothney.

Face and clypeus closely punctured, covered with short white down; the front and vertex almost impunctate. Mesonotum closely and uniformly punctured; scutellum smooth, covered with fuscous pubescence, its sides stoutly

keeled  
base,  
media  
apical  
upper  
mesop  
punct  
petiole  
and th  
wide,

XXIV  
of L  
Pos  
M.A

It is  
genetic  
and th  
ment  
Order  
of com  
paired  
been a  
some c  
the su  
upon th

The  
equal  
form t  
exampl  
much t  
branch  
bivittate  
of *Pelop*  
which I

Secor  
in a fa  
In most

\* 'Cat  
London.  
† Bron  
pl. cxxxiv  
† *ibid.*



apex; the lateral arcæ  
the spiracular closely  
inner side, transversely  
are closely but not  
ave three stout, curved,  
ex of the median seg-  
white hair. Propleuræ  
base, the middle below  
sely punctured, below  
pleuræ punctured like  
abdomen smooth, the  
; the second and third  
of the second striated  
fous at the apex. The  
ickly covered with soft

sp. n.

oints 8-12 underneath,  
ne in the centre of the  
r until it reaches the  
lack in the centre), and  
oadly to the end of the  
e lower half, maxillary  
e edge of pronotum  
, scutellums, the apical  
e mark on the meso-  
e behind the posterior  
pices of the first and  
on either side of the  
yellow. Legs rufous,  
bright lemon-yellow,  
e hinder coxæ black,  
es above on the inner  
lack part rufous; the  
he greater part yellow  
d the base of the tibiæ  
si paler, not so rufous  
black. Wings clear  
♀.

, covered with short  
almost impunctate.  
unctured; scutellum  
ence, its sides stoutly

keeled. Arcola slightly wider than long, rounded at the base, the apex bulging inwardly; the base of posterior median area smooth, the rest transversely striated; the outer apical arcæ are more strongly and widely striated. The upper half of propleuræ closely punctured, as are also the mesopleuræ; the metapleuræ are more closely and strongly punctured. First segment of abdomen aciculated, the post-petiole more strongly and raised in the middle; the second and third segments are closely punctured; the gastrocœli wide, striated, the oblique apex aciculated.

XXIV.—*Preliminary Note on certain Points in the Anatomy of Eryx and other Boidæ, partly indicative of their Basal Position among the Ophidia.* By FRANK E. BEDDARD, M.A., F.R.S.

It is generally believed that the Boidæ occupy phylogenetically a place at or near the base of the Ophidian series; and this view is expressed by Boulenger in a tabular statement of the mutual affinities of the various families of the Order\*. This opinion is largely based upon the persistence of considerable vestiges of the pelvic girdle and upon the paired lungs. In studying the anatomy of snakes I have been able to note a few other points to which little or, in some cases, no attention has been paid and which tend to the support of this conclusion. My observations bearing upon this subject were made upon *Python*, *Eryx*, and *Boa*.

The first point to which I would draw attention is the equal size of the right and left aortic arches, which join to form the dorsal aorta. In at least many other snakes (for example, *Zamenis flagelliformis*) the right aortic arch is so much the smaller that it appears almost as an inconspicuous branch of the left. It would appear, however, that in *Python bivittatus* this is not the case †, though Dr. Gadow's drawing ‡ of *Pelophilus madagascariensis* is in accordance with the facts which I have observed.

Secondly, the intercostal branches of the aorta are arranged in a fashion which appears to me to be distinctly archaic. In most snakes the intercostal arteries are very irregular in

\* 'Catalogue of the Snakes in the British Museum (Natural History),' London, 1893, vol. i. p. 2.

† Bronn's 'Klassen und Ordnungen des Thierreichs,' Bd. vi. Abth. iii. pl. cxxxiv. fig. 2. This figure is copied from Fritsch.

‡ *Ibid.* pl. cxxxv. fig. 1.