



Cantaloupe (Muskmelon) in the Garden

Rick Hefelbower and Dan Drost

Summary

Cantaloupes grow best in sunny locations and in fertile, well-drained soils. Incorporate organic matter and a complete fertilizer into the area before planting. Plant 4-6 cantaloupe seeds directly in the garden when soils are 65°F. Seeds should be planted 1-2 inches deep, in mounds 4 feet apart. Thin the mounds after emergence to 2 plants. Transplant cantaloupe 2 feet apart through black plastic for early maturity. Use row covers or hot caps to protect the plants when planting before the frost-free period. After the vines develop runners, side dress with additional nitrogen fertilizer. Irrigation should be deep and infrequent. Plastic and organic mulches help conserve water and reduce weeding, however, do not apply organic mulches until soils have warmed to 75°F. Control insect and diseases throughout the year. Harvest cantaloupes when the fruits separate from the vine easily and the background color is creamy yellow. Excellent varieties include Ambrosia, Mission, Summit Hybrid, and Rocky Sweet.

Varieties

Ambrosia, Classic Hybrid, Hales Best, Mission, Rocky Sweet and Summit Hybrid are excellent cantaloupe varieties.

How to Grow

Soil: Cantaloupes prefer organic, rich, well-drained, sandy soils for best growth. Most soils will grow cantaloupe provided they are well-drained.

Soil Preparation: Choose a site in your garden that receives full sun. Before planting, determine fertilizer needs with a soil test and then follow the recommendations given with the test report. If

fertilizer applications are warranted, work the fertilizer into the top 6 inches of soil. If you fertilize with compost, apply no more than 1 inch of well-composted organic matter per 100 square feet of garden area.

Plants: Cantaloupe can be grown from seed or transplants. Seed should be planted 1-2 inches deep. Transplants should have 2-3 mature leaves and a well-developed root system. Allow 6 weeks to grow transplants. Transplants mature about 2 weeks before seeded melons and should be used in short growing areas of Utah.

Planting and Spacing: Cantaloupe should be planted when soils are 65°F or after all frost danger has past. Plant 4-6 seeds in mounds 4 feet apart. After they have two leaves, thin to 2 plants per mound. Transplants should be planted 2 feet apart in row, with rows 4-6 feet apart. Avoid damaging the roots when planting which slows establishment and growth.

Mulch: Black plastic mulch warms the soil, conserves water and helps control weeds. Plastic mulches allow earlier planting and maturity, especially with transplants. After laying out the mulch, secure the edges with soil and cut holes for seeds or transplants. When using plastic mulches and row covers, seeds or plants can be set out about 2 weeks before the last frost date. Do not apply organic mulches (grass clippings, straw, newspapers, etc.) until soils are warmer than 75°F. Both plastic and organic mulches help conserve water and control weeds.

Row covers: Row covers enhance growth and earliness. Hotcaps, plastic tunnels, fabric covers,

and other devices help protect seedlings and transplants from cool air temperatures. Use a thermometer to help determine the temperature under row covers. Plants grown under row covers require ventilation when air temperatures exceed 80°F. Covers need to be removed when plants start to flower or when temperatures exceed 90°F. Remove covers when weather has stabilized.

Irrigation: Water cantaloupe deeply and infrequently, 1-2 inches per week. Use drip irrigation if possible. Mulch around the plants will help conserve soil moisture and reduce weed growth. Irrigate so that moisture goes deeply into

the soil. Reduce watering amount as the fruits ripen to improve flavor.

Fertilizer: After the vines develop runners, side dress each plant with 3-4 tablespoons of a nitrogen fertilizer (21-0-0), sprinkled around the plant, then water in the fertilizer.

Problems

Weeds: Plastic and organic mulches effectively control weeds. Healthy vigorous vine growth by cantaloupe will also smother weeds.

Insects and Diseases: For more detailed information on insect and diseases visit the Utah Pests website (www.utahpests.usu.edu).

Insect	Identification	Control
Aphids	Green or black soft-bodies insects that feed on underside of leaves. Leaves become crinkled and curled. May transmit virus diseases. Secreted honeydew makes plants appear shiny, wet or sticky.	Use insecticidal soaps or strong water stream to dislodge insects.
Cucumber Beetles	Adults have stripes or spots and feed on leaves and vines which reduces vigor. They transmit bacterial disease. Larvae bore into roots and stems causing plants to wilt and die.	Application of chemicals at first appearance is needed to control this pest.
Disease	Symptom	Control
Powdery Mildew	White fungal patches start on older leaves. The disease eventually spreads to all plant parts. The foliage dies, exposing fruits to the sun, which causes premature ripening.	Plant resistant varieties.
Wilt Diseases	Leaves wilt on one or more vines. Plants often die. Streaking, slime formation, or gummy exudates visible on stems. Diseases are caused by different pathogens.	Identify causal disease. Treat disease as recommended once identified.
Virus	Leaves are light green, mottled, malformed, dwarfed and curled. Early infection affects fruit shape and flavor. An aphid transmitted disease.	Control aphids. Destroy severely infected plants.

Harvest and Storage

Cantaloupe require 35-45 days to mature from flowering, depending on the temperature and variety. Use the following guide to determine fruit maturity. The netting gets coarse and rough, the stem breaks (slips) away from the vine easily, and the background color of the fruit turns from green to yellow (see photo). Pick melons as they ripen. Cantaloupe will store for 1-2 weeks if held at 45-50°F.



Productivity

Plant 3-4 cantaloupe per person for fresh use and an additional 3-4 plants for juicing or freezing. Expect 100 fruits per 100 feet of row.

Nutrition

Cantaloupe are mostly water. A quarter of a melon has about 50 calories, is low in fat and is an excellent source of vitamin A and vitamin C.

Frequently Asked Questions

Why do the first blossoms drop off my muskmelon plants? The first flowers to appear on the vines are male. The female flowers, which open later, have a swelling at the base that forms the fruit. After bees pollinate these female flowers, the fruit develops.

What causes poor fruit set and low yields? The failure for female flowers to set and develop melons is due to a lack of proper pollination by bees, very hot weather or water stress.

How can I grow muskmelons in a small garden?

Cantaloupe plants can be trained to a fence or trellis or grown in a large pot. After the fruits begin to enlarge they will need some support or the fruit weight may damage the vines.

Do muskmelons cross-pollinate with other vine crops? No. Cantaloupe do not cross-pollinate with cucumbers, watermelon, squash or pumpkins. Different cantaloupe varieties do cross-pollinate, but the effect of this is not evident unless the seeds are saved and planted the following year.

This project is funded in part by USDA-Risk Management Agency and the Utah Department of Agriculture and Foods Specialty Crops Block Grant (SCBG 161039) under a cooperative agreement. The information reflects the views of the author(s) and not USDA-RMA or UDAF.

In its programs and activities, Utah State University does not discriminate based on race, color, religion, sex, national origin, age, genetic information, sexual orientation or gender identity/expression, disability, status as a protected veteran, or any other status protected by University policy or local, state, or federal law. The following individuals have been designated to handle inquiries regarding non-discrimination policies: Executive Director of the Office of Equity, Alison Adams-Perlac, alison.adams-perlac@usu.edu, Title IX Coordinator, Hilary Renshaw, hilary.renshaw@usu.edu, Old Main Rm. 161, 435-797-1266. For further information on notice of non-discrimination: U.S. Department of Education, Office for Civil Rights, 303-844-5695, OCR.Denver@ed.gov. Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Kenneth L. White, Vice President for Extension and Agriculture, Utah State University.

