

DOUBLE-CRESTED CORMORANT AND RING-BILLED GULL DAMAGE MANAGEMENT ON LAKE CHAMPLAIN: ARE BASIN-WIDE OBJECTIVES ACHIEVABLE?

RICHARD CHIPMAN, U.S. Department of Agriculture, Animal and Plant Health Inspection Service,
Animal Damage Control, P.O. Box 1436, Montpelier, VT 05602

DENNIS SLATE, U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Animal
Damage Control, Whitebridge, 91A North State Street, Concord, NH 03301

LARRY GARLAND, Vermont Fish and Wildlife Department, 111 West Street, Essex Junction, VT
05452

DAVID CAPEN, School of Natural Resources, Aiken Center, University of Vermont, Burlington, VT
05402

Abstract: Ring-billed gulls (*Larus delawarensis*) and double-crested cormorants (*Phalacrocorax auritus*) have nested on Lake Champlain since 1949 and 1982, respectively. Recent increases in cormorant nesting populations and pioneering activities of both species to previously uncolonized islands have resulted in impacts related to accumulation of bird guano and interspecific competition with less common species. Of primary concern are: decreases in wildlife and plant diversity on islands; reduced aesthetics and property values of island associated with the loss of trees; and predation or competition for nesting space with other species such as the state-endangered common tern (*Sterna hirundo*). In addition, public concerns have been raised over potential impacts to fish species by cormorants, as well as urban-suburban gull activity impacting human health and safety and property. Nine stakeholder meetings have been held since 1990, to consider prospective management strategies for gulls and cormorants on Lake Champlain, Vermont. Since 1994, limited site-specific control efforts have been conducted by USDA-APHIS-ADC (ADC) at the request of the Vermont Fish and Wildlife Department (VFWD) and private landowners who recently experienced damage from these species. A technique for the removal of cormorant nests in trees has been developed and tested by ADC using portable pumps to deliver pressurized water to remove the nests. This technique has proven successful in tree nests up to approximately 15 m. Also, an alternative method is being tested to selectively remove nesting cormorants from a great blue heron (*Ardea herodias*) rookery. Currently, a draft environmental assessment is being prepared cooperatively by VFWD and ADC that explores alternatives for future management over 5 years beginning in 1998. Extensive public involvement using a variety of approaches will occur during the summer of 1997. VFWD, in cooperation with ADC, will continue to approach damage management on Lake Champlain for these species on a site-specific basis while working toward a more comprehensive interstate and international agreement on goals for basin-wide population and ecosystem management.

Key Words: ADC, double-crested cormorant, Lake Champlain, *Larus delawarensis*, nest removal, *Phalacrocorax auritus*, ring-billed gull, stakeholders

Proc. East. Wildl. Damage Manage. Conf. 8:52
