

## Commentary

### Can the vaquita be saved from extinction?

**GERARDO RODRÍGUEZ-QUIROZ**, Instituto Politécnico Nacional, Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional, Unidad Sinaloa. Blvd. Juan de Dios Bátiz Paredes No. 250, Guasave, Sinaloa 81101, Mexico [grquiroz@ipn.mx](mailto:grquiroz@ipn.mx)

**WENCESLAO VALENZUELA-QUIÑONEZ**, Instituto Politécnico Nacional, Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional, Unidad Sinaloa. Blvd. Juan de Dios Bátiz Paredes No. 250, Guasave, Sinaloa 81101, Mexico

**HÉCTOR A. GONZÁLEZ-OCAMPO**, Instituto Politécnico Nacional, Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional, Unidad Sinaloa. Blvd. Juan de Dios Bátiz Paredes No. 250, Guasave, Sinaloa 81101, Mexico

**ALFREDO ORTEGA-RUBIO**, Centro de Investigaciones Biológicas del Noroeste, Unidad Sonora (CIBNOR), Carretera a San Juan de La Costa “EL COMITAN” Km. 1, Playa Palo de Santa Rita Sur, 23096 La Paz, Baja California Sur, Mexico

**Abstract:** The vaquita (*Phocoena sinus*) is considered the world’s most endangered marine mammal. It is the smallest member of the porpoise family endemic to the upper part of the Gulf of California. The current population is estimated at <30 individuals. The primary reasons for the species decline includes limited habitat and incidental mortalities associated with illegal gillnet fishing activities. Since 2008, the Mexican government has taken environmental and economic actions to protect the vaquitas, focusing on reducing bycatch deaths to zero. In 2015, a federal agreement decreed by the Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación prohibited the use of any fishing gillnets for 2 years, severely affecting local human communities because coastal fisheries in the region represent 40% of the gross domestic product and 50% of the local inhabitants are devoted to this activity. Recently, an economic welfare compensation program is giving monthly to fishermen who have fishing permits if they do not continue with their fishing activities. However, none of these actions have fully considered the range of social and economic solutions for the local inhabitants of this region. The paradigms of the contemporary conservation programs must also focus on the well-being of local fishing communities to prevent the vaquita from becoming the second marine mammal species to disappear due to human activities.

**Key words:** artisanal fisheries, conservation, marine mammal, Mexico, natural protected areas, *Phocoena sinus*, porpoise, Upper Gulf of California, vaquita

**MARINE MAMMALS** are subject to high mortality from accidental capture or bycatch in fisheries (D’agrosa et al. 2000). Global bycatch of marine mammals likely numbers in the hundreds of thousands annually, mostly in gillnets, because they are relatively inexpensive to purchase and can be used in small vessels at little cost (Read et al. 2006). The extinction of the freshwater dolphin baiji (*Lipotes vexillifer*) endemic to the Yangtze River in China was due to fishery bycatches (Read 2008). Highlighting the need to reevaluate strategies intended to reduce marine megafauna bycatch, Senko et al. (2014) suggested more focus should be placed on the exploitation and commercialization of new local fish species, the transformation of the fishing gears supported by an adequate source

funding (Read 2008), and long-term financial strategies to new job alternatives.

The vaquita (*Phocoena sinus*; Norris and McFarland 1958; Figure 1), is endemic to the Upper Gulf of California, and it is critically endangered because of its very small population size and the reduced range of its habitat (Jaramillo-Legorreta et al. 2017). The vaquita suffers from accidental deaths in gillnets because this species shares its habitat with several lucrative fisheries as the totoaba (*Totoaba macdonaldi*), which is highly desired because its bladder is in high demand in the Asian market (D’agrosa et al. 2000, Rojas-Bracho et al. 2006). A passive acoustic monitoring program conducted by the Comité Internacional para la Recuperación de la Vaquita (CIRVA; International Committee



**Figure 1.** Vaquita (*Phocoena sinus*) swimming in the Upper Gulf of California (photo by P. Olson).

for the Recovery of the Vaquita) in 2015 estimated that the entire world populations may consist of <30 individuals (Jaramillo-Legorreta et al. 2017).

The vaquita shares the area where fishermen from 3 Mexican communities, including San Felipe, Baja California; Golfo de Santa Clara, Sonora; and Puerto Peñasco, Sonora (Figure 2) fish for bigeye croaker (*Micropogonias megalops*), Gulf corvine (*Cynoscion othonopterus*), elasmobranchs and blue shrimp (*Litopenaeus stylirostris*). The fishermen's gross profits are related to shrimp fishery, the most economically important fishery, followed by the Gulf corvine. These 2 resources provide employment for fishermen from September to April (Rodríguez-Quiroz et al. 2010). Income for 71% of the fishermen ranged from US\$150–300 per week during the fishing season, and 87% stated that they received <US\$100 per week in other activities when the shrimp and Gulf corvine season ends. These communities depend economically on these 4 fisheries, as alternative jobs are few and poorly paid (Rodríguez-Quiroz et al. 2010).

The World Wildlife Fund (WWF) in 2006 and the Centro de Investigaciones Biológicas del Noroeste in 2012 interviewed approximately 10% of the total fishermen in the 3 fishing communities. Approximately 60% of the fishermen in San Felipe, 50% in Puerto Peñasco, and 40% in Santa Clara stated that they would not stop fishing because it is the only activity in which they feel comfortable and have done for years (Rodríguez-Quiroz et al. 2010).


Redford and Taber (2000) and Knight (2006) argued that to achieve species conservation, all the related social, economic, and cultural communication issues must be addressed. In the case of conserving the vaquita, this implies the development of economically

feasible alternative sources of income for the local inhabitants. This was not the case for the establishment of the biosphere reserve in the Upper Gulf of California, where fishermen's economic necessities and knowledge of the sea were not considered.

Afflerbach et al. (2013) analyzed all the buyout programs implemented in the region and reported that the programs were failing because they undervalued the role of fishing to the livelihood of the communities. The programs focused on changing the fishery activities by providing the fishermen with other economic activities such as tourism and grocery retail (Valdéz-Gardea 2010) without providing them the training needed to be successful in these new alternative jobs.

The Mexican Government has implemented several environmental and economic actions to protect the vaquita. These include: 1) creating the Upper Gulf of California and Colorado River Delta Biosphere Reserve (UGCCRDBR) in 1993 (Diario Oficial de la Federación [DOF] 1993; Figure 1); 2) creating the Vaquita Protected Area (refuge) in 2005, which limits fishing activities (DOF 2005); and 3) through the Secretaría de Medio Ambiente, Recursos Naturales (SEMARNAT), launching the Programa de Acción para la Protección de la Especie: Vaquita program (Program Action for the Conservation of the Species: Vaquita; [PACE]) in 2008 (Avila-Forcada et al. 2012), which was an economic and voluntary program that considered compensation for reducing the fishing effort through the elimination of shrimp driftnet and finfish gillnets (PACE 2008; Morzaria-Luna et al. 2012).

More recently, in 2015 the Mexican Government, through the Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación (SAGARPA) promulgated the "agreement that commercial fishing is temporarily suspended using gillnets, formwork, and/or lines operated with small boats in the Northern Gulf of California." The agreement addressed 4 specific issues: 1) the expansion of the vaquita polygon (the area where the vaquita has been seen the most; see DOF 2005); 2) a 2-year closed season for all fisheries in the site, and providing a monetary compensation to the affected fishermen with fishing permits; 3) an increased enforcement to prevent illegal



**Figure 2.** Location of the Biosphere Reserve of the Upper Gulf of California and Colorado River Delta and the new Vaquita (*Phocoena sinus*) Protection Area. Source: Diario Oficial de la Federación, April 10, 2015.

fisheries; and 4) the introduction of new and ecological gears (DOF 2015; Figure 1).

However, none of these cumulative measures have reversed the vaquita's population decline. Furthermore, these actions have resulted in reductions in fishermen employment and income (Morzaria-Luna et al. 2012) mainly because fishermen without permits and industrial fishery employees were not considered for the economical compensation. These restrictions did not consider catch revenue from other gears or alternative economic activities (Instituto Nacional de Ecología [INE]-WWF) 2006, Aragon-Noriega et al. 2010). The first management actions have not produced the desired conservation results because nobody (governmental and scientific authorities) anticipated that the new job alternatives and fishing gears could fail so unexpectedly.

The SEMARNAT agreement of April 2015

also did not fully consider a social and economic analysis of their conservation research strategies. Government buyout programs have to be adequately enforced so fishermen can find new alternative jobs related to their activity, emphasizing that fishermen are among the social groups in the primary sector with the highest socioeconomic problems and most of them are below the poverty line (Fernández-Carvajal 2013). Additionally, we observed that the lack of an inclusive vision of the human element in the design and act of these reserves by the government institutions has reached levels that encourage poaching of protected species, such as the totoaba, due to the lack of alternative livelihoods for fishermen, who are forced to perform other economic activities to complete their family income for their survival (Rodríguez-Quiroz et al. 2015).

Eliminating all mortality associated with bycatch is an imperative to avert vaquita extinction. We support the

recommendations of scientific organizations such as CIRVA, which suggests that the Mexican Government must enact emergency regulations establishing a gillnet exclusion zone encompassing the full range of the vaquita that is wider than the existing refuge (Gerrodette and Rojas-Bracho 2011, CIRVA-VII 2017). We acknowledge these measures will exacerbate economic impacts that potentially lead to costly and socially divisive conflicts, particularly in regions that rely heavily on fishing or areas where alternative employment opportunities are poor (Mardle and Pascoe 2002).

### Conservation and economic alternatives

Mexican authorities and scientific experts agree that the most immediate actions needed to prevent the extinction of the vaquita are effective enforcement of the fishing ban, continuous

removal of nets that evade the authorities, and the development and implementation of alternative light fishing gear (CIRVA-IV 2012, CIRVA-VII 2017), but these actions will not be fully implemented without changing the fishing culture of the inhabitants (PACE 2008, Rodríguez-Quiroz et al. 2010). To promote fishing culture change for a sustainable fishery in the region and the successful recovery of the vaquita, we make the following recommendations.

1. Adult totoaba could be available for sport fishing with the implementation of management units for wildlife conservation (UMAs) and supported by the Priority Species Recovery Program (PREP) as an alternative to prevent the illegal fishery of this species. The implementation of these programs will allow for the conservation of this species and its exploitation, and it will improve organized fishermen's welfare because fishermen involved in this activity will have benefits in the auction of the permits to fish the totoaba, as is done with the Mexican bighorn (*Ovis canadensis*) and the ejidatarios (Cariño et al. 2004). This will grant a better advantage for the exploitation of resources through controlled management with conservation of the natural resources in the UGCCRDBR (SEMARNAT 2008, Retes López et al. 2010). Recently, Chinese, Mexican, and U.S. authorities had a first meeting in Ensenada, México (SEMARNAT 2017) to implement measures to stop totoaba bladder commercialization (whose exploitation is banned [DOF 1975]), which is not supported by correct ecological fishing and trade guidelines in or outside of Mexico. The Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) and other ecological trade programs like the Marine Stewardship Council could be an option to support the meat and bladder commercialization (Bellchambers et al. 2015).

The vaquita and totoaba share a wide range of common habitat. Thus, emphasizing the UMA program with the totoaba could result in a significant number of traditional fishermen who will change to totoaba sport fishing, reducing vaquita bycatches, and these fishermen will help to enforce regulations on the illegal fishermen, thereby further reducing vaquita bycatches.

2. New ecological or environmentally friendly fishing gears like "suriperas" (modified cast

net) to fish marine species in the biosphere reserve, as suggested by the PACE program, must be given a high product value (Vaquita Safe label), so the use of such gears should equal the amount captured and product value of the marine products they fished with their current fishing gears (Balmori Ramírez et al. 2006, PACE 2008, Instituto Nacional de la Pesca [INAPESCA] 2011). Fishermen should not just receive financial support to change fishing gears (PACE 2008) because, as mentioned in CIRVA-IV (2012), "The conversion of the entire fishing fleet to vaquita-safe methods needs to be accomplished as soon as possible." But this has not been accomplished yet, and the "suriperas" is more effective for brown shrimp than for blue shrimp (INAPESCA 2011), although the latter has a higher price in the international market (Godoy 2008).

3. Government and traders must invest to transform fishing products into high-value products so they can be widely commercialized by establishing long-term contracts with regional and international markets. Such action would reduce fishing effort because there will be well-paid employees in the fish market trade, and that will ensure sustainable fishing activity, particularly for the community of El Golfo de Santa Clara.

4. Even when PACE establishes that fishermen need new job alternatives, fishermen must be well trained in their new qualifications, because many fishermen do not have technical business skills or experience in tourism administration or any other alternative job outside of fishing. Also needed is financial insurance or funds supported by the national and international commerce chambers to prevent a massive return of fishermen to their previous activity, as it is for cattle and farming activity. Today, many of the new alternative jobs like tourism and aquaculture resulted in failures because fishermen were not trained and well-advised, as happened in San Felipe and Golfo de Santa Clara.

## Conclusion

In summary, 2018 will be a crucial year for authorities to find a solution to the demise of the remaining vaquitas. The new agreement published in 2015 does not consider real complementary economic alternatives to support traditional fishermen when there is a

considerable local market for fish and shrimp. We are certain that the application of our 4 recommendations will help to save the vaquita from extinction. Fishermen must be involved in the conservation decision-making beyond being asked to change their fishing gears or increasing the fishing ban in the Upper Gulf of California area, and they must be well-trained and advised in their new alternative jobs.

### Acknowledgments

The authors thank all fishermen who supported our field work. GRQ thanks CONACYT Grant 48445 and the IPN for their support. We thank an anonymous reviewer, and especially T. Messmer, HWI editor-in-chief, who significantly improved an earlier version of this manuscript. This research is the result of the work performed by the Network of Thematic Research in Natural Protected Areas (RENANP) of CONACYT Project 293368.

### Literature cited

- Afflerbach, J., A. Broderick, D. J. Brunkow, S. Herron, J. Sainz, and S. Sohrabian. 2013. An analysis of bioeconomic tradeoffs in vaquita conservation policies. Group project report 2013, Bren School of Environmental Science and Management, University of California Santa Barbara, California, USA.
- Aragón-Noriega, E. A., G. Rodríguez-Quiroz, M. A. Cisneros-Mata, and A. Ortega-Rubio. 2010. Managing a protected marine area for the conservation of critically endangered vaquita (*Phocoena sinus* Norris, 1958) in the Upper Gulf of California. *International Journal of Sustainable Development and World Ecology* 17:410–416.
- Avila-Forcada S., A.L. Martinez-Cruz, and C. Muñoz-Piña. 2012. Conservation of vaquita marina in the Northern Gulf of California. *Marine Policy* 36:613–622.
- Balmori Ramírez, A., J. T. Silva Ramírez, E. Miranda Mier, and A. Rodríguez. 2006. Evaluación de sistemas de captura para una pesca artesanal de camarón eficiente y amigable con el medio ambiente en el alto Golfo de California. Instituto Nacional de la Pesca and World Wildlife Fund Programa Golfo de California, Guaymas, Sonora, Mexico.
- Bellchambers, L. M., B. F. Phillips, and M. Pérez-Ramírez. 2015. From certification to recertification the benefits and challenges of the Marine Stewardship Council: a case study using lobsters. *Fisheries Research* 182: 88–97.
- Cariño, M., J. S. Aceves, C. Rendón, C. Valiente, M. L. Leal, and B. Rodríguez. 2004. La política ambiental mexicana y la conservación del ambiente en Baja California Sur. *Gaceta Ecológica* 70:45–56.
- CIRVA-IV (Comité Internacional para la Recuperación de la Vaquita). 2012. Report of the fourth meeting of the Comité Internacional para la Recuperación de la Vaquita. Scientific Committee Report (SC/66b), Ensenada Baja California, México.
- CIRVA-VII (Comité Internacional para la Recuperación de la Vaquita). 2017. Report of the seventh meeting of the Comité Internacional para la Recuperación de la Vaquita. Scientific Committee Report (SC/66b), Ensenada Baja California, México.
- D'agrosa, C., C. E. Lennert-Cody, and O. Vidal. 2000. Vaquita bycatch in Mexico's artisanal gillnet fisheries: driving a small population to extinction. *Conservation Biology* 14:1110–1119.
- Diario Oficial de la Federación (DOF). 1975. ACUERDO que establece veda para la especie totoaba, *Cynoscion macdonaldi*, en aguas del Golfo de California, desde la desembocadura del Río Colorado hasta el Río Fuerte, Sinaloa en la costa oriental, y del Río Colorado a Bahía Concepción, Baja California, en la costa occidental, Agosto de 1975. Secretaría de Gobernación, Mexico City, Mexico.
- Diario Oficial de la Federación (DOF). 1993. DECRETO por el que se declara área natural protegida con el carácter de Reserva de la Biosfera, la región conocida como Alto Golfo de California y Delta del Río Colorado, ubicada en aguas del Golfo de California y los municipios de Mexicali, Baja California, de Puerto Peñasco y San Luis Río Colorado, Sonora, Junio de 1993. Secretaría de Gobernación, Mexico City, Mexico.
- Diario Oficial de la Federación (DOF). 2005. Programa de protección de la vaquita dentro de área de Refugio ubicada en la porción occidental del Alto Golfo de California, Septiembre del 2005. Secretaría de Gobernación, Mexico City, Mexico.
- Diario Oficial de la Federación (DOF). 2015. ACUERDO por el que se suspende temporal-

- mente la pesca comercial mediante el uso de redes de enmalle, cimbras y/o palangres operadas con embarcaciones menores, en el Norte del Golfo de California, Abril del 2015. Secretaría de Gobernación, Mexico City, Mexico.
- Fernández-Carvajal, D. 2013. Pesca artesanal y pobreza en comunidades aledañas al Golfo de Nicoya. *Revista de Ciencias Sociales* 140(2):137–152.
- Gerrodette, T., and L. Rojas-Brancho. 2011. Estimating the success of protected areas for the vaquita porpoise (*Phocoena sinus*). *Marine Mammal Science* 27:E101–E125.
- Godoy, E. 2008. Camarones congelados. *Revista Poder y Negocios* 50–53.
- Instituto Nacional de Ecología-World Wildlife Fund (INE-WWF). 2006. Conservación de la vaquita y su hábitat en el Alto Golfo de California. Fondo de compensación económica para la pesca con redes de enmalle y el fomento de alternativas a la pesca en el Alto Golfo de California. Instituto Nacional de Ecología, Dirección General de Investigación en Política y Economía Ambiental, WWF programa, México.
- Instituto Nacional de la Pesca (INAPESCA). 2011. Evaluación biotecnológica de la red de arrastre prototipo “RS-INP-MEX” para captura de camarón en el Alto Golfo de California. Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación, Mexico City, Mexico.
- Jaramillo-Legorreta, A., G. Cardenas-Hinojosa, E. Nieto-García, L. Rojas-Bracho, L. J. Ver Hoef, J. Moore, N. Tregenza, J. Barlow, T. Gerrodette, L. Thomas, and B. L. Taylor. 2017. Passive acoustic monitoring of the decline of Mexico’s critically endangered vaquita. *Conservation Biology* 31:183–191.
- Knight, A. T. 2006. Failing but learning: writing the wrongs after Redford and Taber. *Conservation Biology* 20:1312–1314.
- Mardle, S., and S. Pascoe. 2002. Modelling the effects of trade-offs between long and short-term objectives in fisheries management. *Journal of Environmental Management* 65:49–62.
- Morzaria-Luna, H. N., C. H. Ainsworth, I. C. Kaplan, P. S. Levin, and E. A. Fulton. 2012. Exploring trade-offs between fisheries and conservation of the vaquita porpoise (*Phocoena sinus*) using an Atlantis ecosystem model. *PLOS ONE*: 7:e42917.
- Programa de Acción para la Protección de la Especie: Vaquita (PACE). 2008. Estrategia integral para el manejo sustentable de los recursos marinos y costeros en el Alto Golfo de California. Secretaría de Medio Ambiente, Recursos Naturales, Mexico City, México.
- Read, A. J. 2008. The looming crisis: interactions between marine mammals and fisheries. *Journal of Mammalogy* 89:541–548.
- Read, A. J., P. Drinker, and S. Northridge. 2006. Bycatch of marine mammals in U.S. and global fisheries. *Conservation Biology* 20:163–169.
- Redford, K. H., and A. Taber. 2000. Writing the wrongs: developing a safe-fail culture in conservation. *Conservation Biology* 14:1567–1568.
- Retes López, R., M. I. Cuevas González, S. Moreno Medina, F. G. Denogean Ballesteros, F. Ibarra Flores, and M. Martín Rivera. 2010. Unidad de manejo para la conservación de la vida silvestre como alternativa para “los nuevos agronegocios.” *Revista Mexicana de Agronegocios* 27:336–346.
- Rodríguez-Quiroz, G., E. A. Aragón-Noriega, W. Valenzuela-Quifóñez, and H. M. Esparza-Leal. 2010. Artisanal fisheries in the conservation zones of the Upper Gulf of California. *Revista de Biología Marina y Oceanografía* 45:89–98.
- Rodríguez-Quiroz, G., H. A. González-Ocampo, E. Barba-Macías, L. Jiménez-Badillo, M. J. Pinkus-Rendón, M. A. Pinkus-Rendón, and A. Ortega-Rubio. 2015. Aspectos socioeconómicos de la pesca artesanal en las áreas naturales protegidas. Pages 180–195 in A. Ortega-Rubio, M. J. Pinkus-Rendón, and I. C. Espitia-Moreno, editors. *Las áreas naturales protegidas y la investigación científica en México*. Centro de Investigaciones Biológicas del Noroeste SC., La Paz B.C.S., Universidad Autónoma de Yucatán, Mérida, Yucatán and Universidad Michoacana de San Nicolás de Hidalgo, Michoacán, Mexico.
- Rojas-Bracho, L., R. R. Reeves, and A. M. Jaramillo-Legorreta. 2006. Conservation of the vaquita *Phocoena sinus*. *Mammal Review* 36:179–216.
- Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT). 2008. Programa de acción para la conservación de la especie vaquita (*Phocoena sinus*). SEMARNAT, Mexico City, Mexico.
- Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT). 2017. México, China

y Estados Unidos hacen frente común para combatir el tráfico ilegal de totoaba. SEMARNAT, Mexico City, Mexico, <<https://www.gob.mx/semarnat/prensa/mexico-china-y-estados-unidos-hacen-frente-comun-al-trafico-ilegal-de-totoaba>>. Accessed April 20, 2018.

Senko, J., E. R. White, S. S. Heppell, and L. R. Gerber. 2014. Comparing bycatch mitigation strategies for vulnerable marine megafauna. *Animal Conservation* 17:5–18.

Valdéz-Gardea, G. C. 2010. Pesquerías globalizadas: revisitando a la comunidad marítima en el Alto Golfo de California. *Estudios Sociales* 18:135–163.

*Associate Editor: Terry Messmer*

**GERARDO RODRÍGUEZ-QUIROZ** earned his B.S. degree in Oceanology from the Universidad



Autonoma de Baja California, Mexico in 1989, his M.S. degree from El Colegio de la Frontera Norte in 1998, and his Ph.D. degree in 2008 from CIBNOR, SC. He has been a full-time professor and researcher at the Instituto Politécnico Nacional since 1999,

ascribed at the CIIDIR-SINALOA multidisciplinary research center. He has been conducting research for the past 18 years with interests in organic aquaculture and the relation between artisanal fishermen and natural protected areas in Mexico. He has directed 8 theses, which include 2 Ph.D. degrees and 6 M.S. degrees. Since 2010, he has been ratified as National Researcher Level I at CONACYT and has been granted the 1998 Baja California Environmental Award.

**WENCESLAO VALENZUELA-QUIÑONEZ**

earned a degree in fisheries biology from Facultad de Ciencias del Mar of the Universidad Autónoma de Sinaloa in 1994 and his doctoral degree in use, management, and conservation of natural resources from CIBNOR, Mexico. He has been conducting research for the past 19 years with interests in the marine ecology of invertebrates, fish, and other groups of exploited and endangered species at the Interdisciplinary Research Center for Regional Integral Development (Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional [CIIDIR-IPN]) of the National Polytechnic Institute of Mexico. He currently works at researching natural resources and conservation and the graduate-level programming for the Department of Aquaculture at CIIDIR-IPN, Sinaloa.



**HÉCTOR A. GONZÁLEZ-OCAMPO**

earned his B.S. degree in marine biology from the Universidad Autónoma de Baja California Sur, Mexico in 1998 and his NRM's from the CIBNOR, SC, BCS, Mexico in 1998 and 2004, respectively. He is a full-time professor in the second largest university in Mexico, the Instituto Politécnico Nacional, since 2006, ascribed at the CIIDIR-SINALOA multidisciplinary research center. He has been conducting research for the past 14 years with interests in the environmental impact of human activities, as well as sustainable nutritional feeds based in ethnobotanical herbs applied in aquaculture. He has patented different biotechnological products based in the microencapsulation of phenolic extracts from plants, natural antibiotics, and microorganisms used as alternative products in the aquaculture and agropecuary sectors.



**ALFREDO ORTEGA-RUBIO** has a Ph.D. degree in ecology and has been granted 3 national



awards by the President of Mexico for his career in scientific research applied to biodiversity conservation in natural protected areas. He has authored 199 original research articles, published 23 books and 111 book chapters, and led 162 strategic projects of resource management, ecological restoration, and environmental impact assessment,

including those required to create the Sierra de La Laguna, the Vizcaíno Desert, and the Revillagigedo Archipelago Biosphere Reserves. He has directed 66 theses, which include 35 Ph.D. degrees and 25 M.S. degrees. Since 2004, he is Researcher "E," the top researcher level at the CONACYT Research Center System, and he has been ratified by the third consecutive time as National Researcher Level III, the top category.