7-22-1913

Australian Bees. i. A New *Crocisa*, with a List of the Australian Species of the Genus

T. D. A. Cockerell
University of Colorado

Follow this and additional works at: https://digitalcommons.usu.edu/bee_lab_co

Part of the [Entomology Commons](#)

Recommended Citation

Cockerell, T. D. A., "Australian Bees. i. A New *Crocisa*, with a List of the Australian Species of the Genus" (1913). Co. Paper 501.

https://digitalcommons.usu.edu/bee_lab_co/501

This Article is brought to you for free and open access by the Bee Lab at DigitalCommons@USU. It has been accepted for inclusion in Co by an authorized administrator of DigitalCommons@USU. For more information, please contact digitalcommons@usu.edu.

2428

old! W. A. Weber

1129

Ts Rebia

[From the Proceedings of the Linnean Society of New South Wales,
1912, Vol. xxxvii., Part 4, October 30th.]

AUSTRALIAN BEES. I. A NEW CROCISA, WITH A
LIST OF THE AUSTRALIAN SPECIES OF THE
GENUS.

By T. D. A. COCKERELL.

(Communicated by W. W. Froggatt, F.L.S.)

[Issued 22nd July, 1913.]

Hymenoptera. Apoidea. Anthophoridae. Melictinae - Thyreus (= Crocisa)

Systematics

[*From the Proceedings of the Linnean Society of New South Wales,*
1912, Vol. xxxvii., Part 4, October 30th.]

AUSTRALIAN BEES. i. A NEW CROCISA, WITH A
LIST OF THE AUSTRALIAN SPECIES OF THE
GENUS.

By T. D. A. COCKERELL.

(*Communicated by W. W. Froggatt, F.L.S.*)

CROCISA WAROONENSIS, sp. nov.

♂. Length 9 mm., expanse a little over 20. Markings (due to hair) chalk-white; face, occiput and greater part of cheeks covered with white hair, lower part of cheeks and underside of head with black hair; labrum gibbous at sides; ocelli in a straight line; vertex shining, very sparsely punctured; flagellum obscurely brownish beneath; mesothorax shining strongly, unevenly, not densely punctured; anterior third, sides narrowly, and posterior corners of mesothorax covered with long loose white hair; scutellum with sparse small punctures, its posterior margin ~-like, the posterior middle occupied by a quadrate patch of white hair, twice as broad as long, white hair also extending from beneath the margin; sides of metathorax with long white hair; pleura with the upper half densely covered with white hair, the under side of thorax with scanty black hair; tegulae black, with very fine punctures, and a patch of white hair in front; wings with the basal half clear hyaline, the apical dark fuscous, clouded or spotted with paler in the region of the cells; third submarginal cell very narrow, but strongly bulging outwardly; anterior and middle tibiae covered with white hair on outer and posterior side except at apex; hind tibiae with the apical half free from white hair except posteriorly; middle and hind basitarsi with a little white hair; hind basitarsi flattened and curved; abdomen rather closely punctured, ^{and marked} as follows: two transversely placed large hour-glass-shaped ones on first segment; four spots or patches each on second and third, the inner round, the outer (especially on second) large patches; two large patches each on fourth and fifth; venter with lateral spots of white hair. There is no basal spot on first abdominal segment.

Hab.—Waroona, Western Australia, April 4, 1908 (G. F. Berthoud). Froggatt, No.210.

This may be compared with *C. quadrimaculata* Rads., which it resembles in the colour of the wings, but *C. quadrimaculata* has the thorax above with nine white spots, and the maculation of the first abdominal segment is different. It is very much smaller than *C. lugubris* Smith.

I give a list of the known Australian species of *Crocisa*.

i. Blue-spotted Species. For a table, see Entomological News, February 1907, p.46.

C. lamprosoma Boisduval. Queensland.

C. turneri Friese. Queensland and New South Wales.

C. quartinae Gribodo. Confused with *C. caeruleifrons* W. F. Kirby, which is quite distinct.

C. darwini Cockerell. Port Darwin. Blue markings, shining.

C. tincta Cockerell. Toowoomba, Queensland. Markings of abdomen pale blue.

C. beatissima Cockerell. Adelaide. Markings of abdomen bright blue, but not shining.

ii. White-spotted Species. For tables, see Bull. Amer. Mus. Nat. Hist., xxiii. (1907), p.232; and Entomologist, August, 1910, p.217.

C. albopicta Cockerell. Mackay, Queensland.

C. waroonensis Cockerell. Western Australia.

C. rotundata Friese (*albomaculata* Smith, preoccupied). Mackay, Queensland.

C. lugubris Smith.

C. macleayi Cockerell. New South Wales.

C. quadrimaculata Radoszkowski. New South Wales.

Excluded Species.

The following two blue-spotted species have been considered to be Australian. I have seen them only from Amboina, and do not believe they occur in Australia.

C. novae-hollandiae Lepeletier. "New Holland."

C. nitidula Fabricius.