

## PREY AVAILABILITY AND SELECTION BY MOUNTAIN LIONS IN THE ARAVAIPA-KLONDYKE AREA OF ARIZONA

STAN C. CUNNINGHAM, Research Branch, Arizona Game and Fish Department, 2221 W. Greenway Road, Phoenix, AZ 85023

ABSTRACT: Prey selection by mountain lions (*Felis concolor*) in the Aravaipa-Klondyke area (2,000 km<sup>2</sup>) in southeastern Arizona was studied from February 1991 through September 1993. Overall diet from frequency of occurrence as determined from 370 scats was: 48% deer (white-tailed and mule deer combined), 34% cattle, 17% javelina, 6% rabbit (cottontail and jackrabbit), 4% rodent, and 2% desert bighorn. Using a correction factor developed by Ackerman et al. (1984), we also estimated percent biomass and proportion of individuals killed. With respect to biomass consumed, cattle was 44%, deer 40%, javelina 10.9%, rabbits 2.9%, and rodents 0.02%. Based on weights of prey consumed, proportion of individuals eaten changed to rabbits 52.7%, deer 16.3%, rodents 12%, javelinas 10% cattle 8%, and desert bighorn 0.5%. Comparisons with availability as determined from four separate four-day double-count helicopter surveys found that lions selected calves (ate more than expected based on availability alone), killed and ate less deer than expected, and javelina as would be expected.

---

Proc. East. Wildl. Damage Mgmt. Conf. 7:130 1997.