



## Asparagus beetle and spotted asparagus beetle

Erin W. Hodgson  
Extension Entomology Specialist

Dan Drost  
Extension Vegetable Specialist

### What You Should Know

- Asparagus beetle and spotted asparagus beetle are found wherever asparagus is grown in North America.
- Adult and larval feeding can reduce photosynthetic capacity by feeding on young spears and ferns, and negatively affect crop yield.
- Removing weeds and plant debris around asparagus will minimize overwintering and refuge during harvest.

**A**sparagus beetle, *Crioceris asparagi*, and spotted asparagus beetle, *C. duodecimpunctata* are leaf beetles in the family Chrysomelidae. These beetles feed exclusively on asparagus and are native to Europe. Asparagus beetle is the more economically injurious of the two species. Since the early 1900's, they can be found wherever asparagus is grown in the United States and Canada. Although the asparagus beetle dominates in other asparagus-growing regions such as California, both species are equally abundant in Utah.

### Damage

Both the adults and larvae of asparagus beetle and spotted asparagus beetle can damage plants. Adults can consume leaves, and scar spears and stems (Fig. 5). The most serious damage caused by either species is during the spring, when adults are feeding on new spears (Fig. 1). These beetles can occasionally damage plants during the summer, but rarely stunt growth or reduce yield. The spotted asparagus beetle prefers to eat asparagus berries and rarely causes damage to leaves and spears. Generally, larval berry feeding is not economically important unless growing asparagus for seed.

Although not directly damaging to the plant, females cement eggs onto spears or ferns (Fig. 2). Eggs on asparagus is usually an aesthetic problem because eggs cannot be washed away or easily removed.



Fig. 1. Asparagus beetle feeding damage.<sup>1</sup>

### Control Options

**Biological control:** In Utah, there are several natural enemies that target asparagus beetle and spotted asparagus beetle larvae. Predatory beetles and parasitoid wasps will help keep populations to tolerable levels in most situations.

**Sanitation:** Removing volunteer asparagus will prevent population spread over time. Minimizing weeds or other debris near asparagus will make sheltered areas less favorable for overwintering adults.

**Insecticidal control:** For asparagus with persistent beetle damage, insecticides can be used control adults and larvae. Consider reduced risk insecticides first because they have a short residual and are not broad spectrum, including neem oil, kaolin clay (Surround®), diatomaceous earth, pyrethrins (Concern®), and spinosad (Entrust®, Success®). Conventional insecticidal dusts include carbaryl and permethrin.

## Asparagus Beetle

**Life cycle:** There are 2-3 generations of asparagus beetle per year in Utah, but the third generation may be small in colder parts of the state. In Utah, the asparagus beetle causes more spear damage compared to the spotted asparagus beetle. Asparagus beetles overwinter as adults in hollow asparagus stems, under weeds or other plant debris, or under loose tree bark. Temperatures determine beetle emergence every year; adults begin feeding on spears as soon as they break through the soil surface in early spring. Adults mate and females lay eggs on asparagus spear tips or ferns. Eggs are glued on their ends (Fig. 2) and hatch about 7-10 days later. Larvae immediately begin feeding on spears and may contaminate them with their dark frass (excrement). Larvae develop through four instars in about 7-10 days before burrowing into the soil to pupate. Adults emerge from the soil in about 5 days to begin the next generation.

**Description:** Asparagus beetle eggs are brown, oval and 1 mm long, and are individually cemented on asparagus plants. Larvae are plump and brown with a dark head capsule and chewing mouthparts (Fig. 3). Larvae have three pair of legs on the thorax (region behind the head) and fleshy prolegs near their hind end for grasping plants while feeding. Fully grown larvae reach up to 8 mm in length. The pupae resemble the adults except for the poorly developed wings and yellow color. The adults are brightly colored leaf beetles, reaching approximately 5-6 mm in length (Fig. 4). The head, antennae, and legs are bluish black in color. The forewings can be variable in color, but in general, each wing is bluish black with three white spots and orange wing margins. Adults are reluctant to fly away when disturbed, but will drop from plants and “play dead” to avoid predation.



Fig. 2. Asparagus beetle eggs on ferns.<sup>2</sup>



Fig. 3. Asparagus beetle larvae feeding.<sup>1</sup>



Fig. 4. Asparagus beetle.<sup>3</sup>



Fig. 5. Damage asparagus.<sup>1</sup>

## Spotted Asparagus Beetle

**Life cycle:** There are 2 generations of spotted asparagus beetle per year in Utah. Spotted asparagus beetles overwinter as adults in hollow asparagus stems and other sheltered areas. Depending on spring temperatures, adults begin feeding on spears when they break through the soil surface (Fig. 6). Adults mate and females lay eggs on spears. Spotted asparagus beetle eggs can be distinguished from asparagus beetle eggs because they are cemented on their side rather than on their end. In about 7-10 days eggs hatch into larvae. Larval populations build up in late June after the ferns are established, and prefer to feed on asparagus berries when available. Larvae go through four instars in 21-30 days before pupating within the berry. Adults emerge from berries in 12-16 days to begin the next generation. Sometimes females will delay oviposition until flowers are fertilized or berries are present.

**Description:** Spotted asparagus beetle eggs are brown or olive green, oval and 1 mm long. The eggs are individually cemented on asparagus plants. Larvae are plump, and pale yellow or orange in color with chewing mouthparts. Larvae have three pair of legs on the thorax and anal prolegs for grasping onto plants while feeding. Fully grown larvae reach up to 8.5 mm in length. The pupa resembles the adult except for the poorly developed wings and yellow color. The adults are brightly colored leaf beetles, reaching approximately 6-8 mm in length (Fig. 7). The spotted asparagus beetle adult is mostly orange, but has black spots on the forewings and black legs, eyes, antennae and "knee caps" (Fig. 7). Adults will quickly fly away when disturbed and stridulate (make a clicking sounds) loudly if captured.



Fig. 6. Spotted asparagus beetle feeding on new asparagus spears.<sup>1</sup>



Fig. 7. Spotted asparagus beetle.<sup>1</sup>

<sup>1</sup> Images courtesy of Whitney Cranshaw, Colorado State University Extension ([www.ipmimages.org](http://www.ipmimages.org)).

<sup>2</sup> Image courtesy of Debbie Roos, North Carolina Cooperative Extension (<http://chatham.ces.ncsu.edu/growingsmallfarms/>).

<sup>3</sup> Image courtesy of Clemson University, USDA Cooperative Extension Slide Series ([www.ipmimages.org](http://www.ipmimages.org)).

**Precautionary Statement:** All pesticides have benefits and risks, however following the label will maximize the benefits and reduce risks. Pay attention to the directions for use and follow precautionary statements. Pesticide labels are considered legal documents containing instructions and limitations. Inconsistent use of the product or disregarding the label is a violation of both federal and state laws. The pesticide applicator is legally responsible for proper use.

Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran's status. USU's policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions. USU employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran's status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USU-sponsored events and activities. This publication is issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Noelle Cockett, Vice President for Extension and Agriculture, Utah State University.