



Using Degree Days to Time Treatments for Insect Pests

Marion S. Murray, IPM Project Leader


Insecticides that are applied for a perennial insect pest based on a calendar date often result in poor insect control and a waste of resources. Insect activity varies from year to year depending on weather. For example, in Logan, Utah, eggs of the apple pest codling moth began hatching on May 15 in 2005, May 5 in 2006, and April 30 in 2007. If apple growers always spray on May 1, they are not making the most effective insecticide treatment in most years. As long as accurate weather data can be obtained, using degree days to time treatments is more reliable than a calendar date and allows growers to pinpoint a specific treatment date each year.

Degree days (often referred to as "growing degree days") are accurate because insects have a predictable development pattern based on heat accumulation. Insects are exothermic ("cold-blooded") and their body temperature and growth are affected by their surrounding temperature. Every insect requires a consistent amount of heat accumulation to reach certain life stages, such as egg hatch or adult flight. Degree day values interpret that heat accumulation. When used to determine treatment timing, they are an important component of an Integrated Pest Management program, providing a cost effective tool to reduce insect feeding damage.

CALCULATING DEGREE DAYS

Simply put, a degree day (DD) is a measurement of heat units over time, calculated from daily maximum and minimum temperatures. Degree days are based on the rate of an insect's development at temperatures between upper and lower limits for development (see Figure 1). The minimum temperature at which insects first start to develop is called the "lower developmental threshold", or baseline. The maximum temperature at which insects stop developing is called the "upper developmental threshold," or cutoff. The lower and upper thresholds vary among species, and have been determined for many, but not all, major insect pests. For those whose exact values are unknown, including most landscape insect pests, a baseline temperature of 50°F is used. Some insects do not have an upper development threshold.

Figure 1. An insect's development follows a predictable progression based on temperature. When insects reach their upper threshold, development of some species levels off (horizontal cutoff), and for other species, stops (vertical cutoff).



Although degree days are usually calculated for a 24-hour time period, it is the number of accumulated degree days from a starting point, called a biofix, that is most useful. The biofix can be a biological event, such as the date at which moth flight begins, or a calendar date, such as March 1. In northern Utah, we start accumulating degree days for insect pests, such as codling moth, that have a baseline of 50°F on March 1, because there is typically no insect development before that time.

No matter how it is calculated, the degree day value for a 24-hour period is added to the prior day's values, and so on. For an average growing season in northern Utah, areas will accumulate approximately 2500-3500 degree days (with a baseline of 50°F).

Average Method

In general, degree days can be calculated using a simple formula for the average daily temperature, calculat-

ed from the daily maximum and minimum temperatures, minus the baseline (lower developmental threshold):

$$[(\text{daily maximum temperature} + \text{daily minimum temperature})/2] - \text{baseline temperature}$$

For example, a day where the high is 72°F and the low is 44°F would accumulate 8 degree days using 50°F as the baseline:

Example 1: $[(72 + 44)/2] - 50 = 8$

When temperatures do not exceed 50, zero degree days have accumulated. This calculation method is the simplest and least precise.

Modified Average Method

The problem with the average method is that it does not take into account the length of time that the daily temperature may exceed the baseline temperature. In Example 1, results could be skewed if the minimum temperature of 44° F occurred for only 30 minutes out of the 24-hour day while the rest of the time the temperature was above 50° F. Given this, the accumulated degree days using the above calculation would be less than the actual value. To account for situations when the daily minimum temperature is less than the baseline, or the daily maximum temperature is greater than the cutoff, the formula needs to be modified. When either occurs, the lower threshold is used instead of the daily minimum, or the upper threshold is used instead of the maximum. For the above example, we would use 50°F as the daily minimum temperature in the formula instead of 44°F:

Example 2: $[(72 + 50)/2] - 50 = 11$

And for a day with a maximum temperature of 102°F, and a low of 70°F, we would replace the 102 with the upper threshold temperature, if known. We know that it is 88°F for codling moth, so the degree days would be:


Example 3: $[(88+70)/2] - 50 = 29$

Sine Wave Method

A more precise and method of calculating degree days is called the sine wave method. This method still uses the daily minimum, maximum, and baseline temperatures (lower threshold), but also incorporates the upper threshold temperature into the calculation. It is based on the assumption that temperatures of a 24-hour day follow a sine wave curve. The number of degree days is then calculated as the area under this curve within the lower and upper temperature thresholds (see Figure 2).

Because of the somewhat complicated calculus involved, the sine wave formula is not shown here. Degree days calculated using this method are usually determined by a computer.

Figure 2. This diagram is a visual representation of degree days using the sine wave method of calculation, with a horizontal cutoff. The area in black under the curve represents the number of degree days that fall between a lower and upper threshold, for each 24-hour period.



USING DEGREE DAYS

Scouting

Accumulated degree days are useful in timing scouting events such as when to place traps, when to look for damage, when to sample, etc. As an example, codling moth pheromone monitoring traps are placed in the apple orchard at 100 degree days after March 1 in northern Utah to determine initiation of adult moth flight.

Using Insect Models

Scientists have studied biological development over time (phenology) of insects in correlation to accumulated degree days, discovering information on key physiological events, such as egg hatch, adult flight, etc. This predictive information is known as an insect model. Insect models are useful in timing insecticide treatment because the entire life cycle (or certain important events) of the insect is known. Models have been developed for a number of insect pests (see Table 1).

Predicting Treatment Timing

With the development of more targeted, reduced risk insecticides, timing of application is becoming more and more important. Certain life stages of insects are more susceptible to insecticide treatment such as young larvae or scale crawlers. Degree days are used to predict when those life stages will occur (see Table 2). Degree days are “projected” into the future for a given site using either forecasted daily highs and lows or 30-year average highs and lows. This information is only an approximation of a future event, but is highly useful in planning.

Table 1. A partial list of insect pests that occur in Utah for which we have temperature thresholds and degree day models. Those with an asterisk have been validated for Utah.

Target Insect		Lower Developmental Threshold (F)	Upper Developmental Threshold (F)	Availability of Model
Common Name	Scientific Name			
Alfalfa weevil	<i>Hypera postica</i>	50	87	yes
Armyworm	<i>Pseudaletia unipuncta</i>	50	84	yes
Black cutworm	<i>Agrotis ipsilon</i>	50	86	yes
Cabbage maggot	<i>Delia radicum</i>	40	86	yes
Codling moth*	<i>Cydia pomonella</i>	50	88	yes
Corn earworm*	<i>Helicoverpa zea</i>	55	92	yes
European pine shoot moth	<i>Rhyacionia bouliana</i>	28	---	yes
European red mite	<i>Panonychus ulmi</i>	51	---	yes
Greater peachtree borer	<i>Synanthedon exitiosa</i>	50	87	no
Lilac/ash borer*	<i>Podosesia syringae</i>	50	---	yes
Obliquebanded leafroller*	<i>Choristoneura rosaceana</i>	43	85	yes
Peach twig borer*	<i>Anarsia lineatella</i>	50	88	yes
Pear psylla	<i>Cacopsylla pyricola</i>	41	-	no
San Jose scale*	<i>Quadraspidiotus perniciosus</i>	51	90	yes
Strawberry root weevil	<i>Otiorhynchus ovatus</i>	40	103	yes
Variiegated cutworm	<i>Peridroma saucia</i>	45	80	yes
Walnut husk fly*	<i>Rhagoletis completa</i>	41	130	yes
Western cherry fruit fly*	<i>Rhagoletis indifferens</i>	41	130	yes

*Insect model has been validated for Utah

Threshold and model information from: UC-Davis IPM Web site: <http://www.ipm.ucdavis.edu/MODELS/index.html>

Table 2. A partial list of degree day (GDD) accumulations for selected landscape pests that occur in Utah. "DD Min" is the earliest timing for appearance, and "DD Max" is the latest timing.

Common Name	Scientific Name	Life Stage*	DD Min	DD Max
Black pineleaf scale	<i>Dynaspidiotus californica</i>	E	1068	
Bronze birch borer	<i>Agrilus anxius</i>	A	440	800
Cankerworms	<i>Alsophila sp.</i>	L	148	290
European fruit lecanium scale	<i>Parthenolecanium corni</i>	C	800	
European pine shoot moth	<i>Rhyacionia bouliana</i>	L A E	50 700 900	220 800 1000
Honeylocust plant bug	<i>Diaphnocoris chlorionis</i>	N, A	58	246
Lilac/Ash borer	<i>Podosesia syringae</i>	L	148	299
Lilac root weevil	<i>Otiorhynchus meridionalis</i>	A	500	950
Locust borer	<i>Magacyllene robiniae</i>	L, A	2271	2805
Oystershell scale	<i>Lepidosaphes ulmi</i>	C C	363 1600	707 1700
Pine needle scale	<i>Chionaspis pinifoliae</i>	C C	298 1388	448 1917
Spruce spider mite	<i>Oligonychus ununguis</i>	E,L E,L,A E,L,N,A	7 192 2375	121 363 2806
Western tent caterpillar	<i>Malacosoma californicum</i>	L	100	500
Western spruce budworm	<i>Choristoneura occidentalis</i>	L	200	300

*E (eggs), N (nymph), C (crawler), L (larvae), A (adults)

Degree day values determined by: Dr. Warren T. Johnson, Department of Entomology, Cornell University.

The most widely used insect model in Utah is for codling moth (see Table 3). For this pest, it is important to know when 220 degree days after biofix will occur, because this point corresponds to first generation egg hatch, when fruit should begin to be protected.

Table 3. Example of an insect model for **codling moth** showing method of calculation and degree days required for development.

Developmental Thresholds	
Lower: 50 F	
Upper: 88 F	
Calculation Method: Single Sine	
Cutoff Method: Horizontal	
Set out Traps: 100 DD after March 1	
Biofix: First consistent (2+ in a single trap) catch of adults in the pheromone trap	
Degree-Day Accumulations Required for Each Stage of Development	
Event	DD
Generation Time (egg to egg)	880
Generation Time (50% egg hatch to same)	1096
1% egg hatch (1st gen)	220
20% egg hatch (1st gen)	360
50% egg hatch (1st gen)	484
75% egg hatch (1st gen)	610
95% egg hatch (1st gen)	800
5% Adult emergence (2nd gen)	1000
7% egg hatch (2nd gen)	1260
30% egg hatch (2nd gen)	1460
50% egg hatch (2nd gen)	1580
75% egg hatch (2nd gen)	1750
95% egg hatch (2nd gen)	2000

Some digital models will store up to seven days of readings. The thermometer should be calibrated at the start of each season, and placed away from direct sunlight, ideally in a white shelter box. Obtain degree days in one of the following ways:

- Calculate them daily using the average or modified average method.
- Use a degree day look-up table. Degree day values for high and low temperatures are available for certain insects in a look-up table (see Table 4 for example).
- Enter daily maximum and minimum temperatures into a computer spreadsheet that is set up to calculate the values.

2. Biophenometers are instruments that calculate degree days every few minutes based on temperatures and are highly accurate. Many brands allow you to manually input the target pest's upper and lower thresholds. They can be purchased as a stand-alone, or in conjunction with a weather station. Minor setbacks include price (\$300-\$1000), and the fact that the instruments' degree day calculation method provides different results than the modified sine wave method. Typically, the degree days that researchers determined for insect models were calculated using the sine wave method, so values calculated from the biophenometers would be slightly less. They may need to be compared to the values from the sine calculation for one season and readjusted accordingly.

3. USU Extension weekly pest reports (<http://utahpests.usu.edu/ipm/htm/advisories>) provide accumulated and predicted degree days for a variety of sites across northern Utah. Your local county Extension office can also help you with this information.

4. An Internet search for "degree day calculator" can often turn up sites where you can enter your own data, or select a location.

OBTAINING DEGREE DAYS

Whether you are calculating your own degree days, or using information from an instrument or Web site, it is important to know how the degree days were calculated for the target insect, and that your calculation method matches, or is modified to match. There are a variety of ways to acquire degree days:

1. To calculate your own degree days, you will need a thermometer that records maximum and minimum temperatures in a location that closely matches the temperatures that your target pest(s) would encounter. Max-Min thermometers are inexpensive, easily available, and record in digital or mercury.

LIMITATIONS OF DEGREE DAYS

The primary limiting factor in using degree days is obtaining accurate temperature readings. If a thermometer, biophenometer, or weather station location is not representative of the environment in which the target insect occurs, the resultant degree days will not mirror the actual insect development. In addition, temperatures at one site may not be reflective of conditions in another site several miles away. This is particularly true of Utah, where mountains, lakes, and deserts result in a wide variety of microclimates.

Table 4. Example of a degree day look-up table for peach twig borer and codling moth (base 50°F) (not a complete table).

		Minimum Temperature (°F)											
		48	51	54	57	60	63	66	69	72	75	78	81
Maximum Temperature (°F)	52	1	2										
	58	2	3	5									
	61	5	6	8	9	11							
	64	6	8	9	11	12	14						
	70	9	11	12	14	15	17	18	20				
	73	11	12	14	15	17	18	20	21	23			
	76	12	14	15	17	18	20	21	23	24	26		
	79	14	15	17	18	20	21	23	24	26	27	29	
	82	15	17	18	20	21	23	24	26	27	29	30	32
	85	17	18	20	21	23	24	26	27	29	30	32	33
	88	18	20	21	23	24	26	27	29	30	32	33	35
	91	19	21	22	24	25	27	28	30	31	32	34	35

GLOSSARY

Baseline: Equivalent to “lower developmental threshold.”

Biofix: A date that signals the start of growing degree day accumulations (“biological fix”). The date can be represented by a biological event, such as first moth flight, or a calendar date.

Degree days: A measurement of heat units over time, equivalent to the number of degrees that the average temperature is above a baseline value. Also known as “growing degree days” (GDD) to differentiate this value from “heating degree days” or “cooling degree days,” which are used to estimate energy demand.

GDD (50), DD (50), GDD₅₀, DD₅₀, DD (base 50), etc: Terminology used to describe a value of degree days,

using 50° F as the baseline temperature.

Lower Developmental Threshold: A temperature at which insect development begins (also known as “baseline”); determined by laboratory studies.

Phenology: The study of periodic biological events, such as plant flowering, insect development, etc., in relation to environmental factors, such as temperature (translated as: “knowledge of phenomena”).

Upper Developmental Threshold: A maximum temperature of insect development where development levels off or slows down; determined by laboratory studies.

REFERENCES

Agnello, A. M., et al. 1993. Fruit Pest Events and Phenological Development According to Accumulated Heat Units. New York State Agriculture Experiment Station.

Beers, Elizabeth, et al. 1993. “Degree Day Models” in: Orchard Pest Management. Good Fruit Grower. 45-48.

Herns, Daniel A. 2004. “Using Degree Days and Plant Phenology to Predict Pest Activity” in: IPM of Midwest Landscapes. MN Agriculture Experiment Station. 49-59.

Kowalsick, Thomas, and Scott Clark. 2006. Using Growing Degree Days for Insect Pest Management. Cornell Cooperative Extension. 2006.

Precautionary Statement: All pesticides have benefits and risks, however following the label will maximize the benefits and reduce risks. Pay attention to the directions for use and follow precautionary statements. Pesticide labels are considered legal documents containing instructions and limitations. Inconsistent use of the product or disregarding the label is a violation of both federal and state laws. The pesticide applicator is legally responsible for proper use.

In its programs and activities, Utah State University does not discriminate based on race, color, religion, sex, national origin, age, genetic information, sexual orientation or gender identity/expression, disability, status as a protected veteran, or any other status protected by University policy or local, state, or federal law. The following individuals have been designated to handle inquiries regarding non-discrimination policies: Executive Director of the Office of Equity, Alison Adams-Perlac, alison.adams-perlac@usu.edu, Title IX Coordinator, Hilary Renshaw, hilary.renshaw@usu.edu, Old Main Rm. 161, 435-797-1266. For further information on notice of non-discrimination: U.S. Department of Education, Office for Civil Rights, 303-844-5695, OCR.Denver@ed.gov. Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Kenneth L. White, Vice President for Extension and Agriculture, Utah State University.

Fact Sheets: [IPM-General]