

Gunnison Sage-grouse

Centrocercus minimus

Conservation Plan Update

San Juan County, Utah

February 2003

## PREAMBLE

The Gunnison Sage-grouse (*Centrocercus minimus*) inhabits a limited area in Colorado and Utah. San Juan County is the only place in Utah where Gunnison Sage-grouse populations are currently known to occur. On January 26, 2000 the U.S. Fish and Wildlife Service received a petition signed by representatives of the American Lands Alliance, Net Work Associates, The Larch Company, Biodiversity Legal Fund, Wild Utah Forest Campaign, and Sinapu requesting an emergency listing for the Gunnison Sage-grouse. The petitioners cited increasing habitat fragmentation, reduced and limited population distributions, and low and declining localized populations as the primary reasons for classifying the species as endangered. Prior to receiving the petition, the U.S. Fish and Wildlife Service had designated the species as a candidate for listing.

In 1972, a total of 175 males were counted strutting on leks in San Juan County. By 1999, this had dropped to 43. In 1972, the sage-grouse population in San Juan County was estimated to be between 583 and 1,050 birds. In 2002, 35 males were counted on four leks. Based on this count, the current population is estimated to be between 178 and 308 birds.

The San Juan County Gunnison Sage-grouse Working Group (SWOG) was formed in 1996 for the purpose of developing a conservation plan that could be implemented by state and federal wildlife resource agencies, private landowners, and local governments to benefit sage-grouse populations in the county. The conservation plan was finalized and signed in November 2000.

SWOG believes that implementation of the conservation plan will ensure local ownership in future management and land-use decisions; respect private property rights; and embrace community economic, cultural, and social values. This update identifies conservation strategies that have been accomplished since SWOG was organized and the plan was signed. SWOG believes the actions implemented by private and public partners in San Juan County will help to restore Gunnison Sage-grouse habitats and populations while benefiting the local economy and community.

## Table of Contents

**Page**

PREAMBLE .....	ii
TABLE OF CONTENTS.....	iii
LIST OF TABLES AND FIGURES.....	iv
I. PLAN BACKGROUND.....	1
II. PLAN DEVELOPMENT- HABITAT CONSERVATION ASSESSMENT .....	4
III. PLAN IMPLEMENTATION - CONSERVATION STRATEGIES .....	17
IV. REFERENCES .....	31
V. APPENDICES .....	32
A. San Juan County Gunnison Sage-grouse Working Group Members.....	32
B. Vegetation Classifications and Definitions.....	33
C. Habitat Mixture Seeded on CRP Land in San Juan County Gunnison Sage-grouse Conservation Plan Area. ....	36
D. Gunnison Sage-grouse Conservation Brochure.....	37
E. Summary of Gunnison Sage-grouse Conservation Projects and Costs .....	38

## List of Tables and Figures

### Figures

Figure 1. Gunnison Sage-grouse Conservation Area. San Juan County, Utah.....	2
Figure 2. Gunnison Sage-grouse Conservation and Study Area Vegetation Map, San Juan County, Utah.....	6
Figure 3. Gunnison Sage-grouse spring and summer habitat use patterns and seasonal movements, San Juan County, Utah .....	7
Figure 4. Gunnison Sage-grouse Working Group conservation projects, San Juan County, Utah 2002 .....	11

### Tables

Table 1. Landownership in the Gunnison Sage-grouse Conservation Area .....	3
Table 2. Number of males counted on leks in San Juan County, 1972-2002 .....	5
Table 3. Vegetation conditions within the San Juan County Gunnison Sage-grouse Conservation Plan Study Area, 2002 .....	8
Table 4. Percent cover of grasses, forbs, and shrubs at nest and reference sites for 2001 and 2002.....	9
Table 5. Percent availability and use of cover types by sage-grouse hens with broods in 2001 and 2002 in San Juan County, UT .....	12
Table 6. Vegetation height (cm) and percent cover of grasses, forbs, and shrubs at brood-use sites in 2001 and 2002 .....	13
Table 7. Vegetation height (cm) and percent cover of grasses, forbs, and shrubs at brood-use sites and male Gunnison Sage-grouse locations in San Juan County, UT, 2001 .....	14
Table 8. Simultaneous confidence intervals using the Bonferroni approach for utilization of vegetation types by male Gunnison Sage-grouse in 2001 and 2002.....	14
Table 9. Percent cover of grasses, forbs, and shrubs at reference sites	

and male Gunnison Sage-grouse locations in San Juan County,  
UT, 2001 ..... 14

Table 10. Percent cover of grasses, forbs, and shrubs used by broods,  
broodless hens, and male Gunnison Sage-grouse in San Juan  
County, UT, 2002 ..... 15

Table 11. Simultaneous confidence intervals using the Bonferroni approach  
for utilization of vegetation types by non-brood rearing Gunnison  
Sage-grouse in 2002..... 15

## I. PLAN BACKGROUND

### A. Introduction and Purpose

The San Juan County Gunnison Sage-grouse Working Group (SWOG) was formed in 1996 to identify and implement conservation strategies to reverse the decline of sage-grouse populations in the county. From the onset, SWOG has sought local citizen involvement and input. SWOG consists of public management agencies, private landowners, local citizens, and private conservation groups. A current list of SWOG members can be found in Appendix A.

### B. Conservation Area Description and Boundary

San Juan County is located in southeastern Utah and consists of 7,821 square miles (Figure 1). Approximately 6% of the county (324,921 acres) consists of 208 privately-owned farms. In 1994, agriculture enterprises generated slightly over 10% of the county's total personal income. San Juan County ranks 5<sup>th</sup> among all Utah counties in total grain production.

The San Juan County Gunnison Sage-grouse Conservation Area (CA) boundary was delineated using current and historic habitats, sage-grouse observations, and an assessment of the potential for remaining sagebrush areas in the county to provide suitable habitat (Figure 1). The CA encompasses rural areas, rural residential dwellings, and agricultural croplands (Table 1). The CA area consists of 970,660 acres. Thirty-eight percent (314,244 acres) is privately-owned. Within the CA, SWOG has delineated a core conservation area (CCA) consisting of 336,680 of which 89% (218,493 acres) is privately-owned. Within the CCA, SWOG has identified a conservation study area (CSA). The CSA consists of 59,744 acres. Over 93% (55,738 acres) of the SA is privately-owned. The area within the CA, but located outside of the CCA have been identified as a buffer area. While SWOG believes it necessary to include the buffer area in the CA because of the presence of sagebrush, we make no inferences regarding habitat potentials. The buffer area does not presently support Gunnison Sage-grouse.

Table 1. Landownership in the Gunnison Sage-grouse Conservation Area.

Area	Total Acres	Public Acres				Private Acres
		Forest Service	BLM	State	Tribe	
CA (38%)	970,066	195	457,333	65,271	10,862	314,224
CCA (89%)	336,680	-	24,835	3,483	-	218,493
SA	59,744	-	3,024	919	-	55,738 (93%)

CA – Conservation Area, CCA – Core Conservation Area, CSA – Conservation Study Area

Figure I.

## **C. The Process**

Gunnison Sage-grouse in the county depend heavily on private lands for habitat (Table 1). Thus, SWOG is committed to conserving and enhancing Gunnison Sage-grouse populations that occur on privately-owned land in the county and contribute to the economic viability of farms, ranches, and the local community. Landowner involvement in the Plan is strictly voluntary.

The SWOG Conservation Plan (PLAN) consists of two parts. The first part is the Population and Habitat Conservation Assessment. The Assessment describes SWOG's current understanding about the status of Gunnison Sage-grouse distributions, habitat conditions, and factors that may be affecting populations in the county. Research is ongoing to refine this information. The second part of the PLAN is the Conservation Strategy. The Strategy identifies PLAN goals and objectives, conservation actions, implementation schedules and responsibilities, evaluation guidelines, and monitoring requirements.

The PLAN identifies conservation strategies to be implemented in guiding and coordinating management efforts across jurisdictional/land ownership boundaries to improve Gunnison Sage-grouse habitat conditions in the county. SWOG intended that the PLAN be a dynamic document. As new information is obtained or issues identified, SWOG updates the PLAN and modifies conservation strategies accordingly. This document summarizes the progress made to implement the PLAN.

## **II. PLAN DEVELOPMENT - HABITAT CONSERVATION ASSESSMENT**

**A. Current Population Status** In 1972, 175 males were counted strutting on six leks in San Juan County. In 2001 and 2002, 47 and 35 males, respectively, were counted on five and four leks, respectively (Table 2).

### **B. Population Objectives**

#### **1. Spring breeding populations**

One objective of the PLAN is to reestablish a minimum estimated spring breeding population of 500 birds with 6-8 active leks each containing a three-year count average of 20-25 birds per lek. This increase would be measured from 1997 population estimates.

#### *Current Status*

In 1997 the population, based on lek surveys, was estimated to be between 125-140 birds. In 2002, the population was estimated to be between 175-308 birds based on counts of four leks. Only the Roring lek, achieved the 20-25 bird average. The increased number of males counted on this lek may have been the result of birds abandoning the East Seep Wash lek.



The current population estimate is 192-225 birds below the minimum desired population objective. The intent of the PLAN is to achieve this population goal by 2012.

## **2. Lek protection**

All current identified lek sites would be protected from future risk through leases, conservation easements, or in fee title.

### *Current status*

Currently, two of the leks are protected. These include the Bureau of Land Management (BLM) lek (owned in fee title by the BLM) and Hickman Flat lek (part of a perpetual conservation easement on the Adam's property). The Utah Division of Wildlife Resources (UDWR) and the BLM are pursuing conservation easements on the other leks. Currently SWOG is negotiating to secure an easement on the Roring lek and the surrounding areas. Discussions also have been initiated with the East Seep lek landowner (Figure 2)

## **C. Habitat Objectives**

In 2000 and 2001, 14 Gunnison Sage-grouse were captured in the CSA and fitted with radio-collars (8 males and 6 females). We monitored the movements of the birds during the spring and summer. Based on their movements, we have defined the boundaries of the original core conservation study area (CCA) as presented in the PLAN. This area is now referred to as the conservation study area (CSA). The CSA consists of 59,744 acres (Figure 3). This area contains the primary breeding complexes. We are currently monitoring radio-collared birds to determine seasonal habitat use areas.

### **1. Breeding complexes**

An objective of the PLAN is to reestablish desired vegetation conditions on 50-75% of the areas located within two miles of lek sites (Figures 1 and 2). SWOG has defined desired vegetation conditions to include a canopy cover of 20-40% big sagebrush with an average height of 16 inches, 30% minimum grass canopy cover, and 10% minimum forb canopy cover.

### *Current Status*


We have evaluated the existing vegetation conditions in the CSA. This evaluation was conducted by ground truthing Landsat imagery of the CSA. To ground truth the imagery we randomly established 50 training sites within the CSA (Figure 2). These sites were visited and the vegetation conditions documented. Definitions for the vegetation classifications can be found in Appendix B. After completing this fieldwork, we conducted a supervised classification of SA imagery. The results of this are presented in Table 3.

Table 2. Number of strutting males counted on leks in San Juan County, 1972-2002.

YEAR	STRUTTING		GROUNDS		
	RORING	EAST SEEP	HICKMAN FLAT	BLM	DODGE POINT
1970	49	43	9		
1971	51	61	2		
1972	59	64	6		
1973	48	31	7		
1974	41	73	4		
1975	27	51	21		
1976	24	32	33		
1977	18	40	50		
1978	13	30	45		
1979	5	17	39		
1980	4	9	28		
1981		21	39		
1982	2	18	27		
1983	9	15	35		
1984	10	13	28		
1985	7	9	16		
1986	9	6	3		
1987	10	8	3		
1988	11	6	4		
1989	16	11	3		
1990	15	9	4		
1991	11	8	5		
1992	16	14	6		
1993	17	18	3		
1994	18	17			
1995	16	14	8		
1996	14	14			
1997	13	6	6		
1998	15	4	9	4	
1999	22	1	9	6	5
2000	33		12	4	8
2001	25	3	12	4	3
2002	23	0	8	3	1

Lek Counts 1970-2002

Figure 2



**Figure 3**

Table 3. Vegetation classifications of Gunnison Sage-grouse breeding complex , San Juan County, UT, 2002.

Vegetation Classification	Acreage	% Total*	Objective
Non-irrigated Agriculture	4688	17.7	
Black-sage	4287	16.2	
Big Sagebrush > 25 % Canopy	1523	5.7	> 20-40%
Big Sagebrush < 25% Canopy	1706	6.4	
Grass (Conservation Reserve Program)	13,533	51.1	> 30%
Rangeland (Grass)	2908	10.9	

\*These totals are based on 100% of the area. The PLAN calls for achieving the objectives on 50-75% of the area.

Currently the acreages of cover types available for Gunnison Sage-grouse in the area defined as constituting the breeding complex exceed the PLAN objectives, except for the forb canopy cover (see Table 4 reference sites). The suitability of these cover types as nesting habitat is still under study.

To assess Gunnison Sage-grouse habitat use patterns in the CSA and breeding complex, we captured 14 Gunnison Sage-grouse on or adjacent to two separate strutting grounds during March and April each year (6 in 2001, 8 in 2002). In 2001, all birds (1 hen, 5 males) were adults. In 2002 all the males (3/3) were adults and 60 % of the hens were juveniles (3/5).

Four radio-collared females nested (1 in 2002, 3 in 2002). All of the nests were monitored and each hen successfully hatched a brood. All nests hatched between 21 May and 23 May. Allowing for 26 days of incubation, nests were initiated April 25-27.

The only nest monitored in 2001 was located one mile from the nearest lek site. In 2002, nest sites were one-half mile and one and one-half miles from the nearest active strutting grounds. In 2001, the hen nested in black sage (*Artemisia nova*). In 2002, all the hens nested in agricultural land enrolled in the Conservation Reserve Program (CRP). These lands have been reseeded to a grass using a seed mixture provided by the UDWR.

Vegetation measured at nest sites was compared to non-use sites in black sage and CRP/grassland cover classes. The vegetation measured at nest sites differed from non-use sites ( $P = 0.01$ ). Nest sites had fewer grasses and forbs than non-use sites, but had more shrubs

(Table 4). All nests were located under sagebrush canopies. The average height of sagebrush around the nest sites was 8.5 inches (21.5 cm).

Table 4. Percent cover of grasses, forbs, and shrubs at Gunnison Sage-grouse nest and reference sites for 2001-02, San Juan County, UT.

	Percent Cover	
	Nesting	Reference
Grasses	6.0	34.7*
Forbs	0.5	8.8*
Shrubs	27.5	10.4*

\* P-value  $\leq 0.05$  is significantly different

Based on this preliminary analysis, the breeding complex habitat in the CSA meets the minimum desired percent cover requirements established in the PLAN for grass and shrub cover, but is below the desired forb canopy cover objective.

In the late 1980s many landowners within the CCA enrolled their lands in CRP. Most of the decline in non-irrigated agricultural land can be explained by CRP. In 1993, over 43,000 non-irrigated croplands were converted to CRP grasslands. Also, during this period, an additional 10,000 acres of cropland was converted to rangeland.

Many of CRP contracts in the county expired in 1995. Based on new national CRP eligibility requirements, many of these lands and other agricultural lands located in the county would not have been eligible for enrollment in the program. SWOG worked with the Natural Resource Conservation Service (NRCS) and Farm Services Agency (FSA) to have San Juan County designated as a priority conservation area under CRP because of Gunnison Sage-grouse. Designation as a priority conservation area meant that agricultural land submitted for CRP enrollment consideration in the county did not have to meet the CRP erodibility index requirements to be eligible for the program. However, landowners could only qualify for the program if they agreed to implement approved wildlife conservation practices.

As of February 2000, a total of 36,825 acres of private land within the CA had been enrolled in CRP. Approximately 32,667 acres were enrolled as a result of the Gunnison Sage-grouse conservation priority initiative. The UDWR, in conjunction with NRCS, developed a sage-grouse seed mixture for use in San Juan County (Appendix B). The total cost of reseeding these areas was \$531,686. The UDWR and private landowners each paid \$132,921 of this amount. Farm Services Agency (FSA) cost-shared for the remaining \$265,844. The total cost of establishing CRP in San Juan County was \$1,222,728. This includes the seed cost and \$691,042

that was spent to prepare the land for seeding. Half of the costs of land preparation was paid for by the landowners (\$345,521) and half was cost-shared by FSA. CRP leases generate in excess of \$1,000,000 in annual income for participating landowners.

The CRP program enabled SWOG to achieve grass and minimum shrub canopy cover objectives. We believe the forb cover has been affected by the severe drought condition the CA experienced in 2002 and grass competition. Some of the CRP lands in the CSA were grazed in 2002 under an emergency grazing declaration. NRCS considered the grazing to be a stand maintenance treatment. SWOG is monitoring these areas to see what effect the grazing had on canopy cover.

SWOG also has initiated efforts to both hand plant and aerial seed CRP fields with sagebrush in areas that exhibit high bird use to augment existing sagebrush stands thus enhance canopy cover (Figure 4). These efforts are discussed in greater detail under Conservation Strategies.

## **2. Brood-rearing complexes**

The PLAN identifies the need to reestablish brooding rearing-complexes in the CSA. This will be done by enhancing vegetation conditions on 50-75% of the area located within 4 miles of known lek sites within the CCA over the next 10-year period. This area consists of 80,551 acres. Desired conditions include a canopy cover of 20-40% big sagebrush with an average height of 16 inches, 30% minimum grass canopy cover, and 10% minimum forb canopy cover. The height of the vegetation in wet meadow areas is to be greater than 4 inches between June 15- July 31 on over 75% of the area considered to be brood-rearing habitat.

### ***Current Status***

Based on analysis of LandSat imagery, the CCA meets the desired objective percent cover for grass (Table 5). In 2002, we measured vegetation composition in the brood-rearing area. Based on this effort we determined that the percent shrub, forb, and grass composition in the brooding-rearing area to be 11, 9, and 31%, respectively. These data support the LandSat analysis. The 9% forb approximates the 10% PLAN objective.

To determine brood habitat use patterns in the SA, we monitored the three broods of hens that nested in the area. In 2001, one hen produced a successful brood. A brood was considered successful if  $\leq 1$  chick was recruited into the August population. In 2002, one of two broods monitored was successful. In both years, most chicks were lost within the first 14 days post-hatch.

Figure 4.

Table 5. Vegetation classifications for Gunnison Sage-grouse brood-rearing habitat, San Juan County, UT, 2002.

Vegetation Classification	Acreage	% Total	Objective
Non-irrigated	8987	11%	
Black-sage	4287	5%	
Big sage < 25%	4352	5%	
Big Sage > 25%	7704	10%	> 20-40%
Grass/CRP	34,144	42%	30%
Rangeland (Grass)	5536	7%	

Habitat selection by broods was combined for both years because of sample size limitations. Hens with broods selected CRP/grassland and big sagebrush more than any other cover type (Table 6). Agriculture, black sage, grazed rangelands, and bare ground were not used in proportion to their availability. The woodland/mountain shrub cover class was used in equal proportion to its availability. Black sage was only used in 2001; agriculture and woodland/mountain shrub were only used in 2002.

Table 6. Percent availability and use of cover types by sage-grouse hens with broods in 2001 and 2002 in San Juan County, UT.

Cover Type	Year					
	2001			2002		
	Availability	Use	<i>n</i>	Availability	Use	<i>n</i>
Wet meadow	0	0	0	0	0	0
Agriculture	0	0	0	27	11 <sup>-a</sup>	2
Woodland/mtn. shrub	0	0	0	1	5	1
Black sage	85	8-	2	0	0	0
Big sagebrush	3	19+	5	2	11+	2
CRP/grassland	11	73+	19	25	74+	14
Rangeland	1	0-	0	0	0	0
Bare ground	1	0-	0	0	0	0

<sup>a</sup> + = Use greater than availability, no symbol = use in proportion to availability, - = use less than availability ( $P \leq 0.0001$ ) by Bonferroni confidence intervals.



The vegetation composition at brood-use sites in 2001 did not differ from non-use sites ( $P = 0.0001$ ). Brood-use sites exhibited greater forb cover and less grass and shrub cover than non use sites (Table 7). Percent cover of grasses, forbs, and shrubs at brood-use sites did not differ between years ( $P = 0.002$ ). Brood-use sites in 2001 had greater percent canopy cover of all vegetation types (Table 7). Although the vegetation height at brood rearing sites differed between years, the difference was not significant ( $P = 0.07$ ) (Table 6).

Table 7. The vegetation height (cm and in) and percent cover of grasses, forbs, and shrubs at brood-use sites in 2001 and 2002.

	2001	2002
Height	18.8 (7.8 in)	12.2 (5.1 in)
% Grasses	14.8	5.7*
% Forbs	9.5	1.7*
% Shrubs	6.1	2.8*

\*  $P$ -values  $\leq 0.05$  are significantly different.

In 2001, we monitored four radio-collared males. All individuals were captured from the Roring lek site (Figure 2). They were often observed in the same flock. Flock size ranged from 2-18 individuals. We observed no mixed-sex flocks in 2001. In 2002, we monitored six radio-collared males and five radio-collared hens that did not have broods. Three of the males were the same individuals monitored in 2001. The other three males were captured on the Hickman Flat lek site. One of the broodless hens in 2002 was the brooding hen from 2001. The other four broodless hens were captured in 2002, three on the Roring lek and two on the Hickman Flat lek. The birds captured on the Hickman Flat lek were observed in mixed flocks while individuals captured on the Roring lek were only found in single-sex flocks. Mixed-sex flocks ranged in size from 4-18 individuals, single sex flocks had 1-16 individuals.

Percent cover of grasses, forbs, and shrubs at brood-use sites in 2001 differed from sites used by radio-collared males in the same year ( $P = 0.002$ ). Sites used by males had greater grass cover than those used by the brood; the brood used sites with greater forb and shrub cover (Table 7). The height of vegetation at brood sites did not differ at sites used by males in 2001 ( $P = 0.06$ ) (Table 8). Vegetation height did not differ between brood-use sites and sites used by males in 2002 ( $P = 0.5$ ).

In both years, males selected the CRP/grassland and big sagebrush cover types (Table 9). In 2001, agriculture, woodland, and bare ground cover types were avoided; rangelands were used in proportion to their availability. In 2002, wet meadow, agriculture, black sagebrush, and bare ground cover types were avoided; woodlands and rangelands were used in proportion to their

availability.

The percent cover of grasses, forbs and shrubs at sites used by males in 2001 did not differ from reference sites ( $P = 0.1$ ) (Table 9). Reference sites had slightly greater percent cover of grasses, forbs, and shrubs than sites used by Gunnison Sage-grouse males.

Table 8. The vegetation height (cm and in) and percent cover of grasses, forbs, and shrubs at brood-use sites and male Gunnison Sage-grouse locations in San Juan County, UT, 2001.

	Brood	Males
Height	18.8 (7.8in)	23.6 (9.8in)
% Grasses	14.8	28.3*
% Forbs	9.5	8.1*
% Shrubs	6.1	3.3*

\* P-value  $\leq 0.05$  significantly different.

Table 9. Simultaneous confidence intervals using the Bonferroni approach for utilization of vegetation types by male Gunnison Sage-grouse in 2001 and 2002.

Cover Type	2001			2002		
	Confidence Interval		Expected	Confidence Interval		Expected
	Lower	Upper		Lower	Upper	
Wet meadow	0.000	0.000	0.000	0.000	0.000	0.001- <sup>a</sup>
Agriculture	0.113	0.215	0.331-	0.036	0.091	0.104-
Woodland/mtn. shrub	0.000	0.000	0.028-	0.000	0.017	0.003
Black sagebrush	0.000	0.000	0.000	0.099	0.177	0.266-
Big sagebrush	0.092	0.188	0.067+	0.077	0.149	0.055+
CRP/grassland	0.623	0.751	0.565+	0.561	0.671	0.426+
Rangeland	0.000	0.022	0.009	0.010	0.048	0.020
Bare Ground	0.000	0.000	0.001	0.013	0.014	0.127-

<sup>a</sup> + = Use greater than availability, no symbol = use in proportion to availability, - = use less than availability ( $P \leq 0.0001$ ) by Bonferroni confidence intervals.

Table 10. Percent cover of grasses, forbs, and shrubs at reference sites and male

## Gunnison Sage-grouse locations in San Juan County, UT, 2001.

	<u>Males</u>	<u>Reference</u>
Grasses	28.3	33.2
Forbs	8.1	9.1
Shrubs	3.3	10.9

The percent cover of grasses, forbs, and shrubs at brood-use sites in 2002 did not differ from sites used by hens without broods ( $P = 0.5$ ) (Table 10). The height of vegetation at brood-use sites in 2002 did not differ from sites used by hens without broods ( $P = 0.06$ ). In addition, percent cover of grasses, forbs, and shrubs at brood-use sites in 2002 did not differ from sites used by radio-collared males ( $P = 0.4$ ) (Table 11).

Table 11. Percent cover of grasses, forbs, and shrubs used by broods, broodless hens, and male Gunnison Sage-grouse in San Juan County, UT, 2002.

	<u>Brood</u>	<u>Broodless Hen</u>	<u>Males</u>
Height	12.2	20.1	14.5
Grasses	5.7	3.6	4.0
Forbs	1.7	1.2	2.3
Shrubs	2.8	11.6	3.8

Hens without broods did not use wet meadow, black sagebrush, and bare ground cover types in proportion to their availability. Agriculture and big sagebrush cover types were used relative to availability (Table 12).

Table 12. Simultaneous confidence intervals using the Bonferroni approach for utilization of vegetation types by non-brood rearing Gunnison Sage-grouse in 2002.

Cover Type	<u>Confidence Interval</u>		Expected
	Lower	Upper	

Wet meadow	0.0000	0.0000	0.0007 <sup>-a</sup>
Agriculture	0.0372	0.1098	0.0743
Woodland	0.0000	0.0247	0.0017 <sup>+b</sup>
Black sagebrush	0.0734	0.1633	0.2964 <sup>-</sup>
Big sagebrush	0.1008	0.2004	0.1444
CRP/grassland	0.5361	0.6722	0.4402 <sup>+b</sup>
Rangeland	0.0147	0.0712	0.0091
Bare ground	0.0000	0.0000	0.0333 <sup>-</sup>

<sup>a</sup> - = use less than availability, no symbol = use in proportion to availability ( $P \leq 0.0001$ ) by Bonferroni confidence intervals.

<sup>b</sup> + = use greater than availability ( $P \leq 0.05$ ) by Bonferroni confidence intervals.

### *Use of wet meadows*

The PLAN established an objective of maintaining vegetation height in wet meadow areas of 4 inches or greater between June 15 - July 31 on over 75% of the area used as brood-rearing habitat. Based on analysis of LandSat imagery, we classified approximately 3700 acres within the core conservation area a wet meadow habitats (Figure 2). However, the broods we monitored did not frequent these areas. This observation could be attributed to the severe drought the area has been experiencing. This drought has affected water levels and dried up most of the wet meadows adjacent to nest and brood-rearing sites. Activities implemented by SWOG to address this issue are discussed under Conservation Strategies.

### **3. Wintering complexes**

An objective of the PLAN is to reestablish desired vegetation conditions on 50% of the areas located within the conservation study area, and 25% of the vegetation conditions within the buffer areas, over the next 10 year period. Desired conditions for winter habitat include a minimum of 15% canopy cover of big sagebrush vegetation that averages 12 inches in height on southerly and westerly aspects. Big sagebrush in drainages should exhibit a minimum of 30% canopy cover and average 20 inches in height. Small areas that exhibit denser sagebrush canopy cover (40% with an average height of 16 inches) should be interspersed throughout the wintering area on south and west slope aspects.

### *Current Status*

We are currently conducting an assessment of winter habitat in the SA. This work will be conducted through 2005.

### **4. Corridors**

An objective of the plan is reestablish and maintain contiguous travel corridors consisting of big sagebrush exhibiting >25% canopy coverage between breeding, brood rearing, and wintering complexes in the core area. This objective is to be achieved within the next 15 years.

### *Current Status*

We are using information gathered from our Gunnison Sage-grouse habitat-use studies to identify and map corridors used by the San Juan County population to move between seasonal habitats. Based on movements of adult males and females, and juvenile females there appears to be little or no population interchange between birds who frequent the Roaring lek site located in the western part of the SA and birds who use the Hickman lek complexes in the eastern part of the CSA (Figure 3). Location data obtained from recent telemetry flights conducted to determine winter use sites indicate that two birds from the Roring lek have moved in close proximity of birds from the Hickman Flat lek. This suggests that the populations may interchange. Additional effort will be conducted this spring to capture juvenile males. Juvenile males are more likely to disperse to new lek sites.

### III. PLAN IMPLEMENTATION - CONSERVATION STRATEGIES

#### A. Desired Plan Outcomes

Implementation of the PLAN will result in a broad base of local support necessary to coordinate management across land ownership and jurisdictional boundaries to ensure survival of the Gunnison Sage-grouse and the economic viability of San Juan County. To achieve this outcome, the PLAN has been designed to be a dynamic document that will be formally reviewed annually and updated as new information becomes available. Annual progress and quarterly reports have been provided to SWOG members.

The success of this PLAN will be measured by changes in habitat conditions and Gunnison Sage-grouse population numbers in San Juan County. Another measure of success will be increased participation of local landowners, the community, public and private resource management and conservation agency and organization efforts in conservation actions.

#### B. Conservation Strategies

The strategies identified in the PLAN will be implemented and evaluated by SWOG. Although the strategies may be applied to approximately 336,680 acres identified as the core conservation sage-grouse habitat, priority will be placed on areas within the 59,744 acre conservation study area. The conservation study area contains essential Gunnison Sage-grouse breeding, nesting, and brood rearing habitat.

#### C. Status Priority Implementation Actions by Conservation Strategy

##### 1. Develop public support and funding base for the conservation plan.

**Action:** Communicate Conservation Plan goals, objectives, and accomplishments to other stakeholders in the agricultural, natural resources, and legislative community.

**Strategies:** Publish a San Juan County Gunnison Sage-grouse Conservation Plan informational brochure for public distribution.

**Status:** An informational brochure describing the conservation planning efforts was printed and distributed in 2001 (Appendix D)

**Host media interviews with SWOG representatives from the agricultural and wildlife conservation communities.**

**Status:** SWOG members have been interviewed by local newspapers, the Salt Lake Tribune, Deseret News, and other media outlets.

**Organize and conduct an annual San Juan County Gunnison Sage-grouse Conservation Festival to increase public awareness and support for local community efforts to restore the populations.**

**Status:** The Gunnison Sage-grouse Festival concept has been postponed until SWOG is able to assess the effects of increased public visitations on lek attendance. The UDWR in cooperation with Utah State University has hosted small group tours and lectures about Gunnison Sage-grouse conservation. The small, unadvertised tours appear to be meeting the demand for viewing the birds.

**Develop a series of promotional items that carry a designer logo. These promotional items will be sold at local and regional retail outlets to increase the visibility and support of the county conservation efforts and generate revenue to support PLAN implementation.**

**Status:** A local designer was contacted about developing a logo that could be used on caps and other apparel. Three dozen caps were embroidered with the logo and distributed to local landowners. In addition, we contacted a company known as Rag Bag, Inc. ([www.ragbag.com](http://www.ragbag.com)) to develop additional promotional items. The company currently has a series of promotional clothing available that can be ordered on-line.

## **2. Monitoring and Evaluation**

**Action:** **Seek endorsement and funding of the San Juan County Gunnison Sage-grouse Conservation Plan through the Utah Department of Natural Resources Endangered Species Mitigation Fund, the Utah Legislature, and other sources.**

**Strategies:** **SWOG meets as a working group every six months to review PLAN progress and implementation.**

**Status:** SWOG has met seven times since the Conservation Plan was signed in March 2001. The working group uses the meetings to discuss conservation activities and plan future activities.

**SWOG partners will contact their legislative representatives regarding the process and send letters of support to the executive director of the Department of Natural Resources, the Utah Governor's Office, and Utah's congressional delegation.**

**Status:** SWOG members have contacted legislative and congressional representatives. Copies of the signed plan have been distributed to the local legislators, the Utah Governor's Office and Utah's Congressional delegation. SWOG members have written letters to the Executive Director of the Utah Department of Natural Resources to support the use of Utah Endangered Species Mitigation Funds to support the Adam's Conservation Easement. Funding was received from the Fund to support the easement.

**SWOG representatives will testify before the appropriate Utah Legislature committees about the San Juan County Conservation Plan to increase legislator awareness and support for similar efforts in other areas of Utah.**

**Status:** See above comments



**Action:** Monitor impacts of conservation strategies on sage-grouse habitat and population.

**Strategies:** SWOG will initiate an ongoing research program to monitor annual sage-grouse population numbers and trends, monitor sage-grouse habitat use, nest success, mortality, and identify land uses action which may conflict with the goals and objectives of the PLAN. This work began in 2000.

**Status:** Graduate students from Utah State University have been monitoring Gunnison Sage-grouse populations and habitat-use pattern in response to management actions. The results of this research are reported in the update. Utah State University will continue this monitoring effort. The work is supported by funds provided by the UDWR, Bureau of Land Management, Quinney Professorship for Wildlife Conflict Management, Utah State University Extension, and the Jack H. Berryman Institute.

### 3. Species protection and population enhancement

**Action:** Monitor landscape sage-grouse habitat conditions to include land use and vegetation changes in the conservation area.

**Strategy:** Utah State University will update the existing GIS land use database of the CA landscape vegetation and habitat conditions every five years. This update will allow SWOG to compare pre- and post-PLAN time periods to inventory and map habitat changes that resulted because of conservation strategy implementation.

**Status:** This work is continuing. In the update, we report on changes in vegetation conditions that have occurred since 1998.

**Action:** Delineate and map all lek sites within the conservation area, monitor numbers of strutting cocks, estimate population numbers and trends, and determine priority brood rearing and wintering complexes.

**Status:** This work is on-going. The results of our effort to date are reported in this update.

**Strategy:** Conduct annual lek, brood, and winter surveys. Priority areas identified will be added to the SWOG GIS data base. GPS locations of all nesting, brood rearing, and wintering complex will be recorded and sites delineated on the SWOG GIS data base.

**Status:** This work is ongoing and has been reported in this update.

**Action:** Increase the abundance and distribution of Gunnison sage-grouse.

**Strategies:** Enhance sage-grouse habitat conditions (See restoring and improving habitat quality section).

**Status:** This is reported on in the habitat section

**Implement a predation management program.**

**Status:** For FY 2001, USDA Wildlife Services (WS) removed 21 coyotes prior to the breeding season. During this same period they placed 21 DRC-1339 eggs to protect the birds from raven predation. In FY 2002, WS removed 46 coyotes and place 291 DRC-1339 treated eggs. The eggs were placed in draw station to attract ravens. The county also offers a bounty on coyotes.

Currently, the Gunnison Sage-grouse monitored are exhibiting high survival rates. However, we cannot state if the high survival rates are directly attributed to the predation management efforts.

**Reintroduce sage-grouse obtained from Colorado into restored habitats.**

**Status:** SWOG has discussed this option with Colorado. The merits of this action are being assessed and prioritized in the rangewide plan that is being drafted.

#### **4. Restoring and Improving Habitat Quality (Figure 4)**

**Action:** Develop a vegetation management PLAN for the core area.

**Strategies:** Identify and GIS map existing and potential nesting, brood rearing, wintering areas, and travel corridors to include land ownership.

**Status:** Utah State University is continuing to do this work.

**Work with SWOG partners to manage core and buffer areas to achieve defined sage-grouse habitat objectives.**

**Status:** SWOG partners have completed a number of habitat projects (Figure 4). These projects include sagebrush seedings, rabbitbrush treatments on the Adam's Easement and the development of guzzlers and sprinkler systems to provide water sources and wet meadow habitats.

We are continuing to monitor Gunnison Sage-grouse use of the water developments. In 2002, Utah State University purchased two remote sensing cameras to document bird use and other activity at the watering facilities.

A complete list of the projects accomplished and the costs can be found in Appendix E. This work was accomplished under a cost-share agreement with the Natural Resource Conservation Service Wildlife Habitat Improvement Program.

**Work with SWOG partners to develop and implement grazing management plans to achieve Gunnison sage-grouse habitat objectives.**

**Status:** In early 2002, the state of Utah declared a drought emergency for San Juan County. The state petitioned USDA to open CRP land to domestic livestock grazing. The petition was granted. Several of our radio-collared sage-grouse inhabited some of the CRP land in the county that was opened to grazing. We monitored the habitat-use patterns of these birds in response to the presence of livestock. We also build exclosures in each CRP field that was grazed. These exclosures will allow us to assess the effects of grazing on future vegetation growth. These data will be available in the spring of 2003.

In addition, 1280 acres that are part of the Adam's easement are being managed under a prescribed grazing program to enhance Gunnison Sage-grouse habitat conditions. By 2004 an additional 640 acres will be grazed according to established guidelines.

**Work with the USU County Extension Office and the Utah Department of Agriculture and Food to identify and manage noxious weed species to improve sage-grouse habitat and livestock productivity.**

**Status:** We are working with the County Extension Office and the Natural Resource Conservation Service to monitor noxious weed infestation in the county.

**Action:** **Protect critical lek, nesting, brood rearing, and wintering areas.**

**Strategies:** **Secure or acquire important habitats through fee title from willing sellers, land exchanges, conservation easements, tax incentives,**

**voluntary cooperative agreements, CRP leases, grazing lease agreements, etc.**

**Status:** The UDWR purchased a conservation easement from Bruce Adams (Figure 4). The easement covers 2,244 acres. The total cost of the easement was \$336,600. The Department of Natural Resources Endangered Species Mitigation Fund contributed \$182,600 toward the purchase. Other partners included the U.S. Fish and Wildlife Service (\$99,000), The Nature Conservancy (\$5,000), and an anonymous natural gas pipeline company (\$50,000).

The easement protects one lek site and important sagebrush cover. The UDWR is currently negotiating a second easement to protect the Roaring lek site.

## **5. Reducing Physical Disturbance**

**Action:** **Disturbance that negatively impacts sage-grouse will be identified and managed. This includes predation management, recreation use, construction and surface disturbances, and other uses that may conflict with critical biological periods.**

**Strategies:** **Delay or modify construction start up dates or hours to minimize impacts in sage-grouse nesting and brood rearing areas.**

**Status:** When the UDWR reviews construction permits or applications for gas, oil, and/or pipeline developments in the study area, it identifies essential Gunnison Sage-grouse habitats and activity periods that the contractors should avoid.

**Designate OHV use areas and other requirements.**

**Status:** This map is currently being worked on. As soon as data regarding winter use areas are available, the map will be completed.

**Manage off-road travel in key sage-grouse areas.**

**Status:** See the above comment

**Implement predation management in key nesting, brooding rearing and wintering area.**

**Status:** USDA Wildlife Services is conducting a predation management program during critical periods



#### D. Implementation Schedules and Responsibilities (Revised December 2002)

##### San Juan County Gunnison Sage-grouse Working Group (SWOG) Conservation Strategies.

Conservation and Management Strategies	Examples of ways to accomplish	Implementation Schedule	
		When	Who
Develop Public Support and Funding Base			
Provide information to the public, landowners and others that identifies sage-grouse habitat needs, conditions, and sage-grouse population levels. Identify concerns and opportunities to improve conditions for sage-grouse in this area.	Maps, newspaper articles, videos.	2000 - ongoing	SWOG, UDWR, USU, BI
	Meetings with interested landowners.	2000 - ongoing	
	Publish a San Juan County Gunnison Sage-grouse Conservation Plan Brochure.	Completed	
Work with interested parties, landowners, and others to bring about a better understanding of sage-grouse needs, including the value and importance of sage-grouse and sage-grouse habitat, and provide a basis for sharing ideas and reaching agreement on ways to improve sage-grouse habitat and increase populations.	Meetings with interested landowners, government/regulatory entities (e.g. countries).	2000 - ongoing	SWOG
	Developing management plans, cooperative agreements, etc.	2000 - ongoing	SWOG
	Distribute information on: importance of sage-grouse; availability of incentive programs, BMPs, effects of certain land uses on grouse.	2000 - ongoing	SWOG, UDWR, USU Extension, NRCS
	Coordinate sage-grouse conservation actions with management plans for other wildlife species in San Juan County.	2000 - ongoing	SWOG, UDWR, USU Extension
	Continue to work with other groups: Nature Conservancy, Envision Utah, RMEF, Soil Conservation Districts, Utah Farm Bureau, etc., in Colorado to further sage-grouse conservation efforts region wide.	2000 - ongoing	SWOG, UDWR, NRCS, Private Landowner, USU Extension, BLM, NRCS, FWS
	Communicate with other sage-grouse working groups.	2000 - ongoing	SWOG

Conservation and Management Strategies	Examples of ways to accomplish	Implementation Schedule	
		When	Who
Incorporate economic, social, and cultural values into conservation practices.	Communication with San Juan County Commission and Communities.	2000 - ongoing	SWOG
	Encourage voluntary compliance and participation.	2000 - ongoing	SWOG
	Involve landowners and local communities in all aspects of sage-grouse conservation.	2000 - ongoing	SWOG
	San Juan County Sage-grouse Festival and promotional items.	2004	SWOG, Utah Tourism, USU Extension, San Juan County Commissioner, UDWR
Maintain local control.	The Sage-grouse Working Group (must include landowners and local residents) and will act as advisory body to the County Commission and agencies.	2000 - ongoing	SWOG
	Provide for continual public input and involvement.	2000 - ongoing	SWOG
Develop, improve, and encourage credibility and success.	Seek outside scientific review of projects.	As opportunities arise	USU Extension, UDWR, BI
Seek endorsement and funding for conservation plan	Meet with Utah legislature and congressional representatives.	2000 - ongoing	SWOG, UDWR, USU Extension
	Meet with Utah Department of Natural Resources.	2000 - ongoing	SWOG, UDWR, USU Extension

Conservation and Management Strategies	Examples of ways to accomplish	Implementation Schedule	
		When	Who
Monitoring/Evaluation			SWOG
SWOG will meet every 6 months to review Plan implementation and progress	Meetings will be held in San Juan County with the working group and the landowners.	2000 - ongoing	UDWR, USU Extension, BI
Identify important sage-grouse habitat, limiting factors, and activities that have the potential to impact sage-grouse or their habitat. Identify and evaluate critical sage-grouse habitats.	Habitat mapping and monitoring.	2000 - ongoing	
	Meetings with interested landowners.	2000 - ongoing	
	Joint-interagency/landowner evaluation, information sharing.	2000 - ongoing	
Continue to gather or initiate the collection of basic resource data to better understand and document conditions for sage-grouse, including response habitat.	Hire a graduate student to collect baseline habitat and population data.	2000 - ongoing	SWOG, UDWR, USU, BI, WS, USFWS, BLM
	Habitat mapping and monitoring.	2000 - ongoing	
	Meetings with interested landowners.	2000 - ongoing	
	Joint-interagency/landowner evaluation, information sharing.	2000 - ongoing	
Species Protection and Population/Habitat Enhancement			
Develop and encourage incentives for landowners and others to avoid or mitigate loss of sage-grouse habitat.	Land exchanges.	2000 - ongoing	UDWR, FWS, SWOG, BLM, FS, Nature Conservancy, UDWR, NGOs
	Conservation Easements/Leases.	As opportunities arise	UDWR, NGOS
	Transferrable development rights.	As opportunities arise	UDWR, NGOS
	Payment for non-use.	2000 - ongoing	UDWR, FWS, SWOG, NGOS
	Application of specific land-use practices that benefit grouse.	2000 - ongoing	Landowners
	Reintroduce sage-grouse to restored habitats.	As opportunities arise	UDWR, NRCS, SWOG, Private Landowner



Conservation and Management Strategies	Examples of ways to accomplish	Implementation Schedule	
		When	Who
Enhance and restore historic and existing sage-grouse habitat to offset loss of habitat elsewhere.	<p>Reseeding or reclaiming areas, creating or protecting wet meadow areas, and implementing vegetation treatments (i.e., prescribed burning, Dixie harrowing, etc.) to rejuvenate habitats and maintain leks.</p> <p>Reintroduce sage-grouse into restored habitats.</p>	<p>Ongoing</p> <p>As opportunities arise</p>	<p>SWOG, UDWR, NRCS, USFWS Landowners, BLM</p> <p>UDWR</p>
<p>Prevent loss and fragmentation of habitat from construction of roads, utilities.</p> <p>Identify changes to county land use regulations which would benefit sage-grouse.</p>	<p>Relocation or modification of new utility lines, roads, development, etc. in key grouse habitat and provide recommendations to the county or lead agency.</p> <p>Pipeline or power line modifications.</p> <p>For example: seek a county resolution that supports and encourages the use of conservation easements, mitigation of non-critical habitat areas, and preventing loss of critical habitat.</p>	<p>As opportunities arise</p> <p>Ongoing</p> <p>Ongoing</p>	<p>SWOG, San Juan County Commissioner</p> <p>BLM, UDWR, San Juan County Commission</p> <p>SWOG, San Juan County Commission</p>
Restoring or Improving Habitat Quality			
Enhance existing riparian areas by creating or enhancing small wet areas to benefit sage-grouse nesting and brood rearing habitat.	<p>Design and implement livestock grazing management practices to benefit riparian areas.</p> <p>Modify or adapt pipelines/springs to create small wet areas.</p>	<p>2000 - ongoing</p> <p>As opportunities arise</p>	<p>USU Extension, UDWR, FWS, NRCS</p> <p>Pipeline Companies, UDWR, BLM, FWS, and landowners</p>
Reduce or modify situations that cause predation.	<p>Modify power lines and wood fence posts (to remove raptor perches) in critical sage-grouse areas.</p> <p>Cut pinyon-juniper trees near leks and elsewhere within potential sage-grouse habitat to remove raptor perches.</p>	<p>As opportunities arise</p> <p>2000 - ongoing</p>	<p>UDWR, BLM, SWOG, USDA, Wildlife Services, FWS</p> <p>UDWR, BLM, SWOG, USDA, Wildlife Services, FWS</p>

Conservation and Management Strategies	Examples of ways to accomplish	Implementation Schedule	
		When	Who
Develop and use Best Management Practices to guide land uses to increase sage-grouse populations and improve sage-grouse habitat.	Implement local guidelines that describe:	2000 - ongoing	SWOG, UDWR, BLM, USU Extension, NRCS
	Livestock grazing practices that benefit sage-grouse.	As opportunities arise	SWOG, UDWR, BLM, USU Extension, NRCS
	Living with sage-grouse in your backyard (control of dogs, etc.).	Ongoing	SWOG, UDWR, BLM, USU Extension, NRCS
	Restoring and rehabilitating riparian areas.	As opportunities arise	SWOG, UDWR, BLM, USU Extension, NRCS
	Proper land treatment design and construction that reduce impacts to sage-grouse (e.g., how and where to Plan projects).	As opportunities arise	SWOG, UDWR, BLM, USU Extension, NRCS
	Land development options.	As opportunities arise	SWOG, UDWR, BLM, USU Extension, NRCS
Improve sage-grouse habitat quality and improve vegetation cover, especially forbs and grasses in sage-grouse areas.	Developing and using sound grazing management practices.	2000 - ongoing	NRCS, USU Extension, UDWR Landowners
	Planting and re-seeding with a high proportion of forbs.	2000 - ongoing	SWOG
	Designing vegetation treatments in sage-grouse areas to be compatible with sage-grouse needs.	2000 - ongoing	UDWR, NRCS
	Improving ground cover in sage-grouse areas.	2000 - ongoing	UDWR, NRCS, BLM, Landowners
	Managing big game to avoid degrading sage-grouse habitat or recovery.	2000 - ongoing	UDWR
	Integrating weed management with grouse needs.	2000 - ongoing	USU Extension, BLM, Landowner, NRCS

Conservation and Management Strategies	Examples of ways to accomplish	Implementation Schedule	
		When	Who
Increase opportunities for over-winter survival, escape cover near leks, nesting cover.	Improve quality of sagebrush dominated habitats by grazing management.	2000 - ongoing	SWOG, NRCS, UDWR, USU Extension
	Avoiding treatment projects that remove large strands of sagebrush in critical areas.	As opportunities arise	SWOG, NRCS, UDWR, USU Extension
	Developing recommendations for managing sagebrush community as a whole, considering all uses.	Ongoing	SWOG, NRCS, UDWR, USU Extension
Reducing Physical Disturbance to Sage-grouse/Predation Management			
Mitigate or reduce conflicts with sage-grouse during critical biological periods and in critical habitats.	Delay or modify construction start up dates or hours to minimize impacts in sage-grouse nesting and brood rearing areas.	2000 - Ongoing	San Juan County Commission
	Designate OHV use areas and other requirements.	As opportunities arise	San Juan County Commission
	Manage off-road travel in key sage-grouse areas.	2000 - Ongoing	San Juan County Commission
	Implement predation management in key nesting, brooding rearing and wintering area.	As opportunities arise	USDA Wildlife Services, UDWR
	Restrict public observation/lek viewing to 1 or 2 leks with 20-25 breeding males. All lek viewing would be conducted in accordance with established protocols to avoid disturbance	2000 -Ongoing	Landowners, UDWR, BLM

#### IV. REFERENCES

- Beck, J. L. and D. L. Mitchell. 1997. Brief guidelines for maintaining and enhancing sage-grouse habitat in private lands in Utah. Utah Division of Wildlife Resources Report. Salt Lake City, Utah. 9 pp.
- Braun, C. E., T. Britt. And R. O. Wallestad. 1977. Guidelines for maintenance of sage-grouse habitat. Wildlife Society Bulletin 5:99-106.
- Commons, M. L. 1997. Movement and habitat use by Gunnison sage-grouse (*Centrocercus minimus*) in southeastern Colorado. M.S. Thesis. University of Manitoba, Winnipeg. 108 pp.
- Dunn, P.O. and C.E. Braun. 1986. Late summer-spring movements of juvenile sage-grouse. Wilson Bulletin 98:83-92.
- Klebrenow, D.A. 1969. Sage-grouse nesting and brood habitat in Idaho. J. Wildl. Manage. 33:649-662.
- Wallestad, R.O. 1971. Summer movements and habitat use by sage-grouse broods in Central Montana. J. Wildl. Manage. 35:129-136.
- Young, J.R., J.W. Hupp, J.W. Bradbury, and C.E. Braun. 1994. Phenotypic divergence of secondary sexual traits among sage-grouse, *Centrocercus urophasianus*, populations. Anim. Behav. 47:1353-1362.

**V. APPENDICES****A. San Juan County Gunnison Sage-grouse Working Group Members**

**Bureau of Reclamation**  
**Farm Services Agency**  
**Natural Resource Conservation Service**  
**San Juan County Commission**  
**San Juan County Extension Office**  
**San Juan County Landowners**  
**Utah Division of Wildlife Resources**  
**Utah State University Extension Service**  
**Utah State University College of Natural Resources**  
**Jack H. Berryman Institute**  
**U.S.D.A. Wildlife Services**  
**U.S. Fish and Wildlife Service**

## B. San Juan County Gunnison Sage-grouse Conservation Area Vegetation Classifications

1. **Surface Water**--class includes areas of open water.
2. **Wet meadows**--class includes drainages, ephemeral streams, creeks, springs, and other riparian areas. Commonly associated plant species include; *Carex spp.*, *Typha spp.*, *Scirpus spp.*, *Salix spp.*, *Artemisia tridentata*, and other forbs and grasses.
3. **Irrigated agriculture**--class includes irrigated agriculture fields, mainly alfalfa, *Medicago spp.*
4. **Non-irrigated agriculture**--class includes those fields in some sort of dry land farming.
5. **Urban**--class includes urban areas.
6. **Pinyon/Juniper**--class includes those areas where pinyon (*Pinus edulis*) and/or Utah Juniper (*Juniperous osteosperma*) comprise more than 15% of the total vegetation in a given area. Commonly associated plant species include: *Artemisia tridentata*, *Chrysothamnus spp.*, *Quercus gambelii*, *Gutierrezia sarothrae*, *Purshia tridentata*, *Amelanchier alnifolia*, *Opuntia spp.*, *Cordylanthus wrightii*, *Poa spp.*, *Aristida spp.*, *Bromus tectorum*, *Stipa spp.*, *Oryzopsis hymenoides*, *Aster spp.*, and Crypto-gramic crust.
7. **Black Sage**--class includes those areas where *Artemisia nova* is the dominate vegetation. Commonly associated plant species include: *Artemisia tridentata*, *Chrysothamnus spp.*, *Gutierrezia sarothrae*, *Atriplex canescens*, *Opuntia spp.*, *Cordylanthus wrightii*, *Poa spp.*, *Bromus tectorum*, *Agropyron cristatum*, *Agropyron smithii*, *Ceritoides lanata*, *Aster spp.*, and Crypto-gramic crust.
8. **Pinyon/Juniper-mountain shrub**--class includes those areas which contain less than 15% Pinyon pine (*Pinus edulis*) and/or Utah Juniper (*Juniperous osteosperma*) and greater than 25% shrubs. Commonly associated plant species include; *Artemisia tridentata*, *Cercocarpus montanus*, *Chrysothamnus spp.*, *Quercus gambelii*, *Gutierrezia sarothrae*, *Purshia tridentata*, *Amelanchier alnifolia*, *Opuntia spp.*, *Cordylanthus wrightii*, *Ceritoides lanata*, *Wyethia amplexicalis*, *Poa spp.*, *Aristida spp.*, *Bromus tectorum*, *Stipa spp.*, *Oryzopsis hymenoides*, *Agropyron cristatum*, *Agropyron smithii*, *Aster spp.*, and Crypto-gramic crust.
9. **Big Sage >25% canopy cover**--class includes those areas where big sagebrush (*Artemisia tridentata*) is the dominate vegetation type. Commonly associated plant species include; *Chrysothamnus spp.*, *Artemisia nova*, *Gutierrezia sarothrae*, *Atriplex canescens*, *Purshia tridentata*, *Opuntia, spp.*, *Cordylanthus wrightii*, *Ceritoides lanata*, *Wyethia amplexicalis*, *Poa spp.*, *Aristida spp.*, *Bromus tectorum*, *Bromus carinatus*, *Stipa spp.*, *Oryzopsis hymenoides*, *Agropyron cristatum*, *Agropyron smithii*, *Aster spp.*, and Crypto-gramic crust.
10. **Big Sage 15-25% canopy cover**--class contains those areas where big sagebrush (*Artemisia tridentata*) comprises at least 15% of the vegetation but not more than 25% of

the total vegetation type. In some cases this class may be invading some Conservation Reserve Program (CRP) fields. Commonly associated plant species include; *Chrysothamnus spp.*, *Artemisia nova*, *Gutierrezia sarothrae*, *Atriplex canescens*, *Purshia tridentata*, *Opuntia spp.*, *Cordylanthus wrightii*, *Ceritoides lanata*, *Poa spp.*, *Aristida spp.*, *Bromus tectorum*, *Bromus carinatus*, *Stipa spp.*, *Oryzopsis hymenoides*, *Agropyron cristatum*, *Agropyron smithii*, *Aster spp.*, and Crypto-gramic crust.

11. **Big Sage <15% canopy cover**--class contains those areas where big sagebrush (*Artemisia tridentata*) comprises less than 15% of the vegetation in a given area. In some cases this class may be invading some Conservation Reserve Program (CRP) fields. Commonly associated plant species include: *Chrysothamnus spp.*, *Artemisia nova*, *Gutierrezia sarothrae*, *Atriplex canescens*, *Purshia tridentata*, *Opuntia spp.*, *Cordylanthus wrightii*, *Ceritoides lanata*, *Wyethia amplexicalis*, *Poa spp.*, *Aristida spp.*, *Bromus tectorum*, *Bromus carinatus*, *Stipa spp.*, *Oryzopsis hymenoides*, *Agropyron cristatum*, *Agropyron smithii*, and *Aster spp.*
12. **Mountain Shrub**--class is dominated by Gambel's Oak. Commonly associated plant species include: *Pinus edulis*, *Juniperous osteosperma*, *Artemisia tridentata*, *Cercocarpus montanus*, *Chrysothamnus spp.*, *Quercus gambelii*, *Gutierrezia sarothrae*, *Purshia tridentata*, *Amelanchier alnifolia*, *Opuntia spp.*, *Cordylanthus wrightii*, *Ceritoides lanata*, *Wyethia amplexicalis*, *Poa spp.*, *Aristida spp.*, *Bromus tectorum*, *Stipa spp.*, *Oryzopsis hymenoides*, *Agropyron cristatum*, *Agropyron smithii*, *Aster spp.*, and Crypto-gramic crust.
13. **Big Sage CRP mixture**--class contains a fair amount of *Artemisia tridentata*, but is still dominated by some sort of CRP seed mixture mainly, *Agropyron spp.*, *Bromus carinatus*, and *Medicago spp.* Commonly associated plant species include: *Chrysothamnus spp.*, *Gutierrezia sarothrae*, *Poa spp.*, and *Bromus tectorum*.
14. **CRP >70% canopy cover**--class is dominated by CRP grasses and forbs, mainly *Bromus carinatus*, and *Medicago spp.* Commonly associated plant species include: *Agropyron cristatum*, *Agropyron smithii*, *Agropyron intermedium*, *Chrysothamnus spp.*, *Gutierrezia sarothrae*, *Poa spp.*, *Aristida spp.*, and *Bromus tectorum*.
15. **CRP 40-70% canopy cover**--class is dominated by CRP grasses and forbs, mainly *Agropyron spp.* Commonly associated plant species include; *Agropyron cristatum*, *Agropyron smithii*, *Agropyron intermedium*, *Bromus carinatus*, *Medicago spp.*, *Chrysothamnus spp.*, *Gutierrezia sarothrae*, *Poa spp.*, *Aristida spp.*, and *Bromus tectorum*.

16. **CRP 15-40% canopy cover**--class is dominated by CRP grasses and forbs, mainly *Agropyron cristatum*. Commonly associated plant species include: *Agropyron smithii*, *Agropyron intermedium*, *Bromus carinatus*, *Medicago spp.*, *Chrysothamnus spp.*, Commonly associated plant *Gutierrezia*, *Sarothrae*, *Poa spp.*, *Aristida, spp.*, and *Bromus tectorum*.
17. **Rangelands**--class contains various vegetation types but was grazed too close to the ground to allow vegetation to be placed into other classes. Commonly associated plant species include: *Chrysothamnus spp.*, *Gutierrezia sarothrae*, *Opuntia spp.*, *Cordylanthus wrightii*, *Ceritoides lanata*, *Poa spp.*, *Aristida spp.*, *Bromus tectorum*, *Stipa spp.*, *Oryzopsis hymenoides*, *Agropyron cristatum*, *Agropyron smithii*, and *Aster spp.*
18. **Bare ground**--class contains mainly bare ground and rock where vegetation is less than 15% total canopy cover.
19. **Unknown**--class could not be placed into any of the above classes with the few vegetation training sites collected in November 1997.



**C. Habitat mixture seeded on CRP lands in the San Juan County Gunnison Sage-grouse Conservation Plan Area.**

<b>Species</b>	<b>PLS Lbs/acre</b>
<b>Grasses</b>	
Bluebunch wheatgrass	1.0
Thickspike wheatgrass	1.0
Western wheatgrass	1.5
Crested wheatgrass	0.5
Pubescent wheatgrass	1.0
<b>Legumes/Forbs</b>	
Alfalfa (Rambler)	1.0
Alfalfa (Ladak, Normad)	1.5
Western yarrow	0.12
Lewis flax	0.25
Sainfoin	0.5
Small burnet	2.0
<b>Shrubs</b>	
Wyoming big sagebrush	0.5
Forage kochia	0.5
<b>Total</b>	<b>11.37</b>

D. Gunnison Sage-grouse Conservation Brochure

## E. Summary of Gunnison Sage-grouse Conservation Projects and Costs

Project	Location	Acres	Cost \$	Status
Wildlife Watering Facility Pipeline	Adams Easement	N/A	3,151	Completed
Electric/Conventional Fencing for Livestock Exclusion	Adams Easement	N/A	20,034	Completed
Brush Management/ Rabbit	Adams Easement	150	1,800	Completed
Prescribed Grazing	Adams Easement	1,280	-	Completed
Prescribed Grazing	Adams Easement	640	-	2004
Wildlife Watering Facility Solar Pump	Adams Easement	N/A	1,744	Completed
Wildlife Water Development and Tanks	Adams Easement	N/A	3,244	Completed
Range Planting	Adams Easement	100	1,703	Completed
Upland Habitat Improvement	Adams Easement	2,240	-	Ongoing

Wildlife Guzzlers	Study Area	N/A	1,800	Completed
Brush Management	Adams Easement	240	7,600	Completed
Range Planting	Adams Easement	240	4,258	Completed
Brush Management	Adams Easement	80	1,722	2004
Range Planting	Adams Easement	80	1,022	2004
Conservation Reserve Program	Conservation Area	36,825	1,222,728	Completed
Conservation Easement	Adams Property	2,244	336,600	Completed