




St. George, Utah, Turfgrass Management Calendar

Kelly Kopp, Rick Heflebower and Steven Price



Seeding

Cool-season grasses, such as Kentucky bluegrass or the fescues, may be seeded any time from mid-spring to early fall, but late summer is the optimum seeding time in the St. George, Utah, area. The warmth and dryness of the soil in late summer is more advantageous for seed germination than the wet and cool soil conditions of the spring, allowing for faster germination. In addition, cool-season grasses planted in the late summer will have both the cool fall and spring growing seasons to establish before the heat and dryness of the summer months returns. Annual weed pressure is also lower in the late summer and early fall than during the spring months. Warm-season grasses, such as buffalograss and bermudagrass, should be seeded in the spring or early summer months to take best advantage of the warmer temperatures that they prefer.

Refer to [Preparing for Turfgrass Establishment-Southern Utah](#)

[Turfgrass Cultivars for Utah](#)

Figure 1. This St. George, Utah, area turfgrass management calendar applies to Washington County as well as the southern portion of Kane County. (Image credit: Terence Larson)

Management Calendar

	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
Seeding-Cool Season Grasses												
Seeding-Warm Season Grasses												
Overseeding-Cool Season Grasses												
Sodding												
Fertilization-Cool Season Grasses												
Fertilization-Warm Season Grasses												
Aeration- Cool Season Grasses												
Aeration- Warm Season Grasses												
Preemergent Weed Control												
Broadleaf Weed Control												
Irrigation												

* These are general guidelines based on average weather conditions. Adjustment will be necessary during extremes of temperature or precipitation.

Overseeding

Overseeding bare patches or areas of thinning turf may be accomplished in late winter, early spring, or fall. In the spring and fall, a disk-type seeder may be used to drop seed into slits in the soil.

Alternatively, overseeding may be done following routine aeration practices.

Refer to [Preparing for Turfgrass Establishment-Southern Utah](#)

[Turfgrass Cultivars for Utah](#)

Sodding

Sodding may be done at almost any time of the year provided water is available for irrigation. However, sod should not be laid when the soil is frozen. Sod may be laid prior to freezing, but care should be taken to keep it from drying out during the winter

months. Appropriate soil preparation will also improve sod root growth.

Refer to [Preparing for Turfgrass Establishment-Southern Utah](#)

[Turfgrass Cultivars for Utah](#)

Fertilization

Fertilization is the most important management practice for improving the overall quality of turfgrasses. A soil test may be performed to assess the availability and adequacy of phosphorus and potassium in soils and to obtain application recommendations. However, routine soil tests for nitrogen are not available. Instead, recommended nitrogen applications of 2-4 pounds per 1000 sq. ft. annually are based on turfgrass research performed at Utah State University. Standard recommend-

datations are to apply 1 pound of actual N with each fertilization.

Soil testing is available through the [Utah State University Analytical Laboratory](#) and soil testing kits and instructions are available at all USU County Extension offices.

Mowing

Mowing is the most basic management practice and one of the simplest things that may be done to improve grass appearance. In the St. George area, mowing should start when active growth begins in the spring and should continue until the grass stops growing in the fall. Ideally, no more than 1/3rd of the total length of the grass blades should be mowed off at one time and clippings should be recycled into the lawn. For example, if the grass is 4 inches high, mow no lower than 3 inches. Higher mowing heights improve grass resilience to hot weather. Consider a minimum mowing height of 2.5 to 3 inches. The exception would be the last mowing, which should be shorter (1.5 in.) to help avoid disease issues over the winter/early spring months.

Aeration

Aeration removes plugs of turf and soil creating a system of large pores that facilitate air, water and nutrient movement into the soil. Aeration may also improve soil compaction and encourage microbial activity to break down excessive thatch. Aeration should be timed to coincide with periods of active grass growth (early to mid-spring and/or late summer to early fall for cool season grasses and early to mid-summer for warm season grasses). Core aeration using a hollow tine aerator (3-4" length, 1/4-3/4" diam.) is recommended. For best results, aerate in one direction and once again at 90 degrees to the first direction.

Weed Control*

Establishing and maintaining a healthy, dense turfgrass stand is the best defense against weeds. However, there are several options for weed control should weeds become problematic. Annual weed control (crabgrass, spurge) may be accomplished by applying preemergence herbicides prior to weed seed germination in early to mid-spring. Broadleaf weed control may be accomplished by applying postemergence herbicides during periods of active weed growth. For both preemergence and

postemergence herbicides, choose products labeled for control of the weeds of concern and carefully follow label instructions to avoid non-target plant damage.

[Weed Control Options for Residential Lawns in Utah.](#)

Irrigation

Irrigation is typically necessary to support healthy turfgrass growth in northern Utah, though different turfgrass species have different water requirements. The efficient use of water for irrigation is also critical, not only to conserve limited water resources, but also for the health of the grass plants. In addition to following a schedule based on turfgrass water requirements, irrigate early in the morning to avoid evaporative losses of water and avoid windy times of the day. Also, check irrigation systems monthly for leaks, broken or clogged sprinkler heads, and other maintenance issues.

Month	Irrigation Interval
Startup until April 30	Once every 5 days, IF needed
May	Once every 4 days, IF needed
June	Once every 3 days
July	Once every 2 days
August	Once every 3 days
September	Once every 5 days
October	Once every 7 days
November 1 until shutdown	Once every 10 days, IF needed

Reference Simple Sprinkler Performance Testing fact sheets.

[Simple sprinkler performance testing for Kane County](#)

[Simple sprinkler performance testing for Washington County](#)

Salinity

Issues with excess soil salinity, or the amount of salt within the soil, can be prevalent in the hot and dry valleys of southeastern Utah. Areas with high clay

content soils, poor drainage characteristics, or high water tables may be particularly problematic and deserve special attention. Improperly irrigating turf by providing overly frequent but shallow watering not only wastes water resources and does little to promote overall grass health, but can also increase soil salinity to harmful levels. Following recommended irrigation schedules and providing deep watering can help alleviate salt buildup or even reduce soil salinity to acceptable levels. Different types of turfgrasses vary in their tolerance to salinity and areas with persistent salt issues can benefit from selecting salt tolerant grass varieties.

[Growing Turf on Salt-Affected Soils](#)

***Precautionary Statement:** All pesticides have benefits and risks; however, following the label instructions will minimize the risk and maximize the benefit. Pay attention to the directions for use and follow precautionary statements. Pesticide labels are considered legal documents containing instructions and limitations. Inconsistent use of the product or disregarding the label is a violation of both federal and state laws. The pesticide applicator is legally responsible for proper use.

Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran's status. USU's policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions. Utah State University employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran's status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USU-sponsored events and activities. This publication is issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Kenneth L. White, Vice President for Extension and Agriculture, Utah State University.