



An Alternate Method for Setting Codling Moth Biofix

Diane Alston, Entomologist • Marion Murray, IPM Project Leader
Richard Heflebower, Horticulture Agent, Utah State University Extension

Quick Facts

- An alternate method for setting codling moth biofix (first moth emergence) was found to be effective for Utah's fruit production regions.
- The new method, called "fixed-biofix," uses latitude and elevation of orchard sites, rather than trap-capture, to predict biofix and timing of egg hatch during the growing season.
- The fixed-biofix method is simpler to implement and easier to explain to growers and crop consultants, and can be accessed online.



Fig. 1 Codling moth larva exit hole in apple. Fruits are protected from entry of young larvae by proper timing of management practices.

Codling moth, *Cydia pomonella* (L.) (Lepidoptera: Tortricidae), is the key insect pest of apple, pear, and walnut throughout the world, including Utah (apple and pear) (Fig. 1). Effective management practices are based on accurate determination of the first flight of codling moths in the spring, called "biofix." Biofix is the starting point for a heat-driven development model [degree days (DDs)] that predicts the timing of egg hatch and entrance of young larvae into fruit, the target stages for management. An accurate biofix, plus the DD model, provide orchardists (commercial and home garden) with a convenient tool to determine optimal timing for codling moth management.

CHALLENGES IN BIOFIX-SETTING

The traditional method for setting codling moth biofix is through the use of traps baited with the female sex pheromone, codlemone. Traps are placed in the orchard before the first male moths are expected to emerge in the spring, and checked daily until the date at which the first moths are caught.

To address the challenge of setting an accurate biofix date, researchers at Washington State University found that average codling moth emergence time on a DD-scale starting on January 1 each year could be predicted using a site-specific equation, and negated the need to use traps (Jones et al. 2013).

Use of the trap-based method to set codling moth biofix has a number of disadvantages:

1. Too few traps may be placed in an area to represent the diverse microclimates (slopes and aspects) that influence the timing of first emergence.
2. Traps are checked too infrequently (daily is required).
3. Inclement spring weather may delay moth flight.
4. The use of mating disruption in commercial fruit production has decreased codling moth populations in some regions, especially in the spring, causing an inaccurate delay in the first moth capture in traps.

Sites at elevations less than 1,300 ft fit a simple quadratic equation using only latitude; however, for higher elevation sites, such as in Utah, a multiple regression model using latitude and elevation was required.

A study was undertaken in Utah's apple-producing regions to validate the latitude-elevation model, and compare its precision with actual egg hatch and larval entry into fruit, the key life stage targets for insecticide applications.

METHODS

The equation developed by Jones et al. (2013) was based on data from 366 location-years and 33 locations across the U.S. (including Utah) and three other countries. Including data from locations with elevations greater than 1,300 ft (400 m) required both latitude and elevation to accurately represent the data (Fig. 2). The multiple regression equation that best described the entire dataset was:

$$\text{Mean emergence time (DDs)} = 1755.599 - 66.777 \times \text{latitude} + 0.676 \times \text{latitude}^2 - 0.0347 \times \text{elevation (m)}$$

To validate the accuracy of the regression equation to estimate biofix in a broader spectrum of Utah's apple- and pear-producing regions, 30 orchard location-years in northern and southern Utah were used in the study




Fig. 2 Plot of predicted vs. observed mean first seasonal emergence time [in degree-days (DDs)] of codling moth for the latitude-elevation equation (fixed-biofix) vs. trap capture (trap-biofix) from Jones et al. (2013). Locations in Utah represent numbers 18-21 in the graph.

(Box Elder, Weber, Davis, Utah, Iron, and Washington counties) in 2016-2018. Locations were selected based on the availability of a nearby weather station to obtain temperature data for degree-day calculations, and presence of apple orchards conducive for codling moth sampling. At each location, we determined three different biofixes, which corresponded to the date of:

1. First codling moth capture in traps using CM-DA Combo lures (Trécé Inc., Adair, OK) ("trap-biofix").
2. DDs calculated with the equation at left using elevation and latitude ("fixed-biofix"), with a January 1 start date.
3. DDs calculated with the equation at left using elevation and latitude ("fixed-biofix"), with March 1 start date.

Beginning when egg hatch was predicted using the trap-biofix method, for the 1st and 2nd generations of codling moth (220 and 1,120 DDs, respectively), 1,000 fruit were sampled in each orchard one to three times to verify start of egg hatch and first larval entries into fruit (Figs. 3 & 4).

Fruit were returned to the laboratory to check for egg hatch and larval entry status (Fig. 5). Egg hatch data was collected in 2016 and 2017, but not in 2018.



Fig. 3 Codling moth eggs were found on fruit and foliage. Hand lenses (10-20x) were used for viewing. Eggs in the "red-ring" stage indicate imminent hatch.




Fig. 4 Often, larvae do not travel far from their hatching site to enter fruit.

Table 1. The difference in number of days (range) between the new model (fixed-biofix) using March 1 as the start date, and the trap-biofix for predicting seasonal flight and egg hatch in 30 orchard location-years in northern and southern Utah, 2016-2018. Start and end of egg hatch periods for the 1st and 2nd codling moth generations are provided. Positive numbers indicate equation predictions were earlier than traps; negative numbers indicate equation predictions were later than traps.

Location	Biofix	Start 1 st Egg Hatch	End 1 st Egg Hatch	Start 2 nd Egg Hatch	End 2 nd Egg Hatch
Perry ¹	3 to 6	2 to 3	-2 to 3	-3 to 1	1
Kaysville ²	4 to 21	3 to 9	1 to 6	1 to 8	2 to 4
American Fork ³	0 to 13	0 to 5	0 to 2	0 to 2	0 to 2
Payson ³	-1 to 0	-4 to 0	0	0 to -1	0 to -1
Santaquin ³	1 to 6	0 to 1	0 to 2	-1 to 1	0 to 1
West Mountain ³	-2 to 1	-3 to 2	-1 to 1	-1 to 1	-1 to 1
Genola ³	-1 to 1	1 to 2	0 to 4	1 to 3	1 to 3
Cedar City ⁴	-11 to -1	-5 to 0	-3 to 4	-3 to 3	-3 to 3
New Harmony ⁵	-4 to 6	-9 to 2	-2 to 1	-2 to 1	-2 to 1

¹located in Box Elder County

²located in Davis County

³located in Utah County

⁴located in Iron County

⁵located in Washington County

RESULTS

We found that for the two fixed-biofix options (options 2 and 3 on the previous page), using a start date of March 1 correlated more closely to the trap-biofix in each year; therefore, a March 1 start date was used for data analyses.

To compare the accuracy of the DD predictions based on the fixed-biofix to trap-biofix for the three years (2016-2018), we calculated the difference in number of days for critical codling moth phenological events (e.g., start and end of 1st and 2nd generation egg hatch) (Table 1).

We observed variation in dates between the fixed- and trap-biofix methods across the 10 orchard locations in the three years, ranging from 11 days early to 21 days late. However, this early variation in number of days between methods for biofix became less pronounced over the course of the season. The error rate for predicting start

and end of egg hatch periods for the first and second generation of codling moth across all locations and years was ≤ 9 days. The majority of locations across years did not vary more than 5 days between the two methods, and were often within 3 days.

Our fruit sampling data confirmed that start of egg hatch and first larval entry aligned well with using the fixed-biofix for the codling moth DD model in both 2016 and 2017 (egg hatch data were not collected in 2018).

RECOMMENDATIONS

A convenient and accurate method to set codling moth biofix is critical to effective codling moth management. Our target error rate for differences between the two methods was ≤ 5 days. Although there was a greater range in setting biofix dates for most orchard locations, the predictions for 1st- and 2nd-generation start and end of egg hatch did fall within the target range using the fixed-biofix method for most locations (Table 1).

The chance of differing biofix dates between the two methods is greater because daily DD accumulations around the start of flight are low (e.g., few DDs accumulated per day when temperatures are cool in the spring); therefore, even small variations in DDs can result in a large difference in number of days. However, the DD accumulations at biofix are only a small proportion of DD accumulations needed for the key phenological events of start and end of egg hatch that occur later in the season. The resulting "fixed-biofix" management program is simpler and lower cost to implement, and easier to explain to growers and crop consultants than a trap-biofix method. Given the relatively low variation in prediction of egg hatch between the two methods, the fixed-biofix method is appropriate for use in Utah.



Fig. 5 Newly hatched larvae were found on the fruit's surface and initiating tunnels into the fruit.

The results of this Utah study support the validity of using the Jones' equation to set biofix, with a March 1 start date. This alternative biofix-setting option will help balance issues with spring temperature variation, low codling moth populations, and trap operation errors that lead to challenges in setting accurate biofix dates with pheromone traps.

The fixed-biofix model was included in the Utah TRAPs (Temperature Resource and Alerts for Pests) website in 2018, and tested that season for accuracy.

Predicted emergence, or biofix, timings in DDs for the representative apple and pear production areas of northern and southern Utah are presented in Table 2. Training of Utah's tree fruit industry in the efficiency and accuracy of the equation-based biofix-setting model (fixed-biofix) is on-going.

Utah TRAPS



This tool was developed with the Utah Climate Center, and provides accurate dates to implement pest management actions for approximately 70 locations in Utah. Growers can also monitor for frost, wind, soil moisture, and more at 22 of those locations.

On the TRAPs site, the new codling moth model is labelled, "Codling Moth-Fixed Biofix."

climate.usu.edu/traps

Table 2. Predicted emergence time [fixed-biofix from Jones et al. (2013), in degree days (DDs)] for apple and pear production regions of northern and southern Utah.

Location	Latitude	Elevation (ft)	Biofix (DDs)
Northern Utah			
Perry ¹	41.458	4,403	184
Roy ²	41.162	4,541	188
Kaysville ³	41.022	4,344	194
American Fork ⁴	40.392	4,613	202
Payson ⁴	40.036	4,874	205
Santaquin ⁴	39.991	4,844	207
West Mountain ⁴	40.028	4,688	209
Genola ⁴	40.005	4,715	209
Southern Utah			
Cedar City ⁵	37.707	5,627	251
New Harmony ⁶	37.482	5,197	265

¹located in Box Elder County

²located in Weber County

³located in Davis County

⁴located in Utah County

⁵located in Iron County

⁶located in Washington County

REFERENCES

Jones, V. P., M. Doerr, and J. F. Brunner. 2008. Is biofix necessary for predicting codling moth (Lepidoptera: Tortricidae) emergence in Washington state apple orchards? *Journal of Economic Entomology* 101 (5): 1651-1657.

Jones, V. P., R. Hilton, J. F. Brunner, W. J. Bently, D. G. Alston, B. Barrett, R. A. Van Steenwyk, L. A. Hull, J. F. Walgenbach, W. W. Coates, and T. J. Smith. 2013. Predicting the emergence of the codling moth, *Cydia pomonella* (Lepidoptera: Tortricidae), on a degree-day scale in North America. *Pest Management Science* DOI 10.1002/ps.3519 (www.wileyonlinelibrary.com).

Precautionary Statement: Utah State University Extension and its employees are not responsible for the use, misuse, or damage caused by application or misapplication of products or information mentioned in this document. All pesticides are labeled with active ingredients, directions for use, and hazards, and not all are registered for edible crops. "Restricted use" pesticides may only be applied by a licensed applicator. The pesticide applicator is legally responsible for proper use. USU makes no endorsement of the products listed herein.

Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran's status. USU's policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions. Utah State University employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran's status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USU-sponsored events and activities. This publication is issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Kenneth L. White, Vice President for Extension and Agriculture, Utah State University.