



# Parasitoid Wasps of the Invasive Brown Marmorated Stink Bug in Utah

Zachary R. Schumm • Mark Cody Holthouse • Yota Mizuno • Diane G. Alston • Lori R. Spears

## Do You Know?

- In 2012, brown marmorated stink bug (BMSB), an invasive insect pest from eastern Asia, was first detected in Utah in Salt Lake City. As of 2017, it is causing agricultural damage in northern Utah.
- There are few natural enemies of BMSB, allowing populations to increase easily.
- Parasitoid wasps that sting and kill stink bug eggs are the most promising control method. While there are native parasitoid wasps in Utah, none have been effective.
- Samurai wasp, a parasitoid native to eastern Asia, has proven to be effective at controlling BMSB populations. This wasp has been detected in some U.S. states.

The brown marmorated stink bug (BMSB, *Halyomorpha halys* Stål) is an invasive agricultural and nuisance pest native to eastern Asia. It was first confirmed in the U.S. in Allentown, PA, in 1996 and has since spread to 44 U.S. states, many of which have now experienced economic crop damage from this pest (Fig. 1). In Utah, BMSB is now established in five counties (Box Elder, Weber, Davis, Salt Lake and Utah), and has been detected in Cache and Kane counties. While crop damage to peach, apple, squash, and popcorn has been identified, it is currently causing mostly nuisance problems due to overwintering bugs on and inside human structures.




Fig. 1. The current distribution and status of BMSB in North America as of April 2019. For updates, see <http://www.stopbmsb.org/where-is-bmsb/>.

Adult BMSB are marbled brown and black, camouflaging well with woody vegetation. To separate this stink bug from native look-alikes, notice the characteristic white bands on their antennae. Native species do not have this feature. BMSB also has smooth shoulders and a black/white pattern on the edge of the abdomen (Fig. 2).

BMSB is a successful invasive for many reasons: it is polyphagous (feeds on many plant types), highly mobile, has few natural enemies, and adults have a tough exoskeleton that is covered in a waxy, water-repellent cuticle that helps protect them from pesticide applications. Biological control, through the use of egg parasitoids, is the most suitable option for long-term management of BMSB.




Fig. 2. A BMSB adult with quick identification characteristics. The white bands on dark antennae is the most helpful feature.

## GENERAL PARASITOID INFORMATION

There are at least two families of stink bug parasitoids in Utah, Eupelmidae and Scelionidae. These are small, typically black wasps that may be mistaken for small gnats or ants. They will fly in search of stink bug egg masses. Once they find the eggs, they will sting them, depositing one of their own eggs into the stink bug egg. The wasp egg will hatch, and the larva wasp will feed and develop within the stink bug egg, effectively killing the host. The adult wasp will emerge a couple of weeks later.

## PARASITOID WASP FAMILIES IN UTAH

### Eupelmidae – Generalist Egg Parasitoids

Eupelmids are small (3-5 mm), slender wasps that are generalist egg parasitoids. One genus, *Anastatus*, will parasitize BMSB, as well as other stink bugs and insects. They can resemble ants at first glance. Females often have a white band or white triangles on the wings. There are three species of *Anastatus* known to attack BMSB in Utah (Table 1). When




Fig 5. An *Anastatus* adult on BMSB eggs. Notice that the wasp is larger than an individual egg.

seen on a stink bug egg mass, a general rule is that these wasps are much larger than an individual egg (Fig. 5). Females are typically larger than males, and under direct light can exhibit brown, green, or blue iridescence. Males are typically all black, smaller (< 4 mm), and lack wing patterns, making males indistinguishable to species without a microscope.



Fig. 6. An adult *Anastatus* female. Adults typically measure 3-5 mm in length and resemble ants in appearance.

### Scelionidae – Specialist Egg Parasitoids

Scelionids are very small (1-2 mm), but often robust wasps that are specialists on different insect groups (Fig. 7). One genus within this family, *Trissolcus*, only stings stink bug eggs. The wasps attacking BMSB can only be identified to species by using microscopes,



Fig. 7. An adult *Trissolcus* female. Adults measure 1-2 mm with a robust body form.

The Eupelmids attacking BMSB are all generalist parasitoids, meaning that they sting the eggs of a wide variety of insects. Native parasitoids in this group are moderately successful at stinging and developing inside BMSB eggs, but are unlikely to control BMSB populations due to their generalist nature.

The second family, Scelionidae, includes some stink bug specialists, meaning they only sting stink bug eggs. Specialists are more promising as a control agent for BMSB. Although many of the native Scelionid species will sting BMSB eggs, some will not develop into an adult and emerge. Those that can complete development within BMSB eggs have the potential to be more effective control agents.

Stink bug eggs are usually bright in color (Fig. 3) and take 5 to 7 days to develop and hatch. Eggs will develop a triangular egg “burster” shortly before stink bugs emerge from the egg (Fig. 4). However, if parasitized by a wasp, the eggs will turn dark brown or black after about a week. As the wasps develop, the eggs will continue to darken until the adult wasps emerge about 14 days later (Fig. 3).



Fig. 3. Left: An adult BMSB with a freshly laid egg mass; Right top: A parasitoid stings a stink bug egg mass; Right bottom: eggs darken as parasitoids develop.

There is usually a skewed sex ratio in emerging wasps. In a typical stink bug egg mass that consists of 14-28 eggs, one to three wasps will be male, and the rest will be female. Male wasps will emerge first and wait for the females to emerge. Once mated, the females fly off in search of new egg masses to sting.



Fig. 4. A BMSB egg mass with triangular egg bursters. The nymphal stink bugs inside are close to hatching.



Table 1. Parasitoid wasp species found in Utah as of April 2019 from egg mass and yellow sticky card deployments. \*Based on results to-date from surveys in Utah.

Species Name	Family	Collection Method	Actual Size	Can Emerge from BMSB?
<i>Anastatus mirabilis</i>	Eupelmidae	BMSB Eggs	●	Yes*
<i>Anastatus persalli</i>	Eupelmidae	BMSB Eggs	●	Yes*
<i>Anastatus reduvii</i>	Eupelmidae	BMSB Eggs	●	Yes*
<i>Telenomus podisi</i>	Scelionidae	BMSB Eggs / Sticky Cards	-	No*
<i>Trissolcus erugatus</i>	Scelionidae	BMSB Eggs / Sticky Cards	●	No*
<i>Trissolcus euschisti</i>	Scelionidae	BMSB Eggs / Sticky Cards	●	Yes*
<i>Trissolcus hullensis</i>	Scelionidae	BMSB Eggs / Sticky Cards	●	Yes*
<i>Trissolcus parma</i>	Scelionidae	Sticky Cards	●	Unknown
<i>Trissolcus thyante</i>	Scelionidae	Sticky Cards	●	Unknown
<i>Trissolcus strabus</i>	Scelionidae	Sticky Cards	●	Unknown
<i>Trissolcus utahensis</i>	Scelionidae	BMSB Eggs / Sticky Cards	●	No*

as they are entirely black, small, and lack wing patterns or other characteristics to separate them with the naked eye. However, they can be generally identified in the field to family or genus using the tool that they are as small as or smaller than a stink bug egg (Fig. 8). There are at least two genera of stink bug parasitoids in the family Scelionidae in Utah (*Trissolcus* and *Telenomus*), with at least eight different species between these two genera (Table 1).




Fig 8. A *Trissolcus* female on a BMSB egg mass. Notice that the wasp is about the size of an egg.

Surveys are ongoing for other species of parasitoid wasps in Utah, particularly *Trissolcus japonicus* (samurai wasp) (Fig. 9). This wasp is native to eastern Asia, the native range of BMSB. In its native range, BMSB causes minimal economic damage, presumably due to effective biological control by the samurai wasp.

Samurai wasp was collected in China and is undergoing host range-testing in U.S. quarantine facilities to assess non-target effects for release in the U.S. However, samurai wasp has arrived on its own to the U.S. It has been found in 12 states as of January 2019 (Maryland, Pennsylvania, New Jersey, New York, Delaware, Oregon, Ohio, Virginia, West Virginia, Michigan, California, and Washington).

If samurai wasp is found in Utah, it can be reared and redistributed throughout the state to contribute to biological control of BMSB. Until samurai wasp is located in Utah, its release is prohibited.

Samurai wasp is more likely to be found in areas where BMSB are abundant (urban areas of Salt Lake and Utah valleys). However, it could be found in any location with established BMSB populations, making widespread surveys highly valuable.



Fig. 9. The samurai wasp (*Trissolcus japonicus*), a highly effective parasitoid against BMSB. It has been found in 12 U.S. states.

## SURVEYING FOR PARASITOIDS

Methods used to survey for stink bug parasitoids include physical placement of stink bug egg masses on host plants in the field, finding naturally-laid stink bug egg masses laid directly on host plants, and deployment of yellow sticky cards.

### Physical Egg Mass Placements:

Lab-reared stink bug eggs are attached to small squares of cardstock paper. These cards are then clipped to the underside of leaves on common hosts of stink bugs in Utah (fruit trees, vegetables, and ornamental trees such as northern catalpa [*Catalpa speciosa*]) (Fig. 10). Cards are left for 3 to 4 days to attract parasitoids. When collecting cards, parasitoids guarding the eggs are also collected to further assess their efficacy in stinging and developing in eggs.



Fig. 10. A BMSB egg mass clipped to a corn leaf to attract parasitoid wasps.

### Finding Naturally-laid Egg Masses:

Stink bug egg masses can be found on the underside of leaves, on the fruiting structures, and occasionally on the stems of host plants (Fig. 11). Just as with deployed egg masses, parasitoids guarding the egg masses are collected to assess efficacy in killing and developing in stink bug eggs.



Fig. 11. An *Anastatus* wasp on naturally-laid eggs.

### Yellow Sticky Cards:

Yellow sticky cards attract various insects. Cards are deployed for two weeks on the trunks and branches of ornamental and agricultural host plants, after which the wasps are removed from the card and identified (Fig. 12). When placed in areas with high parasitoid wasp diversity, cards are an effective tool for monitoring wasp diversity and density. While cards are effective at locating parasitoid wasps, information regarding the wasp's behavior on BMSB eggs, or their effectiveness at stinging, killing, and sustaining populations within them cannot be determined.



Fig. 12. Yellow sticky card hung on a tree to attract parasitoids.

### If You See a Parasitoid Wasp:

If you see a parasitoid on an egg mass, be mindful that it is beneficial for your garden/crops, as the wasps are potentially killing the stink bugs that would normally emerge and begin eating your plants. You can choose to either leave the stink bug eggs and parasitoids on the plant (preferred), or chase off the parasitoid, remove the stink bug eggs from the plant, and freeze or destroy them before they hatch.

It is possible that if the parasitoid wasps are left on the egg masses, then you may end up with more wasps, and consequently fewer stink bugs negatively impacting your plants. Parasitoids that are known to attack BMSB eggs in Utah can be seen in Figure 13.



Fig. 13. Select native parasitoid wasps stinging BMSB in Utah. From the top left: *Trissolcus erugatus*, *Trissolcus utahensis*, *Anastatus redivivus*, *Telenomus podisi*, and *Trissolcus euschisti*.

## REPORT PARASITOID WASPS

Reports of stink bug egg masses with parasitoid wasps on them are valuable to protect all Utah communities and property owners, and could help find new wasp species that may assist with biological control of BMSB. USU Extension has created a [Report an Invasive Pest in Utah website](https://www.usu.edu/extension/pests/bugs/stinkbug/) to report a finding, upload a photo, and find contact information for submitting a specimen for identification. Collect the egg mass and wasp in a bag or sealable container for specimen submission.



# Parasitoid Survey Locations and Abundances Using Stink Bug Egg Mass and Yellow Sticky Card Deployments




Fig. 14. A map of the current survey sites and parasitoid recovery locations in Utah through September, 2018.

## ADDITIONAL RESOURCES

Bergmann, E.J., Venugopal, P.D., Martinson, H.M., Raupp, M.J., and Shrewsbury, P.M. 2016. [Host plant use by the invasive \*Halyomorpha halys\* \(Stål\) on woody ornamental trees and shrubs](#). PLoS ONE 11(2): e0149975.

Herlihy M.V., Talamas E.J., Weber D.C. 2016. [Attack and success of native and exotic parasitoids on eggs of \*Halyomorpha halys\* in three Maryland habitats](#). PLoS ONE 11(3): e0150275.

Holthouse, M.C., Alston, D.G., Spears, L. R, and Petrizzo, E. 2017. [Brown marmorated stink bug \[\*Halyomorpha halys\* \(Stål\)\]](#) (8 pp). Utah State University Extension Fact Sheet Ent-144-17, Logan, UT.

Spears, L.R., Alston, D.G., and Murray, M. 2018. [Brown marmorated stink bug management for fruits and vegetables in Utah](#) (6 pp). Utah State University Extension Fact Sheet Ent-197-18, Logan, UT.

Spears, L.R., Davis, R., Alston, D.G., and Ramirez, R. 2016. [First detector guide to invasive insects: biology, identification, and monitoring](#) (53 pp). Utah State University Extension, Logan, UT.

Spears, L.R., Davis, R., and Ramirez, R. 2015. [Invasive insect look-alikes: mistaken insect identity](#) (6 pp). Utah State University Extension Fact Sheet Ent-175-15-PR, Logan, UT.

[Stopbmsb.org](#)

A comprehensive website on BMSB identification, management, and new research.

Talamas, E., Herlihy, M., Dieckhoff, C., Hoelmer, K., Buffington, M., Bon, M-C., and Weber, D. 2015. [Trissolcus japonicus \(Ashmead\) \(Hymenoptera, Scelionidae\) emerges in North America](#). Journal of Hymenoptera Research. 43. 119-128.

## PHOTO CREDITS

- |  |   |
|--|---|
| 1 Map courtesy of stopbmsb.org   | 8 Image courtesy of Elizabeth Beers, Washington State University            |
| 2 Image courtesy of Jeff Wildonger, USDA-ARS-BIIR;                     | 9 Image courtesy of Elijah J. Talamas, USDA ARS                             |
| 3 Images courtesy of Zach Schumm, Utah State University                | 10 Image courtesy of Zach Schumm, Utah State University                     |
| 4 Image courtesy of Rutgers New Jersey Agricultural Experiment Station | 11 Image courtesy of Cody Holthouse, Utah State University                  |
| 5 Image courtesy of Ashley Jones, University of Maryland               | 12 Image courtesy of Zach Schumm, Utah State University                     |
| 6 Image courtesy of John Rosenfeld, bugguide.net                       | 13 Images courtesy of Cody Holthouse and Zach Schumm, Utah State University |
| 7 Image courtesy of John Rosenfeld, bugguide.net                       | 14 Map courtesy of Zach Schumm, Utah State University                       |

## FUNDING

Funding for this publication was made possible by the National Institute of Food and Agriculture, U.S. Department of Agriculture, Specialty Crop Research Initiative (under award number 2016-51181-25409), U.S. Department of Agriculture, Utah Agricultural Experimental Station, Animal and Plant Health Inspection Service, Plant Protection and Quarantine (USDA-APHIS-PPQ) (under cooperative agreement number AP18PPQFO000C197), and USU Extension.

Precautionary Statement: Utah State University Extension and its employees are not responsible for the use, misuse, or damage caused by application or misapplication of products or information mentioned in this document. All pesticides are labeled with ingredients, instructions, and risks. The pesticide applicator is legally responsible for proper use. USU makes no endorsement of the products listed herein.

Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran's status. USU's policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions. USU employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran's status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USU-sponsored events and activities. This publication is issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Kenneth L. White, Vice President for Extension and Agriculture, USU.