

Fluoride: From Birth to Adolescence

November 2019

FN/Nutrition/ 2019-04pr

Ben Clark, Dietetics Student; Lydia Bangerter, Dietetics Student; Mateja R. Savoie-Roskos PhD, MPH, RD; Casey Coombs MS, RD; Carrie Durward PhD, RD

Tooth decay is one of the most common chronic diseases among children in the United States, and affects between 60-90% of school-aged children, particularly those in lower socioeconomic groups (Tham et al., 2015). In general, poor oral health can result in pain, infections, and tooth loss, which can negatively impact childhood quality of life (Chou, Cantor, Zakher, Mitchell, & Pappas, 2013). Fortunately, tooth decay is preventable! Fluoride is an essential mineral that helps strengthen tooth enamel and prevent cavities, especially in children (American Dental Association [ADA], 2014). In this fact sheet we will discuss the benefits of using fluoride, the most common concerns about fluoride intake in children, and recommendations from birth to adolescence.



Benefits of Fluoride

The most important benefit of fluoride is its association with improved dental health. More specifically, sufficient fluoride intake is associated with lower cavity rates in nearly every population, from infants to the elderly (ADA, 2014). Fluoride intake during the formative years of tooth development is linked to better overall tooth health as well as improved long-term tooth strength (ADA, 2014). The reduction in cavities will also reduce the need for fillings and tooth removal, which consequently reduces dental costs (Center for Disease Control [CDC], 2018). Studies show that preschool aged children who brushed their teeth with a standard fluoride toothpaste had a 24-29% reduced incidence of cavities (dos Santos, Nadanovsky, & de Oliveira, 2013). It is important to keep in mind that fluoride is a preventative measure and not a treatment for cavities, which is why it is important from a young age (Salas, Nascimento, Huysmans, & Demarco, 2015). If tooth decay or damage has occurred it will not be remedied with additional intake of fluoride (Salas et al., 2015).



Sources of Fluoride



Fluoride is found naturally in the environment in water and food sources, supplements, as well as in dental hygiene products (Kanduti, Sterbenk, & Artnik, 2016). Most freshwater has a small amount of naturally occurring fluoride with a concentration of 0.01-0.3 parts per million (ppm), which is not sufficient to provide protection against cavities. As of 2015, the U.S Public Health Service recommendation is that water has a fluoride concentration of 0.7 ppm or mg/L (U.S. Department of Health and

Human Services (USDHHS,2015). For this reason, many parts of the United States have introduced fluoride into the water supply in order to increase consumption; however, there are many areas in Utah that do not have fluoridated water despite the research and recommendations supporting the practice (CDC, 2018).

Recommendations



Current recommendations from the World Health Organization and the CDC are focused on providing fluoridated water and toothpaste to children (Petersen & Lennon, 2004; CDC, 2018). Fluoride needs vary with age and are different for children and adults (Wong et al., 2010). Table 1 includes the recommended use of fluoridated toothpaste for children of various ages. Before supplementing, it is beneficial to know the fluoride levels in the water you consume regularly. This can be done with home tests or, visit “My Water’s Fluoride” at.

https://nccd.cdc.gov/DOH_MWF/Default/Default.aspx on the CDC website. This can give you the best idea of the baseline amounts being ingested and whether additional supplementation is needed to reach the recommendations under the Affordable Care Act, insurance companies will cover preventative dental care for children, including fluoride treatments (HealthCare.gov, 2019). Talk with your child’s dentist or pediatrician if you have questions.



Table 1. Recommended Use of Fluoride Toothpaste.

Age	Recommendations
Birth-6 months	Not applicable
6 months – 2 years	Brush 2x daily with just water If advised by a dentist, brush with a smear of fluoridated toothpaste
2-6 years	Brush 2x daily with a pea sized amount of fluoridated toothpaste
6 years and over	Brush 2x daily with a peanut sized amount of fluoridated toothpaste

(Kanduti et al., 2016; Wright et al., 2014).

Table 2. Extended Information on Recommendations of Fluoride and Dental Health.

Age	Recommendations
Birth to 6 Months	<ul style="list-style-type: none">• Supplementation is not necessary, as breast milk contains enough fluoride to meet needs (Tham et al., 2015).• Studies show that children who are breastfed up to 12 months of age have a lower risk of developing cavities (Tham et al., 2015).• Teeth typically do not erupt until about 6 months of age, so brushing is not necessary.
6 Months to 2 Years Old	<ul style="list-style-type: none">• Children should visit a dentist every six months to ensure their teeth are healthy and to prevent dental complications (AAPD, 2019).• The ADA recommends that parents brush their children's teeth with only water until two years of age in order to reduce risk of fluorosis (Wright et al., 2014).• If parents choose to use toothpaste at this age, they should consult first with their dentist (ADA, 2014). The United States Department of Health and Human Services (USDHHS) recommends using a small smear of fluoridated toothpaste the size of a grain of rice (Wright et al., 2014).
2 Years to 6 Years Old	<ul style="list-style-type: none">• Children should be encouraged to brush their teeth twice a day with a pea sized amount of fluoridated toothpaste (Kanduti et al., 2016).• Teach children to spit out toothpaste instead of swallowing it while brushing so excess fluoride is not ingested (AAPD, 2019).• If living in an area where the water supply is not fluoridated, supplementation of fluoride is often advised by dentists and pediatricians (ADA, 2014).• Dentists can apply fluoride topically through varnishes, gels, and foam products (AAPD, 2019).• Fluoride products available for home use include toothpastes, mouth rinses, drops, gels, and pastes (AAPD, 2019).
6 Years up to 13 Years Old	<ul style="list-style-type: none">• As permanent teeth grow in, fluoride intake and overall oral health should be monitored carefully to ensure the development of strong permanent teeth (AAPD, 2019).• Children should be encouraged to brush their teeth twice a day with a peanut sized amount of fluoridated toothpaste to protect the tooth enamel (AAPD, 2019).• Children may begin to use fluoride rinses, which can increase fluoride exposure (Kanduti et al, 2016).
13 Years up to 18 Years Old	<ul style="list-style-type: none">• Throughout these years, teeth will become fully grown and require continued maintenance and protection.• Consistent use of fluoride toothpaste throughout adolescence and into adulthood will continue to protect teeth from decay and cavities (AAPD, 2019).

For more information on dental health in children and adolescents, talk with your pediatric dentist or visit the American Academy of Pediatric Dentistry and the American Dental Association websites.

Concerns with Excess Fluoride

The most common concern with excess fluoride is fluorosis, or visible damage and changes to the tooth enamel (USDHHS, 2015; Kanduti et al., 2017). Fluorosis occurs as a result of overexposure to fluoride in the first six years of life, when the process of tooth mineralization and enamel formation are occurring. Milder forms of fluorosis appear as lacy white marks on the teeth and discoloration, while more severe forms of fluorosis can result in pitting in the



teeth (USDHHS 2015). Fluorosis can result from misuse of dental hygiene products and from drinking excessive amounts of fluoridated water in addition to supplementation (Kanduti et al., 2017). Using fluoride treatments as directed by a dentist or physician can prevent these issues from occurring. Fluoride poisoning can also occur if dental products are misused or overconsumed. The probable toxic dose level is defined as being 5 mg of fluoride per kilogram of bodyweight consumed (Kanduti et al., 2017).

References

- American Academy of Pediatric Dentists (AAPD). (2019). Fluoride Therapy. Retrieved from <https://www.aapd.org/research/oral-health-policies--recommendations/fluoride-therapy/>
- American Dental Association (ADA). (2014). Fluoride toothpaste use for young children *The Journal of the American Dental Association*, 145(2), 190–191. doi.org/10.14219/jada.2013.47
- Center for Disease Control (CDC), Division of Oral Health.(2018). Community Water Fluoridation.. Retrieved from <https://www.cdc.gov/fluoridation/index.html>
- Chou, R., Cantor, A., Zakher, B., Mitchell, J. P., & Pappas, M. (2013). Preventing dental caries in children <5 years: Systematic review Updating USPSTF recommendation. *Pediatrics*, 132(2), 332–350. doi.org/10.1542/peds.2013-1469
- dos Santos, A. P. P., Nadanovsky, P., & de Oliveira, B. H. (2013). A systematic review and meta-analysis of the effects of fluoride toothpastes on the prevention of dental caries in the primary dentition of preschool children. *Community Dentistry and Oral Epidemiology*, 41(1), 1–12. doi.org/10.1111/j.1600-0528.2012.00708.x
- Healthcare.gov. (2019). Preventative care benefits for children. Retrieved from <https://www.healthcare.gov/preventive-care-children/>
- Kanduti, D., Sterbenk, P., & Artnik B. (2016). Fluoride: A review of use and effects on health. *Materia Socio Medica*, 28(2), 133. doi.org/10.5455/msm.2016.28.133-137
- Petersen, P. E., & Lennon, M. A. (2004). Effective use of fluorides for the prevention of dental caries in the 21st century: The WHO approach. *Community Dentistry and Oral Epidemiology*, 32(5), 319–321. doi.org/10.1111/j.1600-0528.2004.00175.x
- Tham, R., Bowatte, G., Dharmage, S., Tan, D., Lau, M., Dai, X., ... Lodge, C. (2015). Breastfeeding and the risk of dental caries: A systematic review and meta-analysis. *Acta Paediatrica*, 104, 62–84. <https://doi.org/10.1111/apa.13118>
- U.S. Department of Health and Human Services Federal Panel on Community Water Fluoridation (USDHHS). (2015). U.S. Public Health Service Recommendation for Fluoride Concentration in Drinking Water for the Prevention of Dental Caries. *Public Health Reports*, 130(4), 318–331. doi.org/10.1177/003335491513000408
- Wright, J. T., Hanson, N., Ristic, H., Whall, C. W., Estrich, C. G., & Zentz, R. R. (2014). Fluoride toothpaste efficacy and safety in children younger than 6 years. *The Journal of the American Dental Association*, 145(2), 182–189. doi.org/10.14219/jada.2013.37
- Wong, M., Glenny, A., Tsang, B., Lo, E., Worthington, H., & Marinho, V. (2010). Topical fluoride as a cause of dental fluorosis in children. *Cochrane Library*. doi.org/10.1002/14651858.CD007693.pub2

Acknowledgments

SNAP and FDIPIR State or local agencies, and their subrecipients, must post the following Nondiscrimination Statement: In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, sex, religious creed, disability, age, political beliefs, or reprisal or retaliation for prior civil rights activity in any program or activity conducted or funded by USDA. Persons with disabilities who require alternative means of communication for program information (e.g. Braille, large print, audiotape, American Sign Language, etc.), should contact the Agency (State or local) where they applied for benefits. Individuals who are deaf, hard of hearing or have speech disabilities may contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English. To file a program complaint of discrimination, complete the USDA Program Discrimination Complaint Form, (AD-3027) found online at:

http://www.ascr.usda.gov/complaint_filing_cust.html, and at any USDA office, or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture Office of the Assistant Secretary for Civil Rights 1400 Independence Avenue, SW Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov. This institution is an equal opportunity provider.

Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran's status. USU's policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions. Utah State University employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran's status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USU-sponsored events and activities. This publication is issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Kenneth L. White, Vice President for Extension and Agriculture, Utah State University.