Waterfall formation at a desert river-reservoir delta isolates endangered fishes

Running head: Waterfall formation isolates endangered fishes

Charles N. Cathcart^{1†}, Casey A. Pennock¹, Christopher A. Cheek^{2*}, Mark C. McKinstry³, Peter D. MacKinnon⁴, Mary M. Conner⁵, and Keith B. Gido¹

¹Division of Biology, Kansas State University, 104 Ackert Hall, Manhattan, Kansas 66506, USA

²Navajo Nation Department of Fish and Wildlife, Window Rock, Arizona 86515

³U.S. Bureau of Reclamation, Upper Colorado Regional Office, 125 South State Street, Room 8100, Salt Lake City, Utah 84138, USA

⁴Utah State University, Department of Watershed Sciences, 5210 Old Main Hill, Logan, UT 84322, USA

⁵Utah State University, Department of Wildland Resources, 5230 Old Main Hill, Logan, UT 84322, USA

[†]Present affiliation: Alaska Freshwater Fish Inventory, Alaska Department of Fish & Game, 333 Raspberry Road, Anchorage, AK, 99518, USA.

*Present affiliation: Forestry and Natural Resources, Purdue University, 715 West State Street, West Lafayette, Indiana, 47907, USA

Corresponding author: Charles N. Cathcart, cncathca@gmail.com

Acknowledgements

This project was funded by the Bureau of Reclamation with a grant (R11AC40022) to Kansas State University. Endangered fishes were sampled under the U.S. Fish and Wildlife Service permit TE067729.0. The Navajo Nation provided personnel, equipment, and permitting (Special permit # 1083). The National Park Service - Glen Canyon National Recreation Area permit (GLCA-2017-SCI-0003) allowed work downstream of the waterfall. Fieldwork, data assistance, or a combination of the two was provided by American Southwest Ichthyological Researchers LLC. (S. Platania, H. Brandenburg), Biowest Inc. (R. Rogers, B. Albrecht, R. Keggeries), Kansas State University (J. Brant, C. Ruffing, S. Hedden, L. Bruckerhoff), U.S. Fish and Wildlife Service (N. Franssen, E. Gilbert, S. Durst), Utah Division of Wildlife Resources (B. Hines, K. Creighton, Z. Ahrens, Z. Lowe, J. Dutrow), S. Fassbinder, and Utah State University. C. Ruffing and G. Cooper produced the map. E. Gilbert, S. Hedden, and two anonymous reviewers provided helpful comments on previous drafts. Any use of trade names does not imply endorsement by authors or the U.S. Government.

Abstract - Unforeseen interactions of dams and declining water availability have formed new obstacles to recovering endemic and endangered big-river fishes. During a recent trend of drying climate and declining reservoir water levels in the southwestern United States, a large waterfall has formed on two separate occasions (1989-1995 & 2001-present) in the transition zone between the San Juan River and Lake Powell reservoir because of deposited sediments. Because recovery plans for two large-bodied endangered fish species, razorback sucker (*Xyrauchen texanus*) and Colorado pikeminnow (*Ptychocheilus lucius*), include annual stockings in the San Juan River, this waterfall potentially blocks upstream movement of individuals that moved downstream from the river into the reservoir. To quantify the temporal variation in abundance of endangered fishes aggregating downstream of the waterfall and determine population demographics, we remotely monitored and sampled in spring 2015, 2016, and 2017 when these fish were thought to move upstream to spawn. Additionally, we used an open population model applied to tagged fish detected in 2017 to estimate population sizes. Colorado pikeminnow were so infrequently encountered (< 30 individuals) that population estimates were not performed. Razorback sucker captures from sampling (335) and detections from remote monitoring (943) showed high abundance across all three years. The razorback sucker population estimate for 2017 alone was 755 individuals and, relative to recent population estimates ranging from ~2000 to ~4000 individuals, suggests a substantial population exists seasonally downstream of this barrier. Barriers to fish movement in rivers above reservoirs are not unique, thus the formation of this waterfall exemplifies how water development and hydrology can interact to cause unforeseen changes to a riverscape.

Keywords: fragmentation, waterfall, endangered species, Colorado River Basin, river-reservoir inflow, razorback sucker, climate change

Introduction

Connectivity of freshwater systems and conservation of freshwater animals is challenged worldwide by increasing drought and pervasive water development, often in the form of large dams and excessive water use (Ruhi et al. 2016). Dams and reservoirs disrupt the continuity of rivers (Stanford & Ward 2001) where they create abrupt shifts in physical and biological properties at the junctions of rivers and reservoirs (Galay 1983; Poff et al. 1997; Sabo et al. 2012). Once impaired, fragmented rivers often experience declines or extinctions of fishes disconnected from habitats necessary for the fulfillment of life histories (Minckley & Deacon 1991; Moyle 1995). Ultimately, these disconnections and alterations have contributed to the listing of many fishes or populations under the Endangered Species Act, including a high percentage of native fishes from the Colorado River Basin (Minckley & Deacon 1991; Osmundson 2011). Despite the intrinsic value of native fish and cost of recovery, conservation programs must often consider barriers (especially dams or diversions) as permanent structures to the landscape because of their economic value and importance to water security (Propst & Gido 2004; Coutant & Whitney 2006; Lackey 2013).

Research perspectives have primarily focused on downstream effects of dams, with limited attention paid to changes occurring upstream of impoundments in both fish populations and stream function (Falke & Gido 2006). Inundated lotic habitat upstream of dams can reduce habitat availability, restrict migration, and diminish population viability for riverine species (Osmundson 2011; Hudman & Gido 2013). An upstream perspective may be particularly useful to understand the importance of the river-reservoir interface for both lentic and lotic adapted species (Minckley & Deacon 1991; Stanford & Ward 2001; Birnie-Gauvin et al. 2017). In addition, dynamic reservoir volume alters geomorphological processes structuring delta formation and location (Galay 1983; Stevens et al. 2001; Johnson 2002). Specifically, as reservoir levels recede from decreasing basin water availability or seasonal dam operations, vegetation sequesters sediments in the inflow area (raising elevation of the river channel) slowing inflow and depositing sediment on higher surfaces (Pasternack & Brush 1998; Johnson 2002). In the Colorado River Basin, receding reservoir levels have exposed river-reservoir deltas, altering river channels in alluvial sediments.

Lake Powell, created in 1963, is the second largest reservoir in the US, covering 400-660 km² (1.5-3.0 million hectare meters of storage) and includes the historical confluence of the San Juan and Colorado rivers (Figure 1). Combined sediment deposition and water level declines in Lake Powell have resulted in a geomorphic barrier at the San Juan River inflow to Lake Powell, Utah between 1989 and the present. Lake Powell reservoir experienced dynamic inflows since reaching capacity in 1980, which subsequently led to delta formation and the eventual emergence of waterfall barriers on the San Juan River (Figure 2). These barriers to fish movement, which first appeared as late as 1989, were described by Ryden and Ahlm (1996) as being > 10 m tall depending on river flows. The reservoir then experienced a period of greater storage from higher inflows throughout the mid-1990s, inundating the waterfall by 1995. After further water level

recession in the late 1990s, the river channel again shifted through the newly formed delta and a new waterfall formed in 2001 approximately 3 km downstream from the prior waterfall (Figure 3). This process, referred to as superimposition, involves the river cutting through new deposited sediments as reservoir levels recede, thus creating a new channel. The current waterfall is > 6 m tall and is a complete barrier to upstream fish movement in an area referred to as Piute Farms, UT (Figure 4). Since emerging in 2001, the current waterfall has only been inundated (thus passable) once, in 2011 for two weeks in late-July and mid-August (Durst & Francis 2016).

Two intensively managed endangered species are likely affected by the emergent waterfall. Colorado pikeminnow (*Ptychocheilus lucius*) and razorback sucker (*Xyrauchen texanus*) are large-bodied (> 1 m long), long lived (> 30 years old), highly fecund (mature individuals regularly have > 60,000 eggs), and migratory fishes endemic to large river habitats in the Colorado River Basin that typically spawn in late-spring to mid-summer after snowmelt runoff (Hamman 1985; Hamman 1986). Colorado pikeminnow have a non-augmented wild population in the Upper Colorado River and a stocked population in the San Juan River and are highly migratory in both systems (Osmundson 2011; Durst & Franssen 2014). Besides rivers, razorback sucker inhabit (and spawn in) all major Colorado River Basin reservoirs (Mead, Mohave, Havasu, and Powell). Razorback sucker often spawn on the ascending limb of the hydrograph from mid-March through June at water temperatures between 9-17°C (Tyus & Karp 1990). Successful recruitment to adulthood has only been documented in Lake Mead and we do not understand how reservoir-dwelling razorback sucker life histories may interact with inflowing rivers (Albrecht et al. 2010; Marsh et al. 2015; Albrecht et al. 2017). Lake Powell is both a movement corridor connecting the upper Colorado River and San Juan River basins and a habitat for razorback sucker that are known to make substantial downstream movements after stocking or during larval drift (Zelasko et al. 2010; Durst & Francis 2016; Albrecht et al. 2017). Current management for both species involves stocking (Zelasko et al. 2010), mimicking natural flow regimes (Propst & Gido 2004), and removing nonnative fishes (Franssen et al. 2014).

Over 140,000 razorback sucker and over 50,000 Colorado pikeminnow have been implanted with passive integrated transponder (PIT) tags in the San Juan River basin during stocking or on-river tagging events between 2000 and 2017 (Figure 1). In the Upper Colorado River Basin upstream of Lake Powell (e.g., Colorado, Green, and Gunnison rivers), ~424,000 razorback sucker and ~50,000 Colorado pikeminnow have been PIT tagged and could travel through the reservoir to the waterfall. With few exceptions, razorback sucker are stocked in these rivers with a PIT tag at ~ 300 mm TL. Colorado pikeminnow are stocked in the San Juan River as juveniles (<100 mm TL) and are PIT-tagged at first capture. Intense sampling of tagged endangered fishes in the San Juan River upstream of the waterfall within and across years has allowed population estimates of endangered fishes in the river (USFWS 2017) but does not account for fishes that move downstream to the reservoir. Our main objectives were to measure sex-ratios, quantify temporal patterns of abundance, and estimate population sizes of Colorado pikeminnow and razorback sucker downstream of the waterfall. This research shows how

unforeseen fragmentation alters endangered fish population connectivity and, ultimately, their recovery.

Methods

Fish sampling

Because of limited historical sampling downstream of the Piute Farms waterfall, we performed pilot sampling in 2015 to assess the occurrence of endangered fishes. After confirming the presence of endangered fish, more rigorous sampling in the localized area (0-500 m downstream of the waterfall) was conducted during spring of 2016 (March and April) and 2017 (February and March) with raft-mounted electrofishing. Amount of habitat and sampling effort (two 15 minute “passes”) were similar across days, although total days sampled varied across years (6-13 d). Endangered species were identified, measured for total length (TL), and sexed when possible through observation of sexually dimorphic traits (i.e., gamete expression, tubercle presence, razorback sucker anal fin shape) and were scanned with a PIT tag reader for the presence of prior tags. If a tag was absent, we implanted the fish with a PIT tag (Biomark, Boise, Idaho, 12-mm full-duplex, 134.2 kHz). All individual fish captured in 2015, 2016, and a subset in 2017 were translocated upstream of the waterfall barrier as a conservation action to assist migration and promote spawning.

Temporal variation in abundance

To detect PIT tagged fishes, we deployed a circular (1 m diameter) submersible PIT tag antenna (Biomark, Boise, Idaho) from March 21 through July 6, 2015 (107 d), March 2 through April 7, 2016 (36 d), and Feb 12 to Jun 3, 2017 (111 d). The antenna was deployed in an eddy approximately 10 m downstream of the waterfall on the right bank, over sand and bedrock substrates in water depths from 70 to 160 cm. The antenna typically detected tags within 0.5 m. Detected individuals were identified by relating them to a PIT tag database compiled by the San Juan River and Upper Colorado River recovery programs (STReaMS 2017).

To illustrate environmental cues commonly correlated with fish spawning migrations, we show the relationship of tag detections with mean daily discharge ($\text{m}^3 \text{s}^{-1}$) and mean daily water temperature ($^{\circ}\text{C}$) from the USGS gauge near Bluff, UT (gauge number 9375000), approximately 85 km upstream from the waterfall.

Population size estimates

Extremely low detections of Colorado pikeminnow downstream of the waterfall prevented their population estimation but we estimated population size of razorback suckers in 2017. Capture data from 2015 was inadequate to estimate population size and the sampling period changed between 2016 and 2017; it was 02 March-07 April in 2016 and 12 February-03 June in 2017. We lengthened the sampling period (physical capture plus antenna resight period)

in 2017 to increase sample sizes; sampling period was 3.6 months in 2017 compared to 1.2 months in 2016. The longer sampling period yielded a greater number of unique fish captured, which was 32% higher for 2017 compared to 2016. Thus, we only estimated population size for 2017 as it reasonably encompassed an entire spawning season and had adequate sample size. Translocated fish were not used in the open population size estimates because they could not be recaptured. Due to the long detection period (12 February–03 June), we tested the assumption of population closure for the antenna detection data using Program CloseTest (Stanley & Burnham 1999). This test indicated the assumption of closure was not met. Fish were entering and leaving the study area during the detection period; thus, we estimated population size using POPAN (Schwarz & Arnason 1996), an open population model implemented in Program MARK (Cooch and White 2016). POPAN is a Jolly-Seber model and assumes equal catchability (or detection) among individuals, which means we did not expect there to be a behavioral response to being detected by the antenna.

For the POPAN model, a previously PIT-tagged fish was considered ‘unmarked’ until it was first detected by the antenna, after which it was considered a marked fish. Marked fish could be detected by the antenna continuously during the sampling period. Based on the proportions of unique fish detected, we grouped the data into 4 periods; 12 February–15 March, 16 March–31 March, 01 April–15 April, and 16 April–03 June (the proportions of unique fish detected for each occasion were 0.24, 0.30, 0.23, and 0.23). To account for differences in period length, we used unequal time intervals in Program MARK. The cumulative number of tags detected across the time periods used in the model indicated longer antenna deployment did not result in greater numbers of unique tags detected (Figure S1). We constructed a set of models with capture probability (p), apparent survival probability, which in this situation is the probability of leaving the waterfall area (ϕ), and probability of initial entrance to the waterfall area (p_{ent}) modeled as constant across the 4 sampling periods and variable from period to period. We constructed 8 initial models for all possible combinations of these 3 parameters. We used the “gross” population size from POPAN (Schwarz & Arnason 1996), which is the number of PIT tagged razorback sucker using the waterfall area over the entire study period and includes fish who arrived and departed between occasions. We added the count of translocated fish to the model-averaged estimated population size from POPAN to estimate a minimum total population size of razorback sucker using the waterfall area during the sampling period. This estimate allowed for comparison to razorback sucker population size in the San Juan River upstream of the waterfall (USFWS 2017).

Results

Fish sampling

Below the waterfall, we captured 167 razorback sucker in 2016 and 183 in 2017 (Table 1). Razorback sucker ranged from 403 to 618 mm TL with a minimum weight of 550 g and a maximum of 2800 g. In 2016, about 10% of females and 77% of males that were handled were

freely expressing gametes. Sampling was performed earlier in 2017 and ripe fish were rare. Twenty-four Colorado pikeminnow were captured, and most were sub-adults except for a 571 mm TL fish in 2016.

Temporal variation in abundance

Over three years, we detected 967 unique endangered fish downstream of the waterfall (Table 1). Razorback sucker made up a large proportion (98%) of detected fishes across all years. The majority of detected (and captured) fish were either stocked or tagged in the San Juan River upstream of the waterfall, but several razorback sucker came from the Upper Colorado River Basin (Figure 1), which involves a minimum of 220 km to traverse Lake Powell. The PIT antenna ran continuously during study periods in all three years, except for five days (May 28 to June 3) in 2015 and again in 2017, when ~ 1 m of sediment buried the operating antenna for six consecutive days in late February.

Some fish were detected in multiple years for both species. Of razorback sucker detected in 2015, 51% ($n = 255$) were also detected in 2016 and 64% ($n = 302$) of fish detected in 2016 were then detected in 2017. Eighteen percent (167) of razorback sucker were detected in all three years. Concomitant with their relatively low detection numbers, few Colorado pikeminnow were detected multiple years. One fish was detected in all three years, one fish each was detected in both 2015-2016 and 2016-2017, and a single individual detected in 2015 was detected in 2017.

Water temperatures and flows showed similar patterns across all three years. Water temperatures during antenna deployments included observed spawning temperatures for razorback sucker (Figure 5). Generally, patterns of unique daily razorback sucker detections were similar across all three years. Each year, daily detections were variable but higher earlier in the study period and declined over time with increasing water temperature and river discharge. Peak razorback sucker abundances at the waterfall occurred while water temperatures were below 16°C until warming in mid-April when razorback sucker abundance decreased.

Population estimates

In 2017 we captured and/or detected 689 unique individual razorback sucker. Of these, 183 were physically captured (27%) and 506 were PIT tagged but only detected by the antenna (73%). Of the 183 fish physically captured, 34 did not have a PIT tag (19%). All physically captured fish were moved upstream of the waterfall area. The eight candidate models estimating razorback sucker population using only fish only detected by the antenna were ranked by Akaike's information criterion. The top POPAN model included $\phi(.)$ $p(t)$ $pent(t)$ and had a model weight (w_i) of 0.81 (Table S1). Detection probabilities were high, ranging from 0.64 to 0.91. The model-averaged estimated population size for 2017 was 572 (SE = 11.7, 95% CI = 549–595). Adding the minimum count of physically captured fish indicated that at least 755 razorback suckers used the waterfall area in 2017.

Discussion

Although we expected to capture endangered fish downstream of the waterfall based on past occurrence records, the large number of razorback sucker we sampled was surprising and showed that a substantial proportion of the fish stocked in the river moved downstream to the reservoir. Using PIT antennas continuously in a novel, albeit discrete and fine-scaled, location within the Colorado River Basin further illustrated how remote sensing can more accurately measure populations compared to spatially continuous yet temporally discrete active sampling events (Webber and Beers 2014). That these fish migrated back upstream and aggregated below the waterfall in spring enhanced our ability to detect individuals and then accurately represent and estimate the population of razorback suckers here. USFWS (2017) population estimates from 112.5 km of the upper river in 2015 ranged from 2296-4073 fish compared to our 2017 population estimate of 755 fish. Thus, the proportion of the San Juan River population using habitat downstream of the waterfall was between 19 and 33%. Given that 5,800 adult razorback sucker in the San Juan River are necessary for downlisting them from endangered to threatened, the cumulative populations in habitats upstream and downstream of the waterfall represent significant progress toward reaching that recovery criterion. Barring rare waterfall inundation during high river flow events synchronized with elevated reservoir pool such as late-summer 2011 (that wouldn't assist spring spawning migrations anyway), this > 6 m tall waterfall is a barrier to all fishes attempting to swim upstream (see Meixler et al. 2009). Although it seems limited, quantifying spawning habitat (i.e., confluences of washes, areas with coarse substrates) in the ~25 km reach between Lake Powell and the waterfall would be a considerable first step toward understanding the potential of this river-reservoir transition area to support the life history of razorback sucker isolated from the upper San Juan River.

While the waterfall certainly impedes connectivity of adult fishes, recruitment of early life stages upstream of the waterfall could also be compromised by this fragmentation. The abundance of mature, gamete-spewing razorback sucker repeatedly detected and captured coincident with observed spawning temperatures implies the waterfall blocks annual spawning migrations into the upper San Juan River. Historical and contemporary monitoring indicates the presence of young-of-the-year (larval and transformed juvenile) razorback sucker and Colorado pikeminnow just upstream of the waterfall as well as in the inflow area where the San Juan River transitions into Lake Powell (Platania et al. 1991; Pennock unpublished data). Larval fish could accumulate in the inflow following drift from hatching locations upstream in the San Juan River and over the waterfall. Flow regulation and invasive Russian olive (*Elaeagnus angustifolia*) have channelized the river, thereby reducing larvae-retaining habitats (inundated floodplains and backwaters) and increasing larval drift distance (e.g., Robinson et al. 1998). Generational losses to upstream reaches from isolated downstream populations could also occur when upstream migrations cannot occur to offset larval drift (e.g., Perkin & Gido 2011).

Ryden & Ahlm (1996) suggested the first waterfall in the San Juan River disrupted Colorado pikeminnow migrations. Our sampling from late winter to early summer may have

missed movements to or over the waterfall that could occur at other times of year. Given their tendency for long-distance migrations as adults (Tyus & McAda 1984), downstream winter migrations as sub-adults (Durst & Franssen 2014), and the fact they are stocked at small sizes without PIT tags, the Piute Farms waterfall presents a major challenge to Colorado pikeminnow recovery in the San Juan River if downstream migrating fish swim too far and become “trapped” below the waterfall.

The discontinuity of a desert river caused by an emergent waterfall in a reach between two large dams is likely reconcilable. Connecting habitats through fish passage (including barrier removal, bypass, or capture and translocate) could allow hundreds of endangered fish to move seasonally (*sensu* Pess et al. 2014). Fish passage systems mitigate barriers to migratory fish if designed correctly, but they can also negatively interact with some species, including suckers, by preventing or delaying movements (McLaughlin et al. 2013; Hatry et al. 2016). Regardless, total functional connectivity of the river is not necessarily preferred by recovery programs that devote substantial resources to removing nonnative fish that are considered a primary threat to endangered Colorado River Basin fishes (Minckley & Deacon 1991; Franssen et al. 2014). In fact, the Piute Farms waterfall also blocks upstream movement of nonnative predatory fishes such as striped bass (*Morone saxatilis*) and walleye (*Sander vitreus*). Thus, alternative methods (e.g., selective fish passage such as translocation) would maintain downstream isolation of nonnative fishes (Rahel 2013). Lake Powell requires >85% fullness to inundate the waterfall, suggesting this will likely remain a barrier to fish movement for the foreseeable future (Bureau of Reclamation, *unpublished data*). If connectivity is desired, our study pinpoints effective times to manage for passage, especially for razorback sucker.

The barrier-forming geomorphological processes described here (and in Ryden & Ahlm 1996) are not unique to the San Juan River and are currently creating fragmentation issues upstream of another large southwestern American reservoir. A volatile large rapid formed via interactions of reservoir volume and superimposition processes in the mid-2000s at the Colorado River inflow to Lake Mead where the river exits the Grand Canyon at Pearce Ferry (Martin & Whitis 2013). Formation of this rapid created such a hazard to river runners the National Park Service constructed a multi-million-dollar road and takeout area upstream of the rapid to allow users to exit safely (Video S2). Pearce Ferry Rapid is younger than Piute Farms waterfall but may be approaching a similar result: a barrier to endangered fish movements between Lake Mead and the Grand Canyon. The importance of connectivity between Lake Mead and Grand Canyon to razorback sucker is unknown and should be considered as Pearce Ferry Rapid develops.

The effects (and threat) of fragmentation on freshwater fishes are well documented and include community structure changes, population reduction, enhanced negative species interactions, and species extirpation upstream and downstream of barriers (Sanches et al. 2006; Perkin & Gido 2011; Guy et al. 2015; Gido et al. 2016). Despite the acknowledgment of fragmentation effects in conceptual models of riverine function (e.g., Stanford & Ward 2001)

and negative interactions of reservoirs with large river fish recruitment (Guy et al. 2015), current models treat reservoirs separately from the rivers they impound, which could explain the limited number of studies assessing upstream effects of reservoirs. Studies on fish distributions between or within reservoir and riverine habitats treat reservoirs as strictly lentic habitats and often consider these artificial systems as barriers themselves (Taylor et al., 2001; Falke & Gido 2006; Buckmeier et al., 2014). In reality, there is not an abrupt change from riverine to reservoir environments but more gradual change as one moves through the riverine, transition, and lacustrine zones within a reservoir (Thornton et al. 1990). This gradient of ecosystem novelty (e.g., Gandy & Rehage 2017) along the river-reservoir continuum could provide productive habitats (e.g., floodplain connectivity) no longer seen in upstream portions of regulated rivers (Volke et al. 2015) and benefit fish (e.g., razorback sucker) able to utilize the lentic-lotic interface (Gido et al. 2002; Da Silva et al. 2015). However, the consequences of being isolated in these habitats are largely unknown. These contemporary barrier formation events illustrate how fragmentation and isolation can metastasize in alluvial rivers when delta formation processes interact with increased water use, historical fragmentation, and natural topography. Depending on when, where, and what these emergent features can affect (such as fish or public safety), awareness and action can assist resource managers in adapting to them.

Literature Cited

- Albrecht, B. A., Holden, P. B., Kegerries, R. B., & Golden, M. E. (2010). Razorback sucker recruitment in Lake Mead, Nevada-Arizona, why here? *Lake and Reservoir Management*, 26, 336-344. <https://doi.org/10.1080/07438141.2010.511966>
- Albrecht, B.A., Mohn, H.E, Kegerries, R., McKinstry, M.C., Rogers, R., Francis, T., Hines, B., Stolberg, J., Ryden, D., Elverud, D. Schleicher, B., Creighton, K., Healy, B., & Senger, B. 2017. Use of inflow areas in two Colorado River basin reservoirs by endangered razorback sucker (*Xyrauchen texanus*). *Western North American Naturalist*.
- Birner-Gauvin, K., Aarestrup, K., Riis, T. M. O., Jepsen, N., & Koed, A. (2017). Shining a light on the loss of rheophilic fish habitat in lowland rivers as a forgotten consequence of barriers, and its implications for management. *Aquatic Conservation Marine and Freshwater Ecosystems*, 27, 1345-1349. <https://doi.org/10.1002/aqc.2795>
- Buckmeier, D. L., Smith, N. G., Fleming, B. P., Bodine, K. A. (2014). Intra-annual variation in river-reservoir interface fish assemblages: implications for fish conservation and management in regulated rivers. *River Research and Applications*, 30, 780-790. <https://doi.org/10.1002/rra.2667>
- Cooch, E. G., & White, G. C. (2016). Program MARK: "A gentle introduction", 14th Edition. <http://www.phidot.org/software/mark/docs/book/>.

- Coutant, C. C., & Whitney, R. R. (2006). Hydroelectric system development: effects on juvenile and adult migration. In R. N. Williams (Eds.), *Return to the River: restoring salmon to the Columbia River* (pp. 249-324). Elsevier Academic Press, San Diego, California.
- Da Silva, P. S., Makrakis, M. C., Miranda, L. E., Makrakis, S., Assumpção L., Paula, S., Dias, J. H. P., & Marques, H. (2015). Importance of reservoir tributaries to spawning of migratory fish in the upper Paraná River. *River Research and Applications*, 31, 313-322. <https://doi.org/10.1002/rra.2755>
- Durst, S. L., & Frannsen, N. R. (2014). Movement and Growth of Juvenile Colorado Pikeminnows in the San Juan River, Colorado, New Mexico, and Utah. *Transactions of the American Fisheries Society*, 143, 519-527. <https://doi.org/10.1080/00028487.2013.869258>
- Durst, S. L., & Francis, T. A. (2016). Razorback Sucker trans-basin movement through Lake Powell, Utah. *The Southwestern Naturalist*, 60, 60-63. <https://doi.org/10.1894/0038-4909-61.1.60>
- Falke, J.A., & Gido, K. B. (2006). Effects of reservoir connectivity on stream fish assemblages in the Great Plains. *Canadian Journal of Fisheries and Aquatic Sciences*, 63, 480-493. <https://doi.org/10.1139/f05-233>
- Franssen, N. R., Davis, J. E., Ryden, D. W., & Gido, K. B. (2014). Fish community responses to mechanical removal of nonnative fishes in a large Southwestern river. *Fisheries*, 39, 352-363. <https://doi.org/10.1080/03632415.2014.924409>
- Gandy, D. A., & Rehage, J. S. (2017). Examining gradients in ecosystem novelty: Fish assemblage structure in an invaded Everglades canal system. *Ecosphere*, 8, e01634. <https://doi.org/10.1002/ecs2.1634>
- Gido, K. B., Hargrave, C.W., Matthews, W.J., Schnell, G.D., Pogue, D.W., & Sewell, G. (2002). Structure of littoral-zone fish communities in relation to habitat, physical, and chemical gradients in a southern reservoir. *Environmental Biology of Fishes*, 63, 253-263. <https://doi.org/10.1023/A:1014359311188>
- Gido, K. B., Whitney, J. E., Perkin, J. S., & Turner, T. F. (2016). Fragmentation, connectivity and fish species persistence in freshwater ecosystems. In G. P. Closs, M. Krkosek, & J. D. Olden (Eds.) *Conservation of freshwater fishes* (pp. 292-323). Cambridge University Press.
- Greathouse, E. A., Pringle, C. M., McDowell, W. H., & Holmquist, J. G. (2006). Indirect upstream effects of dams: consequences of migratory consumer extirpation in Puerto Rico. *Ecological Applications*, 16, 339–352. <https://doi.org/10.1890/05-0243>

- Guy, C.S., Treanor, H.B. Kappenman, K.M., Scholl, E.A. Ilgen, J.E., & Webb, M.A.H. (2015). Broadening the regulated-river management paradigm: a case history of the forgotten dead zone hindering pallid sturgeon recovery. *Fisheries*, 40, 6-14. <https://doi.org/10.1080/03632415.2014.987236>
- Hatry, C., Thiem, J. D., Hatin, D., Dumont, P., Smokorowski, K. E., & Cooke, S. J. (2016). Fishway approach behaviour and passage of three redhorse species (*Moxostoma anisurum*, *M. carinatum*, and *M. macrolepidotum*) in the Richelieu River, Quebec. *Environmental Biology of Fishes*, 99, 249-263. <https://doi.org/10.1007/s10641-016-0471-3>
- Hamman, R. L. (1985). Induced spawning of hatchery-reared razorback sucker. *The Progressive Fish Culturist*, 47, 187-189. [https://doi.org/10.1577/1548-8640\(1985\)47<187:ISOHRS>2.0.CO;2](https://doi.org/10.1577/1548-8640(1985)47<187:ISOHRS>2.0.CO;2)
- Hudman, S. P. & Gido, K. B. (2013). Multi-scale effects of impoundments on genetic structure of creek chub (*Semotilus atromaculatus*) in the Kansas River Basin. *Freshwater Biology*, 58, 441-453. <https://doi.org/10.1111/fwb.12079>
- Johnson, W. C. (2002). Riparian vegetation diversity along regulated rivers: contribution of novel and relict habitats. *Freshwater Biology*, 47, 749-759. <https://doi.org/10.1046/j.1365-2427.2002.00910.x>
- Lackey, B. T. (2013). Saving wild salmon: a 165-year policy conundrum. Dubach Workshop: Science and Scientists in the Contemporary Policy Process. Oregon State University, October 3-4, Portland, Oregon.
- Marsh, P. C., Dowling, T. E., Kesner, B. R., Turner, T. F., & Minckley, W. L. (2015). Conservation to stem imminent extinction: the fight to save Razorback Sucker *Xyrauchen texanus* in Lake Mohave and its implications for species recovery. *Copeia*, 2015, 141–156. <https://doi.org/10.1643/OT-14-030>
- Martin, T. & Whitis, D. (2013). Guide to the Colorado River in the Grand Canyon: Lees Ferry to South Cove, 5th edition. Vishnu Temple Press, Flagstaff, AZ.
- McLaughlin, R. L., Smyth, E. R. B., Casto-Santos, T., Jones, M. L., Koops, M. A., Pratt, T. C., & Vélez-Espino, L.-A. (2013). Unintended consequences and trade-offs of fish passage. *Fish and Fisheries*, 14, 580-604. <https://doi.org/10.1111/faf.12003>
- Meixler, M.S., Bain, M.B., and Walter, M.T. (2009). Predicting barrier passage and habitat suitability for migratory fish species. *Ecological Modelling*, 220, 2782-2791. <https://doi:10.1016/j.ecolmodel.2009.07.014>
- Minckley, W. L., & Deacon, J. E. (1991). Battle against extinction: native fish management in the American West. University of Arizona Press, Tucson.

- Moyle, P. B. (1995). Conservation of native fresh-water fishes in the Mediterranean-type climate of California, USA – a review. *Biological Conservation*, 72, 271-279. [https://doi.org/10.1016/0006-3207\(94\)00089-9](https://doi.org/10.1016/0006-3207(94)00089-9)
- Osmundson, D. B. (2011). Thermal regime suitability: assessment of upstream range restoration potential for Colorado Pikeminnow, a warmwater endangered fish. *River Research and Applications*, 27, 706-722. <https://doi.org/10.1002/rra.1387>
- Perkin, J. S., & Gido, K. B. (2011). Stream fragmentation thresholds for a reproductive guild of Great Plains fishes. *Fisheries*, 36, 371-383. <https://doi.org/10.1080/03632415.2011.597666>
- Pess, G. R., Quinn, T. P., Gephard, S. R., & Saunders, R. (2014). Re-colonization of Atlantic and Pacific rivers by anadromous fishes: linkages between life history and the benefits of barrier removal. *Reviews in Fish Biology and Fisheries*, 24, 881-900. <https://doi.org/10.1007/s11160-013-9339-1>
- Pasternack G. B. & Brush G. S. (1998). Sedimentation cycles in a river-mouth tidal freshwater march. *Estuaries*, 21, 407-415. <https://doi.org/10.2307/1352839>
- Platania, S. P., Bestgen, K. R., Moretti, M. M., Brooks, J. E., & Propst, D. L. (1991). Status of Colorado squawfish and razorback sucker in the San Juan River, Colorado, New Mexico, and Utah. *The Southwestern Naturalist*, 36, 147-150. <https://doi.org/10.2307/3672136>
- Poff, N. L., Allan, J. D., Bain, M. B., Karr, J. R., Presegaard, J. L., Richter, B. D., Sparks, R. E., & Stromberg, J. C. (1997). The natural flow regime: a paradigm for river conservation and restoration. *BioScience*, 47, 769-784. <https://doi.org/10.2307/1313099>
- Pringle, C. M. (1997). Exploring how disturbance is transmitted upstream: going against the flow. *Journal of the North American Benthological Society*, 16, 425–438. <https://doi.org/10.2307/1468028>
- Propst, D. L., & Gido, K. B. (2004). Responses of native and nonnative fishes to natural flow regime mimicry in the San Juan River. *Transactions of the American Fisheries Society*, 133, 922–931. <https://doi.org/10.1577/T03-057.1>
- Rahel, F. J. (2013). Intentional fragmentation as a management strategy in aquatic systems. *BioScience*, 63, 362-372. <https://doi.org/10.1525/bio.2013.63.5.9>
- Robinson A. T., Clarkson, R. W., & Forrest, R. E. (1998). Dispersal of larval fishes in a regulated river tributary. *Transactions of the American Fisheries Society*, 127, 772-786, [https://doi.org/10.1577/1548-8659\(1998\)127.0.CO;2](https://doi.org/10.1577/1548-8659(1998)127.0.CO;2)
- Ruhi, A., Olden, J.D. & Sabo, J. L. (2016) Declining streamflow induces collapse and replacement of native fish in the American Southwest. *Frontiers in Ecology and the Environment*, 14, 465-472. <https://doi.org/10.1002/fee.1424>

- Ryden, D. W., & Ahlm, L. A. (1996). Observations on the distribution and movements of Colorado squawfish, *Ptychocheilus lucius*, in the San Juan River, New Mexico, Colorado, and Utah. *The Southwestern Naturalist*, 41, 161-168.
www.jstor.org/stable/30055101
- Sabo, J. L., Bestgen, K., Graf, W., Sinha, T., & Wohl, E. E. (2012). Dams in the Cadillac Desert: downstream effects in a geomorphic context. *Annals of the New York Academy of Sciences*, 1249, 227-246. <https://doi.org/10.1111/j.1749-6632.2011.06411.x>
- Sanches, P. V., Nakatani, K., Bialetzki, A., Baumgartner, G., Gomes, L. C., & Luiz, E. A. (2006). Flow regulation by dams affecting ichthyoplankton: the case of the Porto Primavera Dam, Paraná River, Brazil. *River Research and Applications*, 22, 555-565.
<https://doi.org/10.1002/rra.922>
- Schwarz, C., & Arnason, A. (1996). A general methodology for the analysis of capture-recapture experiments in open populations. *Biometrics*, 52, 860-873.
<https://doi.org/10.2307/2533048>
- Stanford, J. A., & Ward, J. V. (2001). Revisiting the serial discontinuity concept. *Regulated Rivers: Research and Management*, 17, 303-310. <https://doi.org/10.1002/rrr.659>
- Stanley, T. R., & Burnham, K. P. (1999). A closure test for time-specific capture-recapture data. *Environmental and Ecological Statistics*, 6, 197-209.
<https://doi.org/10.1023/A:1009674322348>
- Stevens, L. E., Ayers, T. J., Bennett, J. B., Christensen, K., Kearsley, M. J. C., Meretsky, V. J., Phillips III, A. M., Parnell, R. A., Spence, J., Sogge, M. K., Springer, A. E., & Wegner, D. L. (2001). Planned flooding and Colorado River riparian trade-offs downstream from Glen Canyon Dam, Arizona. *Ecological Applications*, 11, 701-710.
<https://doi.org/10.2301/3061111>
- Species Tagging, Research and Monitoring System (STReaMS). (2017). Accessed via the internet at <https://streamsystem.org> on 7/20/2017.
- Taylor, C. A., Knought, J. H., & Hiland, T. M. (2001). Consequences of stream impoundment on fish communities in a small North American drainage. *River Research and Applications*, 17, 687-698. <https://doi.org/10.1002/rrr.629>
- Tyus, H. M., & McAda, C. W. (1984). Migration, movements, and habitat preferences of Colorado squawfish, *Ptychocheilus lucius*, in the Green, White, and Yampa rivers, Colorado and Utah. *The Southwestern Naturalist*, 29, 289-299.
<https://doi.org/10.2307/3671360>

505 Tyus, H. M., & Karp, C. A. (1990). Spawning and movements of Razorback Sucker,
 506 *Xyrauchen texanus*, in the Green River basin of Colorado and Utah. *The Southwestern*
 507 *Naturalist*, 35, 427-433. <https://doi.org/10.2307/3672041>

508 U.S. Fish and Wildlife Service (USFWS). (2017). Population abundance estimates of Colorado
 509 pikeminnow and razorback sucker in the San Juan River. U. S. Fish and Wildlife
 510 Service, San Juan River Basin Recovery Implementation Program, New Mexico
 511 Ecological Services Field Office, Albuquerque, New Mexico 87113.

512 Volke, M. A., Scott, M. L., Johnson Carter, W., & Dixon, M. D. (2015). The ecological
 513 significance of emerging deltas in regulated rivers. *BioScience*, 65, 598-611.
 514 <https://doi.org/10.1093/biosci/biv040>

515 P. A., Webber, & Beers, D. 2014. Detecting Razorback Suckers using passive integrated
 516 transponder tag antennas in the Green River, Utah. *Journal of Fish and Wildlife*
 517 *Management* 5, 191–196. <https://doi.org/10.3996/122012-JFWM-104>

518 Zelasko, K. A., Bestgen, K. R., & White, G. C. (2010). Survival rates and movement of
 519 hatchery-reared razorback suckers in the Upper Colorado River Basin, Utah and
 520 Colorado. *Transactions of the American Fisheries Society*, 139, 1478-1499.
 521 <https://doi.org/10.1577/T09-107.1>

522

523 Table 1. Number of individual fish detected by a Passive Integrated Transponder antenna or captured during sampling efforts
 524 downstream of a waterfall barrier on the San Juan River, Utah. Since fishes could be both detected and sampled, the “Number unique”
 525 column indicates the total number of unique fishes recorded from all sampling and detection data.

Species	Year	Days detecting	Days sampling	Number detected	Number captured	Number unique	Percent female	Total length (mm) (mean $\pm$ SD)	Weight (g) (mean $\pm$ SD)
Razorback									
sucker	2015	107	6 ^a	499	16	507	---	---	---
	2016	36	6	472	167	523	53%	483 $\pm$ 39	1251 $\pm$ 323
	2017	111	13	615	183	689	48%	502 $\pm$ 36	1340 $\pm$ 348
	<i>Total unique</i>			943	335	1015			
Colorado									
pikeminnow	2015			15	6	19	---	---	---
	2016			8	6	13	---	330 $\pm$ 126	418 $\pm$ 613
	2017			7	6	13	---	214 $\pm$ 95	122 $\pm$ 186
	<i>Total unique</i>			24	18	39			

526 ^aNote: Sampling in 2015 was a pilot effort of multiple gears including castnets, gillnets, and beach seines but not raft electrofishing.
 527 Consequently, effort was less intensive compared to 2016 and 2017.

528

529

Figure 1. Study area showing the stocking or tagging event location and relative abundance of passive integrated transponder tagged endangered fishes detected or captured downstream of the waterfall (shown by black line labeled ‘waterfall’) in 2015, 2016, or 2017. Tags were matched to records in the Species Tagging Research and Monitoring System (STReamS 2017, accessed 7/20/2017, <https://streamsystem.org>). Lake Powell is shown at full pool.

537

538 Figure 2. Lake Powell reservoir surface elevation metrics (maximum and minimum annual
 539 elevation) and thresholds (full pool and waterfall elevations) since Glen Canyon Dam operations
 540 began. Lake Powell and the San Juan River inflow are characterized by four phases since 1963:
 541 1) filling to capacity, 2) elevation declines leading to emergence of first waterfall, 3) refilling of
 542 reservoir inundating the initial waterfall, and 4) subsequent declines and prolonged water
 543 shortage leading to the current waterfall. The star indicates a two-week period of waterfall
 544 inundation in July-August 2011 that was not captured by mean annual reservoir elevation.
 545 Shaded phases indicate times when the waterfall is present and a barrier to fish passage.

546

Figure 3. Time series of photos of the San Juan River arm of Lake Powell showing the dynamic water levels at the inflow area since 1985. The location of the current waterfall, shown in all photos, is indicated by the white filled circle. The plus and minus signs next to years indicates the presence (+) or absence (-) of a waterfall, respectively. Open circle in 1992 indicates location of the first waterfall that existed from the late-1980s to the mid-1990s. Arrow indicates Clay Hills Crossing, UT.

Figure 4. A photo of the Piute Farms waterfall in 2015 looking downstream towards Lake Powell reservoir (~177 km upstream of Glen Canyon Dam).

559

560 Figure 5. Passive integrated transponder (PIT) tag detections of razorback sucker at a
 561 submersible PIT tag antenna stationed immediately downstream of the Piute Farms waterfall
 562 (top) and coinciding environmental conditions of the San Juan River from 2015, 2016, and 2017.
 563 Dashed lines in the middle panel represent the upper and lower bounds of observed spawning
 564 temperatures for razorback sucker (Tyus and Karp 1990).

Supplemental figure 1. Data used to estimate the population size of passive integrated transponder tagged razorback sucker downstream of the Piute Farms waterfall from February 11 to June 3, 2017. Cumulative number of unique detected razorback sucker (line, left y-axis) over the four detection periods. Bars (right y-axis) represent the proportion of new fish detected during each period. From left to right, the corresponding time intervals for each column are 32 d, 15 d, 14 d, and 49 d.

Supplemental Table 1. Full model set for population size estimation using POPAN, an open population model, for razorback suckers detected by a passive integrated transponder antenna during sampling efforts downstream of a waterfall barrier on the San Juan River, Utah, 12 February-03 June 2017.

Model ^a	K	AIC _c	ΔAIC _c	w _i	Deviance
$\phi(.) p(t) p_{ent}(t)^b$	9	1393.022	0.000	0.814	-1161.401
$\phi(t) p(t) p_{ent}(t)$	11	1395.978	2.956	0.186	-1162.525
$\phi(t) p(.) p_{ent}(t)$	8	1424.293	31.272	0.000	-1128.096
$\phi(.) p(.) p_{ent}(t)$	6	1431.155	38.134	0.000	-1117.177

^a Key to model notation: K = no. of parameters; AIC_c = Akaike Information Criteria corrected for small sample size and lack of model fit; ΔAIC_c = difference between the model listed and the AIC_c of the best model; w_i = model weight based on model AIC_c compared to all other model AIC_c values; . = constant time; t = sampling occasion; φ = probability of leaving the waterfall population, p = probability of detection, p_{ent} = probability of entering the waterfall population.

^b Models without time variation in p_{ent} (i.e., p_{ent}(.)) failed to converge.