Utah State University

DigitalCommons@USU

All Graduate Theses and Dissertations

Graduate Studies

5-1960

Hydrogen Bonding Between the Carbonyl Group and Wyoming Bentonite

Robert A. Kohl

Utah State University

Follow this and additional works at: <https://digitalcommons.usu.edu/etd>

Part of the [Soil Science Commons](#)

Recommended Citation

Kohl, Robert A., "Hydrogen Bonding Between the Carbonyl Group and Wyoming Bentonite" (1960). *All Graduate Theses and Dissertations*. 3804.

<https://digitalcommons.usu.edu/etd/3804>

This Thesis is brought to you for free and open access by the Graduate Studies at DigitalCommons@USU. It has been accepted for inclusion in All Graduate Theses and Dissertations by an authorized administrator of DigitalCommons@USU. For more information, please contact digitalcommons@usu.edu.

HYDROGEN BONDING BETWEEN THE CARBONYL

GROUP AND WYOMING BENTONITE

by

Robert A. Kohl

A thesis submitted in partial fulfillment
of the requirements for the degree

of

MASTER OF SCIENCE

in

Soil Physics

UTAH STATE UNIVERSITY
Logan, Utah

1960

3782

K823

C.2

ACKNOWLEDGMENTS

I am very grateful to Dr. S. A. Taylor for his helpful suggestions. I also wish to express my appreciation to Dr. L. G. Tensmeyer for his advice and assistance and to Dr. R. L. Smith and Dr. J. T. Spence for their suggestions.

Robert Kohl

TABLE OF CONTENTS

	Page
Introduction	1
Review of literature	2
Experimental methods	7
Experimental results	10
Discussion	11
Summary	21
Literature cited	22
Appendix	25
Suggestions for future work	30

LIST OF FIGURES

Figure	Page
1. The spectra of hydrogen bentonite (A), acetic acid (B), and acetic acid adsorbed on hydrogen bentonite (C) . . .	13
2. The spectra of VAMA and VAMA adsorbed on calcium bentonite in various concentrations	14
3. The spectra of VAMA and VAMA adsorbed on hydrogen bentonite in various concentrations	15
4. The spectra of the acid form of HPAN and HPAN adsorbed on hydrogen bentonite in two concentrations	16
5. The spectra of diethyl ketone (A) and diethyl ketone adsorbed on calcium bentonite (B)	18
6. The spectra of polygalacturonic acid (A) and 4 g of polygalacturonic acid on 100 g of calcium bentonite (B)	20
7. The spectra of HPAN as the calcium salt (A), as the acid (B), and as the natural sodium salt (C)	27
8. The spectra of HPAN adsorbed on calcium bentonite in various concentrations	28
9. The spectra of IBMA as the calcium salt (A), as the natural ammonium salt (B), and as the acid (C)	29
10. The spectra of IBMA adsorbed on calcium bentonite in two concentrations and on hydrogen bentonite in two concentrations	31

INTRODUCTION

The vibrational frequencies of atom to atom bonds within a molecule are a function of the bond energies. Each bond has its characteristic frequency, and most of these frequencies can be detected with the infra-red spectro-photometer. When one compound reacts with another or is adsorbed on the surface of a solid, detectable frequency changes or shifts may occur. These changes or shifts yield valuable information about the bonds which are formed or broken.

Infra-red spectroscopy was used in this study to investigate the nature of the bonds that might form between organic molecules and silicate minerals.

REVIEW OF LITERATURE

Early in the study of factors affecting soil aggregation, good correlations were found to exist between aggregate stability and the organic and clay contents of the soil (1, 22). Inorganic oxides were also found to be important in the aggregation of some soils. In a detailed study Kroth and Page (18) showed that inorganic oxides (iron and aluminum), fats, waxes, and resins formed a continuous matrix which bound the soil particles together by physical forces alone.

Soil microbiologists found that the products of active microbial decomposition were much more effective aggregators than fresh organic matter or completely composted organic matter (5, 19, 20, 21, 25). These products often carry several active groups, the most common of which are the hydroxyl, carboxyl, and amino groups. Few studies have given direct evidence as to the methods by which these groups may enhance aggregation.

Workers who have encountered difficulty in extracting organic matter from the soil noted that polyuronides were not removed during the extracting process and that the C/N ratio in the residue was lower than that of the original soil (32). The lower C/N ratio in the residue suggested that nitrogenous organic compounds form strong bonds with inorganic colloids.

When large organic cations were adsorbed on montmorillonite the c-axis spacing was greatly increased (13, 16). Only other large organic cations were able to replace the adsorbed cations. It was concluded that organic molecules which are cationic in nature, perhaps as a result

of amino groups, are attached to the inorganic colloid by coulombic forces (13, 16).

Several workers (9, 10, 29) have presented further evidence that amino cations form bonds with clay minerals by noting that bacteria encountered difficulty in attacking proteins which were adsorbed on montmorillonite.

Amino acids and free amino groups have been found to exist in soil organic matter (27, 31). It is therefore possible that long chain organic molecules containing amino acids may bind soil particles together into aggregates by being adsorbed at cation exchange sites on the clay minerals and by acting as bridges between the particles.

X-ray defraction analyses have shown that organic anions do not enter between the montmorillonite lattice. However, since synthetic polymers which contain carboxyl groups (soil conditioners) stabilize inorganic soil colloids, it has been postulated that organic anions are adsorbed on the edges of the crystals where anion adsorption sites are known to exist (7, 8, 14, 15, 17, 26, 30). Definite evidence to support this theory has not appeared in the literature as yet.

Using electron micrographs, Kroth and Page (18) noted that polar substances which resulted from the decomposition of fresh organic matter were uniformly distributed throughout natural and synthetic aggregates. The dried aggregates were not easily disrupted on rehydration. The authors concluded that the polar organic substances probably formed physicochemical bonds with the surface active clays.

A similar linkage was proposed by Michaels (23), who used the sediment volume technique. Noting that the best flocculation was obtained with polyelectrolytes which were 33 percent hydrolyzed, he postulated that the polymers effected aggregation by the adsorption (via

ester formation or hydrogen bonding) of hydroxyl or amide groups on the solid surfaces, each polymer chain adsorbing on, and bridging between, more than one solid particle.

Recently Emerson (8) compared the effects of synthetic soil conditioners and soil organic matter on aggregate stabilization. He used three polymers, polyvinyl alcohol, a non-ionic polymer, and two carboxylated polymers, high molecular weight sodium alginate and vinyl acetate maleic acid (VAMA). Because sodium hydroxide and an excess of sodium pyrophosphate displaced the carboxylated polymers to various degrees, it was concluded that these polymers were hydrogen bonded to the edge of the clay materials. Polyvinyl alcohol was not displaced by sodium hydroxide because its hydrogen bonds are probably independent of pH. Since the alcohol entered between the clay lattice, it was protected from the effect of sodium pyrophosphate as the pyrophosphate anion could not enter between the lattice due to its charge. Strong sodium hydroxide and sodium pyrophosphate did not displace the binding organic matter from the natural soil aggregates. Emerson concluded that natural aggregates are bound together by organic molecules between the crystal lattice joined by hydrogen bonding.

Peterson (28) noted an increase in the thixotropy of clay suspensions to which he had added calcium and pectin. Based on the research of Bower and Truog (4) who found that polyvalent cations hydrolyze in the following manner

pure compound by noting the frequencies at which absorption takes place and recalling the bonds which absorb at those particular frequencies. In practice this can only be approximated as resonance and coupling take place between bonds with similar frequencies.

It is of importance in this study to note that group frequencies are altered as a result of hydrogen bonding. When the carbonyl group undergoes hydrogen bonding, the oxygen carbon double bond is slightly weakened which results in a slight lowering in its vibration frequency. This shows up as a shift in the infra-red band attributed to the carbonyl group.

Infra-red absorption has been used to study the adsorption of organic and inorganic substances on the surfaces of metals, oxides, and silicates (6, 11, 34).

Infra-red absorption was used by Holmes and Toth (17) to determine a possible linkage between a synthetic soil conditioner and clay minerals. It was noted that the absorption band indicating free OH groups on the clay mineral disappeared when large amounts of the conditioner were adsorbed. This indicated that hydrogen bonding had taken place between the soil conditioner and the clay mineral, but it was not determined which of the available groups, the amide, phenolic, or the carboxyl groups, were active.

It appears that further use of infra-red spectra may provide information about the actual mechanisms by which organic molecules are linked to silicate minerals.

EXPERIMENTAL METHODS

Montmorillonite no. 26 (bentonite from Clay Spur, Wyoming) was obtained. The bentonite, as received, contained significant amounts of quartz. Therefore, the bentonite was dispersed in deionized water and centrifuged in a Sharples Super Centrifuge. Only that portion below 0.2μ was retained for use, as it was relatively free of quartz.

Using an International Centrifuge, size 2, the centrifuged bentonite samples were saturated with calcium and hydrogen using calcium chloride and hydrogen chloride in dilute solutions. Enough calcium chloride or hydrogen chloride was added to the bentonite solution to cause flocculation. The samples were centrifuged and the supernatant liquid was decanted and discarded. Deionized water was added, the samples were shaken for five minutes and recentrifuged. The supernatant liquid was again discarded and the process repeated until the suspensions were dispersed again. The flocculation, centrifuging and dispersion was repeated four times to assure saturation with the desired ion. After washing with deionized water to the point of dispersion the last time, the calcium saturated bentonite had a pH of 6.6 and the hydrogen saturated bentonite a pH of 1.7.

Several organic compounds were used to determine a possible linkage between organic materials and montmorillonite. The organic compounds used were acetic acid, diethyl ketone, vinyl acetate maleic acid (VAMA), a copolymer of isobutylene and the half ammonium-half amide salt of maleic acid (IBMA), the sodium salt of hydrolyzed polyacrylonitrile (HPAN), and polygalacturonic acid. The structural formulas of the less

common organic compounds are:

VAMA

HPAN

IBMA

Polygalacturonic acid

Samples were prepared by placing various quantities of the adsorbate in a 50 ml Erlenmeyer flask containing 5 ml of deionized water. Five ml of a solution containing 0.021 g of bentonite per ml were added to the flask. The mixture was shaken on a wrist action shaker for one hour and allowed to stand for 16 hours.

The samples were poured into a petri dish and dried at 55° C in an oven. The residue was ground with a mortar and pestal and then subjected to infra-red inspection by the KBr pressed-disk technique (33), using a

Beckman die of diameter one-half inch. A Perkin-Elmer Model 21 Double Beam Recording Spectro-photometer was used to obtain the spectra.

EXPERIMENTAL RESULTS

The spectra of several organic compounds adsorbed on centrifuged Wyoming bentonite were obtained. That portion of the spectra of these mixtures which showed a change due to adsorption is given in figures 1 through 10. Most of the compounds showed activity in the 5.5μ to 6.5μ region though some were active over the 5.5μ to 7.5μ region.

The band which appears at 6.16μ in all spectra containing bentonite is due to the adsorbed water on the bentonite itself.

The units on the absorbance scale of the figures are not necessary as quantitative data was not sought. It was found convenient to space the spectra so as to get several in one figure. Therefore it would not be convenient to place units on the absorbance scale as several scales would be required in each figure.

DISCUSSION

The infra-red absorption band arising from the C=O stretching vibration has been extensively studied, and much is known of the factors which influence its frequency. This band lies in the region between 5.8μ and 5.9μ (3) unless it is hydrogen bonded, part of an anhydride linkage, or in the dipole field of a halogen.

Hydrogen bonding or the forces attracting a proton to the carbonyl oxygen causes a slight weakening in the C=O bond. This weakened bond has a lower frequency of vibration than the original carbonyl double bond, and results in a small increase of the resulting wave length. Thus, the absorption band arising from the carbonyl frequency will be shifted toward the right or to a longer wave length. The literature reviewed indicated that only hydrogen bonding could lead to a 0.05μ or greater shift to the right in the carbonyl wave length.

When two carboxyl groups form an anhydride, two carbonyl frequencies show up in the infra-red spectrum, one between 5.60μ and 5.75μ , and one between 5.40μ and 5.55μ . The two bands are usually about 0.2μ apart. If the anhydride linkage is part of a strained cyclic structure, the two carbonyl frequencies are slightly greater than the open-chain frequencies. The reason for the splitting of the carbonyl band has not yet been adequately solved. The most likely explanation is that the two carbonyl frequencies are coupled together, leading to resonance splitting (3).

An α -halogen-substituted carbonyl containing compound shows an increase in the carbonyl frequency. Bellamy (3) suggests that the near

approach of a negatively charged halogen atom to the readily polarisable negatively charged oxygen of the carbonyl group would result in a mutual induction of opposite charges. The negative character of both the halogen and oxygen atoms would be diminished and the C-X and C=O bonds would become less polar. This would result in a rise of their vibrational frequencies.

Figure 1 shows the spectra of hydrogen bentonite, acetic acid, and the acid adsorbed on hydrogen saturated bentonite. The carbonyl wave length of 5.90μ for the acid has shifted to 5.97μ in the adsorbed spectrum. This suggests that hydrogen bonding has taken place between the double bonded oxygen of the carbonyl group and the clay mineral. This shift in the carbonyl band could not represent the formation of an ionic bond with an adsorbed polyvalent cation as the ionized carboxyl group does not absorb radiation at this wave length.

Figures 2 and 3 show the spectra of VAMA adsorbed on calcium and hydrogen saturated bentonite. The C=O band at 5.82μ in the acid shifts to 5.89μ when VAMA is adsorbed on bentonite. This suggests that strong hydrogen bonds are formed between the carbonyl groups of VAMA and the bentonite. As larger amounts of VAMA were added to a constant quantity of bentonite, the 5.82μ band of the free carbonyl group shows up in the adsorbed spectra. This indicates that some C=O groups are present which are not bonded to the clay.

Comparing figures 2 and 3, VAMA showed a similar degree of hydrogen bonding through the carbonyl group to calcium bentonite as to hydrogen bentonite. This indicates that the adsorbed cation does not affect the hydrogen bonding of the carbonyl group in VAMA.

Figure 4 shows the acid form of HPAN and HPAN adsorbed on hydrogen saturated bentonite. The carbonyl band at 5.80μ shifts to 5.85μ when

FIGURE 1. THE SPECTRA OF HYDROGEN BENTONITE (A), ACETIC ACID (B),
AND ACETIC ACID ADSORBED ON HYDROGEN BENTONITE (C)

FIGURE 2. THE SPECTRA OF VAMA AND VAMA ADSORBED ON CALCIUM BENTONITE IN VARIOUS CONCENTRATIONS

FIGURE 3. THE SPECTRA OF VAMA AND VAMA ADSORBED ON HYDROGEN BENTONITE IN VARIOUS CONCENTRATIONS

FIGURE 4. THE SPECTRA OF THE ACID FORM OF HPAN AND HPAN
ADSORBED ON HYDROGEN BENTONITE IN TWO CONCENTRATIONS

HPAN is adsorbed on the clay mineral. This also suggests hydrogen bonding.

Diethyl ketone was studied in an attempt to isolate the effects of the carbonyl group. Figure 5 shows the spectra of diethyl ketone and the ketone adsorbed on calcium saturated bentonite. It is noted that the strong carbonyl band of the ketone at 5.85μ shifts to 5.94μ when the ketone is adsorbed on bentonite. This indicates that the carbonyl group is active in hydrogen bonding to the clay mineral.

Since the carbonyl group does hydrogen bond to the clay mineral, one might speculate as to the site on the clay mineral which is taking part in the bond formation.

Several workers (7, 15, 17, 30) have shown by X-ray defraction that organic anions such as the ones used in this study do not enter between the crystal lattice of montmorillonite type clays. Since a hydrogen atom is necessary and since one attached to an oxygen atom has been shown to be effective in forming dimers and polymers in alcohols and acids, it is proposed that a hydroxyl group exposed on the broken edge of the clay crystal is the reactive site.

When a hydroxyl group is involved in hydrogen bonding, it has been shown that the OH stretching frequency is reduced and the OH deformation frequencies are increased. These frequency shifts were not observed due to the masking of these regions by two factors: (a) by the many internal hydroxyl groups of the clay mineral which are not in a position to take part in the proposed mechanism and (b) by the adsorbed water on the clay mineral.

Several authors (7, 8, 17, 26, 30) have proposed hydrogen bonding to the edge faces of the clay crystal but have not suggested that the

FIGURE 5. THE SPECTRA OF DIETHYL KETONE (A) AND DIETHYL KETONE ADSORBED ON CALCIUM BENTONITE (B)

carbonyl group is responsible. They have either failed to suggest the reactive groups altogether or have speculated that the hydroxyl group of an organic acid was the reactive group.

The data presented in this paper agree in principle with the above authors except for showing that the carbonyl group is active.

It has been found that polygalacturonic acid is a constituent of soil organic matter (12). Figure 6 shows the spectra of polygalacturonic acid and the acid adsorbed on hydrogen saturated bentonite. The 5.78μ band of polygalacturonic acid shifts to 5.88μ when adsorbed on hydrogen bentonite. Again, this indicates that the carbonyl group has hydrogen bonded to the bentonite.

Since polyuronides may represent 30 percent of the soil organic matter (12), this suggests a possible mechanism by which organic matter may bind soil particles together into aggregates. However, it should be emphasized that other mechanisms are also possible and that this study does not indicate their presence nor confirm their absence.

Since postulations on the mechanisms involved in binding the remaining compounds to the clay mineral are highly speculative, they are included in the appendix along with suggestions for further study.

FIGURE 6. THE SPECTRA OF POLYGALACTURONIC ACID (A) AND 4 g
OF POLYGALACTURONIC ACID ON 100 g OF CALCIUM
BENTONITE (B)

SUMMARY

Several organic compounds were adsorbed on centrifuged Wyoming bentonite. Their infra-red spectra were obtained and studied for a possible mechanism by which they were bound to the clay mineral.

A definite shift to the right in the carbonyl stretching frequency has been shown. This shift is the result of hydrogen bonding between the double bonded oxygen of the carbonyl group and a hydrogen atom in the clay crystal. It is postulated that an exposed hydroxyl group on the broken edge of the clay crystal is the reactive group on the bentonite.

Polygalacturonic acid, a polyuronide, also indicated hydrogen bonding through its carbonyl group. Since polyuronides may represent 30 percent of the soil organic matter, it is suggested that hydrogen bonding between the carbonyl group of an organic compound and an exposed hydroxyl group on the edge of the clay crystal is a possible mechanism by which organic matter may bind soil particles together into aggregates.

LITERATURE CITED

- (1) Alderfer, R. B. and Merkle, F. G. The comparative physical constitution of water-stable granules and of the originating soil. *Soil Sci.* 53:105-113. 1942.
- (2) Allison, L. E. Effect of soil conditioning polymer on the cation-exchange capacity. *Soil Sci.* 83:391-397. 1957.
- (3) Bellamy, L. J. *The Infra-red Spectra of Complex Molecules.* 2nd ed. New York, John Wiley & Sons, Inc. 1958.
- (4) Bower, C. A. and Truog, E. Base exchange capacity determination as influenced by nature of cation employed and formation of basic exchange salts. *Soil Sci. Soc. Amer. Proc.* 5:86-89. 1941.
- (5) Chesters, G., Attoe, O. J., and Allen, O. N. Soil aggregation in relation to various soil constituents. *Soil Sci. Soc. Amer. Proc.* 21:272-277. 1957.
- (6) De Mumbrum, L. E. and Jackson, M. L. Infrared absorption evidence of exchange reaction mechanism of copper and zinc with layer silicate clays and peat. *Soil Sci. Soc. Amer. Proc.* 20:334-337. 1956.
- (7) Emerson, W. W. Synthetic soil conditioners. *J. Agr. Sci.* 47:117-121. 1956.
- (8) Emerson, W. W. A comparison between the mode of action of organic matter and synthetic polymers in stabilizing soil crumbs. *J. Agr. Sci.* 47:350-353. 1956.
- (9) Ensminger, L. E. Resistance of clay-adsorbed proteins to proteolytic hydrolysis. *Soil Sci.* 53:205-209. 1942.
- (10) Ensminger, L. E. and Gieseking, J. E. The absorption of proteins by montmorillonitic clays. *Soil Sci.* 48:467-471. 1939.
- (11) Eyring, E. M. and Wadsworth, M. E. Differential infrared spectra of adsorbed monolayers-hexanethiol on zinc minerals. *Mining Eng.* 8:531-535. 1956.
- (12) Fraser, G. K. Soil organic matter. Chapter 5. *Chemistry of the Soil.* F. E. Bear, editor. New York, Reinhold Publishing Corp. 1955.
- (13) Gieseking, J. E. Mechanism of cation exchange in the montmorillonite-beidellite-nontronite type of clay minerals. *Soil Sci.* 47:1-13. 1939.

- (14) Grim, R. E. Clay Mineralogy. New York, McGraw and Hill. 1953.
- (15) Hagin, J. and Bodman, G. B. Influence of the polyelectrolyte CRD-186 on aggregation and other physical properties of some California and Israeli soils and some clay minerals. Soil Sci. 78:367-378. 1954.
- (16) Hendricks, S. B. Base exchange of the clay mineral montmorillonite for organic cations and its dependence upon adsorption due to Van der Waals forces. J. Phys. Chem. 45:65-81. 1941.
- (17) Holmes, R. M. and Toth, S. J. Physico-chemical behavior of clay-conditioner complexes. Soil Sci. 84:479-487. 1957.
- (18) Kroth, E. M. and Page, J. B. Aggregate formation in soils with special reference to cementing substances. Soil Sci. Soc. Amer. Proc. 11:27-34. 1946.
- (19) Martin, J. P. Microorganisms and soil aggregation: I. Origin and nature of some of the aggregating substances. Soil Sci. 59:163-174. 1945.
- (20) Martin, J. P. Microorganisms and soil aggregation: II. Influence of bacterial polysaccharides on soil structure. Soil Sci. 61:157-166. 1946.
- (21) McCalla, T. M., Haskins, F. A., and Frolik, E. F. Influence of various factors on aggregation of Peorian loess by microorganisms. Soil Sci. 84:155-161. 1957.
- (22) McHenry, J. R. and Russell, M. B. Elementary mechanics of aggregation of puddled materials. Soil Sci. Soc. Amer. Proc. 8:71-78. 1943.
- (23) Michaels, A. S. Aggregation of suspensions by polyelectrolytes. Ind. Eng. Chem. 46:1485-1490. 1954.
- (24) Mortensen, J. L. Adsorption of hydrolyzed polyacrylonitrile (HPAN) on kaolinite: I. Effect of exchange cation and anion. Soil Sci. Soc. Amer. Proc. 21:385-388. 1957.
- (25) Myers, H. E. and McCalla, T. M. Changes in soil aggregation in relation to bacterial numbers, hydrogen-ion concentration, and length of time soil was kept moist. Soil Sci. 51:189-199. 1941.
- (26) Packter, A. Interaction of montmorillonite clays with polyelectrolyte. Soil Sci. 83:335-343. 1957.
- (27) Packer, D. I., Sowden, F. J., and Atkinson, H. J. The nitrogen distribution and amino acid content of certain soil organic matter fractions. Sci. Agr. 32:163-169. 1952.
- (28) Peterson, J. B. Calcium linkage, a mechanism in soil granulation. Soil Sci. Soc. Amer. Proc. 12:29-34. 1947.

- (29) Pinck, L. A., Dyal, R. S., and Allison, F. E. Protein-montmorillonite complexes, their preparation and the effects of soil microorganisms on their decomposition. *Soil Sci.* 78:109-118. 1954.
- (30) Ruehrwein, R. A. and Ward, D. W. Mechanism of clay aggregation by polyelectrolytes. *Soil Sci.* 73:485-492. 1952.
- (31) Sowden, F. J. Note on the occurrence of amino groups in soil organic matter. *Can. J. Soil Sci.* 37:143-144. 1957.
- (32) Sowden, F. J., Parker, D. I., and Atkinson, H. J. Comparison of organic matter fractions from three soil types. *Sci. Agr.* 32:127-134. 1952.
- (33) Stimson, M. M. and O'Donnell, M. J. The infrared and ultraviolet absorption spectra of cytosine and isocytosine in the solid state. *J. Amer. Chem. Soc.* 74:1808. 1952.
- (34) Tensmeyer, L. G., and Wadsworth, M. E. Quantative determination of adsorbed sulfates on thoria by means of infrared spectroscopy. Tech. report III. University of Utah. Salt Lake City, Utah. 1959.
- (35) Tiulin, A. F. Colloidal chemical investigation of soils for agricultural purposes. Moscow: The Lenin Academy of Agricultural Sciences. 1946.

APPENDIX I

APPENDIX

Two of the compounds that were studied did not show hydrogen bonding through the carbonyl group when the carboxyl group was ionized.

Figure 7 shows the spectra of HPAN as the natural sodium salt, as the calcium salt, and as the acid. The band at 6.42μ is a result of the anti-symmetrical vibration and the one at 7.10μ or 7.16μ is a result of the symmetrical vibration of the COO- structure. The band at 6.90μ is due to the CH₂ deformation mode. The symmetrical, ionized carboxyl band in the calcium salt lies to the left of that of the sodium salt. A small unidentified band at 6.02μ is present in the sodium salt form of HPAN.

Figure 8 shows the spectra of HPAN adsorbed on calcium saturated bentonite in various concentrations. It is noted that when the natural sodium salt of HPAN is adsorbed on calcium saturated bentonite, the symmetrical ionized carboxyl band resembles that of the calcium salt of HPAN. This might suggest a linkage similar to that proposed by J. B. Peterson (28). However, the evidence is meager and further work should be carried out using bentonite saturated with various mono- and polyvalent ions and using various salts of HPAN.

It is interesting to note the increased prominence of the unidentified band at 6.02μ . If this band could be identified, it would help in determining the mechanism by which HPAN is bonded to montmorillonite type clay minerals.

Figure 9 shows the spectra of IBMA as the calcium salt, as the

FIGURE 7. THE SPECTRA OF HPAN AS THE CALCIUM SALT (A), AS
THE ACID (B), AND AS THE NATURAL SODIUM SALT (C)

FIGURE 8. THE SPECTRA OF HPAN ADSORBED ON CALCIUM
BENTONITE IN VARIOUS CONCENTRATIONS

FIGURE 9. THE SPECTRA OF IBMA AS THE CALCIUM SALT (A), AS
THE NATURAL AMMONIUM SALT (B), AND AS THE ACID (C)

natural ammonium salt, and as the acid. The band at 5.85μ is due to the carbonyl group. The two bands at 6.43μ and 7.16μ are due to the anti-symmetrical and symmetrical vibrations of the ionized carboxyl group. The bands at 5.44μ and 5.65μ in the spectrum of the acid form of IBMA seem to be due to the formation of an anhydride.

Figure 10 shows the spectra of IBMA adsorbed on calcium and hydrogen saturated bentonite. The anhydride bands show up only slightly when IBMA is adsorbed on calcium saturated bentonite but are quite pronounced when adsorbed on hydrogen saturated bentonite. The ionized carboxyl bands are present in the case of the calcium bentonite and the carbonyl band is present with hydrogen bentonite. However, in both cases there is a new band at 7.02μ .

There is a possibility that this new band is the result of a shift in the symmetrical ionized carboxyl vibration. Such a shift might be caused by the approach of the ionized carboxyl group to a positive exchange site on the edge of the clay mineral.

Suggestions for future work

Further study should include the adsorption of IBMA and HPAN in various salt forms onto bentonite saturated with different ions. This should provide data to either confirm or contradict the linkage proposed by J. B. Peterson (28).

It should be of interest to study a few samples using clay of a smaller size fraction. This should provide more surface area per gram of bentonite and make it possible to take full advantage of the double beam infra-red spectro-photometer.

FIGURE 10. THE SPECTRA OF IBMA ADSORBED ON CALCIUM BENTONITE
IN TWO CONCENTRATIONS AND ON HYDROGEN BENTONITE IN
TWO CONCENTRATIONS.