

4-15-2015

Thermal Evidence of Flatslab Subduction Perturbations in the Western US

Michael Berry
Utah State University

Follow this and additional works at: https://digitalcommons.usu.edu/phys_capstoneproject

 Part of the [Physics Commons](#)

Recommended Citation

Berry, Michael, "Thermal Evidence of Flatslab Subduction Perturbations in the Western US" (2015). *Physics Capstone Project*. Paper 19.

https://digitalcommons.usu.edu/phys_capstoneproject/19

This Report is brought to you for free and open access by the Physics Student Research at DigitalCommons@USU. It has been accepted for inclusion in Physics Capstone Project by an authorized administrator of DigitalCommons@USU. For more information, please contact dylan.burns@usu.edu.



Thermal Evidence of Flat-slab Subduction Perturbations in the Western US

Student: Michael Berry

Faculty Mentor: Anthony Lowry

2014

Publication Draft: Further results and analysis will be found in a publication Summer 2015

Abstract

The Laramide Orogeny, a late Cretaceous mountain building event in the western United States, has been postulated to result from flat-slab style subduction of the Farallon plate^{1,2,3}. Possible consequences of flat-slab style subduction on elevation, tectonism and volcanism associated with expected changes in temperature and hydration state of the mantle in the western U.S. has been of considerable interest^{x1}. However, the impact of flat-slab subduction on modern mantle temperature, water and tectonic stability remains poorly-understood. Here we show that thermal perturbation by Farallon flat slab subduction is still evident today in Moho temperatures of the Rocky Mountain and Great Plains regions of the U.S. Measurements of temperature in the uppermost mantle from tomographic P-wave velocity images⁴ are compared to models of surface heat flow measurements assuming steady-state lithospheric thermal transfer and reasonable rock thermal properties. Misfit between modeled and measured Moho temperature indicates a violation of the modeling assumptions, likely dominated by transient perturbations on timescales less than the ~100 Myr required for a conductive thermal pulse to traverse the lithosphere. Misfit is dominated by anomalously low measurements in regions deformed by thick-skin tectonism during the Laramide, suggesting the temperature discrepancy may reflect thermal perturbation by Farallon flat-slab subduction. Shallow subducted slab would serve both to chill the base of the lithosphere and insulate it from deeper convective heating. The measurements indicate perturbation of mantle lithospheric temperature to distances of more than 1800 km from the modern plate boundary, where Laramide-style tectonism was weakly expressed. These results may have significant implications for history, evolution and stability of continental lithosphere within EarthScope's USArray footprint⁵.

Introduction

The subduction and subsequent removal of the proposed Laramide flat slab subducted plate dramatically impacted the geologic landscape of much of the North American Cordillera. The hypothesised removal of the plate caused the so-called "ignimbrite flare-up" of the Mid-Tertiary, and re-introduction of higher temperature mantle to the base of the lithosphere is one likely contributor to uplift of the Colorado Plateau and other Cordilleran regions⁶. Thermal perturbation of the lithosphere by shallowly-dipping subducted slab is expected to be significant because the time required to thermally equilibrate an initial slab temperatures to the ~1350°C potential temperature of asthenospheric mantle are large relative to the time required for transport to distances of order 1000 km from the trench, with final temperatures of subducted oceanic crust



dependent primarily on age of the oceanic lithosphere and rate of subduction convergence⁷.

Thermal modeling

Two important pieces of informational constraint here: surface heat flow and Pn-derived Moho temperature. Need to describe (at least cursorily) important features of heat transfer that determine the geotherm: radiogenic heat production in the crust, temperature dependence of thermal conductivity, advective heat transfer processes.

We modeled the geothermal gradient using data from surface heat flow, surface temperatures, radiogenic heat production in the crust, and crustal thickness measurements (Lowry et al.). To arrive at best fitting parameters for conductivity, parameters A and B were grid-searched for each map point for the western US. The values were used to calculate a geothermal gradient for each of the ~50,000 points, this geotherm was then compared against Pn wave inferred moho temperature estimates to produce an overall RMS misfit. Significant efforts were made to reduce the RMS misfit, by varying radiogenic length parameters (*lrad*), surface heat production (A_0), to minimize the RMS. These efforts were unfruitful as they eventually produced non-physical results. Therefore fixed *lrad* (14km) and mean A_0 values were used.

Numerically searched conductivity parameters (*q*) were compared to laboratory values established by Kukkonen et al¹⁰ (See Figure 1). The mapped RMS on a A versus B parameters grid searched yielded consistent trends with lab values. Plotted conductivity parameters of various compositions follows a trend going from quartz poor (more mafic) to quartz rich (more granitic). With the admitted limitations picking a singular best fitting values of $A=.23$ and $B=5.4 \cdot 10^{-4}$ creating a compositionally uniform crust.



Citations

- ¹Coney, P. J. & Reynolds, S. T. Cordilleran Benioff zones. *Nature* **270**, 403–406 (1977).
- ²Henderson, L. J., Gordon, R. G., & Engebretson, D. C. Mesozoic aseismic ridges on the Farallon plate and southward migration of shallow subduction during the Laramide orogeny. *Tectonics* **3** 121-132 (1984).
- ³Humphreys, E. D. Post-Laramide removal of the Farallon slab, western United States. *Geology* **23**, 987-990 (1995).
- ⁴Schutt, D. L., Lowry, A. R., & Buehler, J. S., Moho temperature and mobile lower crust in the western United States. *Geology* submitted (2014).
- ⁵Williams, M. L. et al. Unlocking the secrets of the North American continent: an EarthScope science plan for 2010–2020. <http://www.earthscope.org/ESSP> (2010).
- ⁶Roy, M., Jordan, T. H. & Pederson, J. Colorado Plateau magmatism and uplift by warming of heterogeneous lithosphere. *Nature* **459**, 978-982 (2009).
- ⁷English, J. M., Johnston, S. T., & Wang, K. Thermal modelling of the Laramide orogeny: testing the flat-slab subduction hypothesis. *Earth and Planetary Science Letters* **214**(3), 619-632 (2003).
- ⁸Bunge, H. P., & Grand, S. P. Mesozoic plate-motion history below the northeast Pacific Ocean from seismic images of the subducted Farallon slab. *Nature* **405** 337-340 (2000).
- ⁹Humphreys, E.D, Hessler, E., Dueker, K., Farmer, G. L., Erslev, E., & Atwater, T. How Laramide-age hydration of North American lithosphere by the Farallon slab controlled subsequent activity in the western United States. *International Geology Review* **45**, 575-595 (2003).
- ¹⁰Kukkonen, I. T. & Šafanda, J. Palaeoclimate and structure: the most important factors controlling subsurface temperatures in crystalline rocks. A case history from Outokumpu, eastern Finland. *Geophysical Journal International* **126**, 101-112 (1996).
- ¹¹Lipman, P. W. & Glazner, A. F. Introduction to middle Tertiary cordilleran volcanism: Magma sources and relations to regional tectonics. *Journal of Geophysical Research* **96** 13193-13199 (1991).
- ¹²Lowry, Anthony R., and Marta Pérez-Gussinyé. The role of crustal quartz in controlling Cordilleran deformation. *Nature* **471** 353-357 (2011)