

2011

Rank 2 distributions of Monge equations: Symmetries, equivalences, ex-tensions

Ian M. Anderson
Utah State University

B. Kruglikov

Follow this and additional works at: https://digitalcommons.usu.edu/mathsci_facpub

 Part of the [Mathematics Commons](#)

Recommended Citation

Ian Anderson, Boris Kruglikov, Rank 2 distributions of Monge equations: Symmetries, equivalences, extensions, *Advances in Mathematics*, Volume 228, Issue 3, 20 October 2011, Pages 1435-1465, ISSN 0001-8708, 10.1016/j.aim.2011.06.019.

This Article is brought to you for free and open access by the Mathematics and Statistics at DigitalCommons@USU. It has been accepted for inclusion in Mathematics and Statistics Faculty Publications by an authorized administrator of DigitalCommons@USU. For more information, please contact dylan.burns@usu.edu.



RANK 2 DISTRIBUTIONS OF MONGE EQUATIONS: *SYMMETRIES, EQUIVALENCES, EXTENSIONS*

IAN ANDERSON, BORIS KRUGLIKOV

ABSTRACT. By developing the Tanaka theory for rank 2 distributions, we completely classify classical Monge equations having maximal finite-dimensional symmetry algebras with fixed (albeit arbitrary) pair of its orders. Investigation of the corresponding Tanaka algebras leads to a new Lie-Bäcklund theorem. We prove that all flat Monge equations are successive integrable extensions of the Hilbert-Cartan equation. Many new examples are provided.

INTRODUCTION AND MAIN RESULTS

In his paper of 1895 [L], S. Lie made the important observation that the simplest differential equations, beyond those of finite type, are equations with general solution depending upon 1 function of 1 variable, and he indicates that they are integrable by ODE methods.

Such differential equations can generally be encoded as rank 2 distributions or equivalently as codimension 2 Pfaffian systems on k -dimensional manifolds. For $k = 5$, Goursat showed [G] that these distributions can be realized as Monge equations

$$y' = F(x, y, z, z', z''), \quad (1)$$

while E. Cartan [C1] realized them as reductions of involutive pairs of 2nd order PDEs in 1 dependent and 2 independent variables. The interplay between the Monge and PDE realizations of rank 2 distributions is an important aspect of their geometry.

From the Goursat-Monge viewpoint, one can therefore interpret Cartan's 1910 paper as a detailed analysis of the geometry of Monge equations (1). In what is still considered to be a computational *tour de force*, Cartan solved the equivalence problem for (1) and, as a corollary, deduced that the celebrated¹ Hilbert-Cartan equation

$$y' = (z'')^2 \quad (2)$$

¹We write this in retrospect [H, C2]; in [C1] only the PDE model was explicitly constructed.

is the Monge equation of type (1) with the largest finite-dimensional Lie algebra of symmetries.

Despite Lie's observation and Goursat's contribution, or perhaps because of the complexity of Cartan's analysis, little progress has been made in the symmetry analysis of the general Monge equations

$$y^{(m)} = F(x, y, y', \dots, y^{(m-1)}, z, z', \dots, z^{(n)}) \quad (3)$$

(here and beyond $y = y(x), z = z(x)$ and we assume $m \leq n$).

These underdetermined ordinary differential equations are important in the variational calculus and control theory, in reductions of compatible PDEs and in the symmetry approach to Darboux integrability. Their geometry is the subject of the present article.

The first goal of the paper is to describe the most symmetric equations of type (3). There are different kinds of symmetries and Lie-Bäcklund type theorems clarify the relations between point, contact, generalized (higher) and internal symmetries whenever possible, see [C1, KLV, AKO, GK].

Below we restrict to purely internal symmetries. We will assume $m > 0$, otherwise (3) has an infinite-dimensional algebra of symmetries.

Theorem 1. *Among equations \mathcal{E} of type (3) with fixed $m \leq n$ and $F_{z^{(n)}z^{(n)}} \neq 0$, the most symmetric (realizing $\max \dim \text{sym}(\mathcal{E})$) are internally equivalent to the equation*

$$y^{(m)} = (z^{(n)})^2. \quad (4)$$

The dimension of the symmetry algebra $\text{sym}(\mathcal{E})$ for this equation equals $2n + 5$ for $m = 1, n > 2$ and $2n + 4$ for $m > 1$.

Let us denote by $\mathcal{E}_{m,n}$ the model equation (4). Notice that the equation $\mathcal{E}_{1,1}$ is internally equivalent to the pure jets $J^2(\mathbb{R}, \mathbb{R})$ with (infinite-dimensional) contact symmetry algebra, while $\mathcal{E}_{1,2}$ is the Hilbert-Cartan equation (2) with symmetry algebra $\mathfrak{g}_2 = \text{Lie}(G_2)$ of dimension 14. These are the exceptional cases omitted in Theorem 1.

Remark 1. *As a by-product of the proof of Theorem 1 we conclude that Monge equations \mathcal{E} with $F_{z^{(n)}z^{(n)}} \neq 0$ are strongly regular in the sense of Tanaka [T]. This implies the existence of a Cartan connection for (3) as a $\mathfrak{t}_{m,n}$ -valued 1-form on a natural bundle over the equation \mathcal{E} , where $\mathfrak{t}_{m,n}$ is the Tanaka algebra calculated in Section 3.2.*

As is well-known, equation (3) defines a submanifold \mathcal{E} in the space of jets $J^{m,n}(\mathbb{R}^1, \mathbb{R}^2) \simeq \mathbb{R}^{m+n+3}(x, y, y_1, \dots, y_m, z, z_1, \dots, z_n)$, with $\dim \mathcal{E} = m + n + 2$. The canonical rank 3 distribution

$$\mathcal{C} = \text{Ann}\{dy_i - y_{i+1} dx, dz_j - z_{j+1} dx \mid 1 \leq i < m, 1 \leq j < n\}$$

(Cartan distribution) on the jet space $J^{m,n}(\mathbb{R}^1, \mathbb{R}^2)$ induces a rank 2 distribution Δ on \mathcal{E} . Internal symmetries of (3) are, by definition, symmetries of the pair (\mathcal{E}, Δ) .

As noticed above, rank 2 distributions were investigated in many classical papers. Recently Doubrov and Zelenko [DZ] described their internal geometry via a canonical frame on a bundle over the manifold \mathcal{E} , providing a unique solution for maximally symmetric model among all rank 2 distributions from some generic stratum (called maximal class in loc.cit). This class does not appear to have a clear description for distributions, arising for differential equations (3).

Theorem 1 yields maximally symmetric distributions in a large class of distributions corresponding to Monge equations of a fixed dimension.

Corollary 1. *Among all rank 2 distributions Δ corresponding to equations (3) with fixed $m+n = k+1$ and $F_{z^{(n)}z^{(n)}} \neq 0$, the most symmetric are locally equivalent to the canonical distribution of the equation $\mathcal{E}_{1,k}$.*

The non-degeneracy condition $F_{z^{(n)}z^{(n)}} \neq 0$ is related to the finite-dimensionality of $\text{sym}(\mathcal{E})$ and will be explained in Section 2. It implies that the growth vector of the corresponding distribution starts $(2, 1, 2, \dots)$ and that the distribution is not de-prolongable.

Let us note that, according to [AKO], internal symmetries for equations of type (3), in the particular case $m = n = 2$, coincide with first order generalized symmetries. We can prove much more than this, namely, we prove a new Lie-Bäcklund type theorem.

Theorem 2. *For equations of type (3), with $F_{z^{(n)}z^{(n)}} \neq 0$, $0 < m \leq n$ and (m, n) being different from the exceptional cases $(1, 1)$ and $(1, 2)$, the algebra of internal symmetries coincides with the algebra of external symmetries, induced by Lie transformations of the ambient mixed jet space $J^{m,n}(\mathbb{R}, \mathbb{R}^2)$.*

Lie transformations of $J^{m,n}(\mathbb{R}, \mathbb{R}^2)$ will be described in Section 4. Let us remark that Theorem 2 is the first instance of Lie-Bäcklund type theorem whose proof entirely depends on the structure of the graded Lie algebra associated to the distribution, and cannot be derived by the usual de-prolongation arguments.

To prove our results we exploit and elaborate upon the Tanaka theory [T]. Distributions corresponding to equation (4) are flat in its sense. There exist however underdetermined ODE systems, for which the respective rank 2 distributions cannot be realized as non-degenerate Monge equations. Their flat models can nevertheless be calculated via the same technique.

In Section 5 we relate the algebraic theory of central extensions of Lie algebras to the geometric theory of integrable extensions. We provide a very simple and constructive description of all flat models.

Theorem 3. *Any flat rank 2 distribution with growth vector $(2, 1, 2, \dots)$ (be it represented as a nondegenerate single Monge equation or underdetermined system of ODEs) is a result of successive integrable extensions of the Hilbert-Cartan equation (2).*

This allows us to calculate many examples of flat rank 2 distributions and the corresponding ODE systems.

At the end of the paper we show that our theory does not restrict to ODEs only. We demonstrate relations of our results to integration of determined and overdetermined PDEs. In particular, we construct a flat non-trivial integrable extension of Cartan's most symmetric involutive PDE model of 1910.

The results of this paper indicate that the method of E. Cartan from his seminal paper [C1] can be extended to a much broader class of PDE systems. This generalization and further applications to the method of Darboux [AFV] and compatible overdetermined systems [KL] will be presented in a separate publication.

ACKNOWLEDGMENT. We are grateful to Valentin Lychagin for discussions during the initial stage of the project. Hospitality of Institut Mittag-Leffler (Stockholm) in 2007, MFO (Oberwolfach) in 2008 and Banach center (Warsaw) in 2009 is acknowledged. I.A. was supported by NSF grant DMS-0713830. The computations in Sections 5 and 6 were done, in part, using the DifferentialGeometry package of MAPLE.

1. TANAKA ALGEBRA OF A DISTRIBUTION

Given a distribution $\Delta \subset TM$, its *weak derived flag* $\{\Delta_i\}$ is given via the module of its sections by $\Gamma(\Delta_{i+1}) = [\Gamma(\Delta), \Gamma(\Delta_i)]$ with $\Delta_1 = \Delta$. The *strong derived flag* is given by the filtration $\nabla_{i+1} = \partial\nabla_i$ with $\Gamma(\nabla_{i+1}) = [\Gamma(\nabla_i), \Gamma(\nabla_i)]$ and $\nabla_1 = \Delta$.

We will assume throughout this paper that the distribution is *completely non-holonomic*, i.e. $\Delta_k = TM$ for some k , and we also assume that Δ is *regular*, so that the ranks of both Δ_i and ∇_i are constant.

The quotient sheaf $\mathfrak{m} = \bigoplus_{i < 0} g_i$, $g_i = \Delta_{-i}/\Delta_{-i-1}$ has a natural structure of the graded nilpotent Lie algebra (GNLA) at any point $x \in M$. The bracket on \mathfrak{m} is induced by the commutator of vector fields on M . Δ is called *strongly regular* if the GNLA $\mathfrak{m}(x)$ does not depend on $x \in M$ (but we do not impose this property a priori).

The *growth vector* of $\Delta = g_{-1}$ is the sequence of dimensions $(\dim g_{-1}, \dim g_{-2}, \dots)$. (In other sources, the growth vector is one of the sequences $(\dim \Delta_1, \dim \Delta_2, \dots)$ or $(\dim \nabla_1, \dim \nabla_2, \dots)$ but, as our main tool is the Tanaka theory, we choose this convention).

The Tanaka prolongation $\mathfrak{g} = \hat{\mathfrak{m}}$ is the GNLA with negative graded part \mathfrak{m} , whose non-negative part is defined successively by the rule [T, Y]: g_k consists of all $u \in \bigoplus_{i < 0} g_{k+i} \otimes g_i^*$, acting as derivations: $u([X, Y]) = [u(X), Y] + [X, u(Y)]$, $X, Y \in \mathfrak{m}$. Notice that since \mathfrak{m} is fundamental, i.e. g_{-1} generates the whole GNLA \mathfrak{m} , the grading k homomorphism u is uniquely determined by the restriction $u : g_{-1} \rightarrow g_{k-1}$.

The space $\mathfrak{g} = \bigoplus g_i$ is naturally a graded Lie algebra, which we will call the *Tanaka algebra* of Δ (it depends on the point $x \in M$, unless Δ is strongly regular). By contrast, the GNLA $\mathfrak{m} = \bigoplus_{i < 0} g_i$ will be called the *Carnot algebra* of Δ (it is also called Levi-Tanaka or symbol algebra in the literature).

We wish to indicate that $\mathfrak{g} = \hat{\mathfrak{m}}$ can alternatively be defined via right extensions for modules, just as right extensions² for Lie algebras are governed by $H^1(\mathfrak{m}, \mathfrak{m})$ [SL, F]. First notice that the cohomology H^i of GNLA is graded. We denote the grading by a subscript $H^i = \bigoplus H_k^i$.

Then $g_0 = H_0^1(\mathfrak{m}, \mathfrak{m})$ and we define $\mathfrak{g}^0 = \mathfrak{m} \oplus g_0$. On the next step we get $g_1 = H_1^1(\mathfrak{m}, \mathfrak{g}^0)$ and we define $\mathfrak{g}^1 = \mathfrak{g}^0 \oplus g_1$. This is equivalent to adding variables w_i to kill the cohomology classes $[\omega_i] \in H_1^1(\mathfrak{m}, \mathfrak{g}^0)$, that is, let $dw_i = \omega_i$. The \mathfrak{m} -action naturally extends to \mathfrak{g}^1 making it into a representation and we continue: $\mathfrak{g}^p = g_{-k} \oplus \dots \oplus g_p$ with $g_p = H_p^1(\mathfrak{m}, \mathfrak{g}^{p-1})$. The prolongation of \mathfrak{m} is $\mathfrak{g} = \lim \mathfrak{g}^p$.

The importance of the Tanaka algebra is in majorizing the symmetry algebra $\text{sym}(\Delta)$ of the distribution. The following is the Corollary of Theorem 8.4 from [T]:

Theorem 4. *Let (M, Δ) be a manifold equipped with distribution, \mathfrak{m} its Carnot GNLA and $\mathfrak{g} = \hat{\mathfrak{m}}$ the Tanaka algebra. Then*

$$\dim \text{sym}(\Delta) \leq \dim \mathfrak{g}.$$

Moreover, provided \mathfrak{g} is finite-dimensional, this inequality is an equality if and only if the distribution Δ is locally flat.

Some remarks concerning this theorem are of order. First, this theorem is formulated (and proved in [T]) for strongly regular systems, and

²Both fundamental extension constructions in the theory of Lie algebras play a role in this paper: right extensions are important for Tanaka prolongation, while central extensions appear in Section 5 in connection with integral extensions.

this will suffice for our purpose. But it remains true if we understand the right hand size as $\sup_x \dim \mathfrak{g}(x)$.

Second, the last part of the theorem is not formulated in [T] as we have, but it follows from the construction of an absolute parallelism on the prolongation manifold of the structure.

Finally, locally flat means that the structure functions of the absolute parallelism vanish. This is equivalent to the claim that the distribution is locally diffeomorphic to the standard distribution on the affine realization of Lie group corresponding to the GNLA \mathfrak{g} , see [T].

2. FINITE-DIMENSIONALITY OF THE SYMMETRY ALGEBRA

In this section we give a sufficient condition for $\dim \mathfrak{g}$, and hence $\dim \text{sym}(\Delta)$, to be finite in the case $\dim \Delta = 2$. The statement has been surprisingly unknown until now.

Notice that for a rank 2 non-holonomic distribution the growth vector always begins like $(2, 1, \dots)$ and that the first number after a sequence of 1's can be only 2.

Theorem 5. *Let Δ be a completely non-holonomic rank 2 distribution with growth vector $(2, 1, \dots, 1, 2, \dots)$. Then the algebra $\text{sym}(\Delta)$ is finite-dimensional.*

Proof. Let $\mathfrak{g} = \oplus g_i$ be the Tanaka algebra of Δ evaluated at any point $x \in M$. In view of Theorem 4 it is enough to verify that $\dim \mathfrak{g} < \infty$.

To this end we use Corollary 2 of Theorem 11.1 from [T], which states that if the subalgebra

$$\mathfrak{h}_0 = \{v \in g_0 : [v, g_r] = 0 \ \forall r < -1\}$$

is zero, then \mathfrak{g} is finite-dimensional.³

Let $-k$ be the grading of the second two-dimensional space g_{-k} in \mathfrak{m} ($k > 3$). Choose a basis e_1, e'_1 of g_{-1} that generates basis elements $e_2 = [e_1, e'_1]$ of g_{-2} , $e_3 = [e_1, e_2]$ of g_{-3} , \dots , $e_{k-1} = [e_1, e_{k-2}]$ of g_{1-k} and $e_k = [e_1, e_{k-1}]$, $e'_k = [e'_1, e_{k-1}]$ of g_{-k} .

Let $v \in \mathfrak{h}_0$ and write $[v, e_1] = a e_1 + b e'_1$, $[v, e'_1] = c e_1 + d e'_1$. Then $[v, e_{k-1}] = 0$ implies $[v, e_k] = [a e_1 + b e'_1, e_{k-1}] = a e_k + b e'_k = 0$ and $[v, e'_k] = [c e_1 + d e'_1, e_{k-1}] = c e_k + d e'_k = 0$. Whence $a = b = c = d = 0$ and therefore $v = 0$. The claim follows. \square

Consider the growth vector of Δ . If it starts $(2, 1, 2, \dots)$ then, by Theorem 5, the symmetry algebra is finite-dimensional. The other

³The meaning of the condition $\mathfrak{h}_0 = 0$ is this: the complex characteristic variety of the Lie equation for $\text{sym}(\Delta)$ is empty.

possibility, namely $(2, 1, 1, \dots)$, has a clear geometric meaning which we provide in the next theorem.

Recall that the (geometric) prolongation of a rank 2 distribution $\bar{\Delta}$ on \bar{M} is (we consider only the local picture) the rank 2 distribution $\Delta = \langle \cos t \cdot U + \sin t \cdot V, \partial_t \rangle$ on $M = \bar{M} \times S^1$, where U, V are generators of $\bar{\Delta}$ and t the coordinate on S^1 .

If the distribution $\bar{\Delta}$ has preferred (vertical) section $V \in \Gamma(\bar{\Delta})$, then the (affine) prolongation $\Delta = \bar{\Delta}^{(1)}$ is simply $\Delta = \langle U + tV, \partial_t \rangle$ on $M = \bar{M} \times \mathbb{R}^1$, where t is the coordinate on \mathbb{R}^1 . This coincides with (Spencer) prolongation in the geometric theory of PDEs ([KLV, MZ]).

The reverse operation is not generally possible but, in the case when $\Delta = \bar{\Delta}^{(1)}$, we shall call the operation $\Delta \mapsto \bar{\Delta}$ de-prolongation.

Theorem 6. *Let Δ be a rank 2 non-holonomic distribution with the derived rank 3 distribution $\Delta_2 = [\Delta, \Delta]$. The growth vector of Δ is $(2, 1, 1, \dots)$ if and only if there exists a vector field $\xi \in \Gamma(\Delta)$ which is a Cauchy characteristic for the distribution Δ_2 . In this, and only in this case, Δ can be locally de-prolonged.*

Recall that the Cauchy characteristic ξ is a symmetry of the distribution Δ and its derived flag. So de-prolongation in this case is simply the quotient of the first derived distribution by the Cauchy characteristic

$$(\bar{M}, \bar{\Delta}) = (M, \Delta_2) / \xi.$$

The above theorem is due to E. Cartan [C2]. We present a new proof here for the convenience of the reader.

Proof. The claim about the existence of a Cauchy characteristic is obvious. We shall only demonstrate that $\bar{\Delta}^{(1)} = \Delta$.

For this, let us write M locally as the product $\mathbb{R}_x^{n-1} \times \mathbb{R}_t^1$ in such a way that $\xi = \partial_t$. We take the other generator X of Δ from $T\mathbb{R}_x^{n-1}$ with coefficient dependent on (x, t) . The derived flag equals

$$\Delta_1 = \Delta = \langle X, \partial_t \rangle, \quad \Delta_2 = \langle X, \dot{X}, \partial_t \rangle, \quad \Delta_3 = \langle X, \dot{X}, \ddot{X}, [X, \dot{X}], \partial_t \rangle,$$

where dot denotes the differentiation by t (or equivalently, the Lie derivative along ∂_t). Since the growth vector is $(2, 1, 1, \dots)$ and ∂_t is the Cauchy characteristic, we must have

$$\ddot{X} = \alpha X + \beta \dot{X}$$

for some t -dependent functions $\alpha, \beta \in C^\infty(\bar{M} \times \mathbb{R})$. Let $W = uX + v\dot{X}$ with $u, v \in C^\infty(\bar{M} \times \mathbb{R})$. Then solution to the system

$$\dot{W} = (\dot{u} + \alpha v)X + (u + \dot{v} + \beta v)\dot{X} = 0$$

is equivalent (provided $\text{rank}(X, \dot{X}) = 2$) to the ODE system

$$\frac{d}{dt} \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 0 & -\alpha \\ -1 & -\beta \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}.$$

By choosing a basis of fundamental solutions, we get two t -independent vector fields U, V such that $\langle U, V \rangle = \langle X, \dot{X} \rangle$. Since generators are defined up to scaling, we can assume $X = U + bV$. Regularity implies $\dot{b} \neq 0$. By reparametrizing t we can assume that $b = t$. Thus we have:

$$\Delta = \langle U + tV, \partial_t \rangle, \quad \Delta_2 = \langle U, V, \partial_t \rangle, \quad \bar{\Delta} = \langle U, V \rangle,$$

so that the claim follows from the definition of the prolongation. \square

De-prolongation does not just result in the removal of 1 from the growth vector at the first component and shift. It can significantly change the weak derived flag.

Example. Consider the Hilbert-Cartan equation $\mathcal{E}_{1,2}$. It has growth vector $(2, 1, 2)$ and the Tanaka algebra is the exceptional Lie algebra \mathfrak{g}_2 with grading (the number in the sequence below means the dimension while the subscript - the grading):

$$(2_{(-3)}, 1_{(-2)}, 2_{(-1)}, 4_{(0)}, 2_{(1)}, 1_{(2)}, 2_{(3)}).$$

The prolongation of $\mathcal{E}_{1,2}$ is a 6-dimensional manifold with rank 2 distribution of growth $(2, 1, 1, 1, 1)$. Here we see that de-prolongation is not just removal of 1 and shift.

The Tanaka algebra of this 6-dimensional model is still \mathfrak{g}_2 , but now with the grading

$$(1_{(-5)}, 1_{(-4)}, 1_{(-3)}, 1_{(-2)}, 2_{(-1)}, 2_{(0)}, 2_{(1)}, 1_{(2)}, 1_{(3)}, 1_{(4)}, 1_{(5)}).$$

The next prolongation has growth vector $(2, 1, 1, 1, 1, 1)$, and at this point the Tanaka prolongation is the contact (infinite-dimensional) algebra. This means that the 2nd (Spencer) prolongation of $\mathcal{E}_{1,2}$ is not Tanaka-flat.

Since the algebras $\text{sym}(\bar{\Delta})$ and $\text{sym}(\bar{\Delta}^{(1)})$ are isomorphic, Theorems 6 and 5 imply the following important statement (it can be also deduced in a non-trivial way from Theorem 7.1 of [MT]).

Corollary 2. *A regular germ of a rank 2 distribution Δ has infinite-dimensional symmetry algebra only in one of the following cases.*

- (1) Δ completely non-holonomic: then it is equivalent to the Cartan distribution \mathcal{C}_k on jet-space $J^k(\mathbb{R}, \mathbb{R})$, with $k = \dim M - 2$.
- (2) $\Delta_i \neq TM$ for any i : then there exists a codimension 1 foliation \mathcal{F} of M with $T\mathcal{F} \supset \Delta$ such that all leaves (\mathcal{F}, Δ) are equivalent.

Recall that distributions Δ with strong growth vector $(2, 1, 1, \dots, 1)$ are called Goursat distributions [G]. Their regular points can be characterized by the condition that the growth vector is $(2, 1, 1, \dots, 1)$ in both the weak and the strong sense. Near such points the normal form

$$\mathcal{C}_k = \text{Ann}\{dy_i - y_{i+1} dx \mid 0 \leq i < k\} \subset TJ^k(\mathbb{R}^1, \mathbb{R}^1)$$

is provided by the von Weber - Cartan theorem [W, C2], see also [GKR].

This concludes our discussion of distributions with growth vector $(2, 1, 1, \dots)$. From this point forward we assume that the distribution is not de-prolongable, i.e. its growth is $(2, 1, 2, \dots)$.

3. MOST SYMMETRIC RANK 2 DISTRIBUTIONS

In this section we study the Tanaka algebra of a single Monge ODE and prove our main result.

3.1. Calculation of the Carnot algebra. Let us consider equation (3) with fixed $m \leq n$ as a submanifold in the space of jets

$$\mathcal{E} \subset J^{m,n}(\mathbb{R}, \mathbb{R}^2) \simeq \mathbb{R}^{n+m+3}(x, y, \dots, y_m, z, z_1, \dots, z_n),$$

given as a hypersurface $y_m = F(x, y, \dots, y_{m-1}, z, z_1, \dots, z_n)$. Internally $\mathcal{E} \simeq \mathbb{R}^{n+m+2}(x, y, \dots, y_{m-1}, z, z_1, \dots, z_n)$ is canonically equipped with the rank 2 distribution

$$\mathcal{C} = \langle \mathcal{D}_x = \partial_x + y_1 \partial_y + \dots + F \partial_{y_{m-1}} + z_1 \partial_z + \dots + z_n \partial_{z_{n-1}}, \partial_{z_n} \rangle.$$

Its (weak) derived flag has growth vector $(2, 1, 2, \dots)$ if and only if

$$F_{z_n z_n} \neq 0.$$

In what follows we will assume this *non-degeneracy condition* which, according to Theorem 5, implies $\dim \text{sym}(\mathcal{C}) < \infty$.

In this case the derived distribution is $\mathcal{C}_2 = \mathcal{C} + \langle \partial_{z_{n-1}} + F_{z_n} \partial_{y_{m-1}} \rangle$ and the second derived is

$$\mathcal{C}_3 = \langle \mathcal{D}_x, \partial_{y_{m-1}}, \partial_{z_n}, \partial_{z_{n-1}}, \partial_{z_{n-2}} + F_{z_n} \partial_{y_{m-2}} \rangle.$$

The next (weak) derived⁴ are:

$$\begin{aligned} \mathcal{C}_{i+1} &= \langle \mathcal{D}_x, \partial_{y_{m-1}}, \dots, \partial_{y_{m-i+1}}, \partial_{z_n}, \dots, \partial_{z_{n-i+1}}, \partial_{z_{n-i}} + F_{z_n} \partial_{y_{m-i}} \rangle, \\ \mathcal{C}_{j+1} &= \langle \mathcal{D}_x, \partial_{y_{m-1}}, \dots, \partial_y, \partial_{z_n}, \dots, \partial_{z_{n-j}} \rangle, \quad i \leq m < j. \end{aligned}$$

⁴These formulas imply that the weak and strong flags of \mathcal{C} coincide, as was mentioned in Remark 1; this plays a role in Section 4.

Thus the dimension grows by two from grading -3 till grading $-m-2$ (resp. $-m-1$ if $n = m$) and then grows by one. The growth vector is:

$$(2, 1, \underbrace{2, \dots, 2}_m, \underbrace{1, \dots, 1}_{n-m-1}) \quad \text{for } m < n$$

and

$$(2, 1, \underbrace{2, \dots, 2}_{n-1}, 1) \quad \text{for } m = n.$$

Proposition 7. *Carnot algebra of equation (3) does not depend on a point $x \in \mathcal{E}$ and is equal to the Lie algebra $\mathfrak{f}_{m,n}$ given by the following table of brackets (only upper-diagonal non-zero entries are shown). For $m < n - 1$:*

	e_1	e'_1	e_2	e_3	e'_3	e_4	e'_4	\dots	e_{m+2}	e'_{m+2}	e_{m+3}	e_{m+4}	\dots	e_{n+1}
e_1		e_2	e_3	e_4	e'_4	e_5	e'_5	\dots	e_{m+3}		e_{m+4}	\dots	e_{n+1}	
e'_1			e'_3	e'_4		e'_5	\dots							
e_2														
\vdots														

The GNLA structure is given by the rule: $\deg e_i = -i = \deg e'_i$.

For $m = n$ the structure equations change to:

	e_1	e'_1	e_2	e_3	e'_3	e_4	e'_4	\dots	e_m	e'_m	e_{m+1}	e'_{m+1}	e'_{m+2}
e_1		e_2	e_3	e_4	e'_4	e_5	e'_5	\dots	e_{m+1}	e'_{m+1}		e'_{m+2}	
e'_1			e'_3	e'_4		e'_5	\dots	e'_{m+1}		e'_{m+2}			
e_2													
\vdots													

And for $n = m + 1$ the table is:

	e_1	e'_1	e_2	e_3	e'_3	e_4	e'_4	\dots	e_{m+1}	e'_{m+1}	e_{m+2}	e'_{m+2}
e_1		e_2	e_3	e_4	e'_4	e_5	e'_5	\dots	e_{m+2}	e'_{m+2}		
e'_1			e'_3	e'_4		e'_5	\dots	e'_{m+2}				
e_2												
\vdots												

In other words, $\mathfrak{f}_{m,n}$ is obtained through m successive 1D central extensions (this has sense to be called one central extension of order m) of the Heisenberg algebra $\mathcal{H}_{n+2} = \langle e_1, e'_1, e_2, e_3, \dots, e_{n+1} \rangle$.

Proof. The generators are given by the formulae (we suppose $F_{z_n z_n} \neq 0$; we use the same symbol for the vector and its image under quotient to

GNLA)

$$\begin{aligned} e_1 &= -\mathcal{D}_x, \quad e'_1 = \partial_{z_n}, \quad e'_i = F_{z_n z_n} \partial_{y_{m-i+2}} \quad (2 < i \leq m+2), \\ e_j &= \partial_{z_{n-j+1}} + F_{z_n} \partial_{y_{m-j+1}} + c_j \partial_{y_{m-j+2}} \quad (2 \leq j \leq m+2), \\ e_k &= \partial_{z_{n-k+1}} \quad (m+2 < k \leq n+1) \end{aligned}$$

with coefficients $c_2 = 0$, $c_j = F_{z_{n-1}} + F_{z_n} F_{y_{m-1}} - (j-2)\mathcal{D}_x(F_{z_n})$ ($3 \leq j \leq m+2$) playing no role in what follows. For $n = m+1$ the last line in the above formula is void, and for $n = m$ we shall additionally modify the range of the middle line ($2 \leq j \leq m+1$). The structure equations follow. \square

3.2. Calculation of the Tanaka algebra. Now let us describe the Tanaka algebra of equation (3) with $F_{z_n z_n} \neq 0$.

Proposition 8. *Prolongation $\mathfrak{t}_{m,n} = \widehat{\mathfrak{f}_{m,n}}$ of the GNLA $\mathfrak{f}_{m,n}$ has the following graded structure (the number means the dimension while the subscript - the grading):*

$$(1_{(-m-2)}, 2_{(-m-1)} \dots, 2_{(-4)}, 2_{(-3)}, 1_{(-2)}, 2_{(-1)}, 2_{(0)}) \quad \text{for } n = m;$$

then for $n > m > 1$:

$$(1_{(-n-1)} \dots 1_{(-m-3)}, 2_{(-m-2)} \dots 2_{(-3)}, 1_{(-2)}, 2_{(-1)}, 3_{(0)}, 1_{(1)} \dots 1_{(n-m-1)});$$

and for $n > 2$, $m = 1$ we get:

$$(1_{(-n-1)}, \dots, 1_{(-4)}, 2_{(-3)}, 1_{(-2)}, 2_{(-1)}, 3_{(0)}, 2_{(1)}, 1_{(2)}, \dots, 1_{(n-2)}).$$

Let us notice that for $m = 1, n = 2$ the prolongation is different. Indeed, in this case $\widehat{\mathfrak{f}_{1,2}}$ is isomorphic to the Lie algebra \mathfrak{g}_2 with the grading

$$(2_{(-3)}, 1_{(-2)}, 2_{(-1)}, 4_{(0)}, 2_{(1)}, 1_{(2)}, 2_{(3)}).$$

In the case $m = n = 1$ the prolongation is infinite, corresponding to the contact algebra of symmetries of $J^1(\mathbb{R}, \mathbb{R})$:

$$(1_{(-2)}, 2_{(-1)}, 4_{(0)}, 6_{(1)}, 9_{(2)}, 12_{(3)}, 16_{(4)}, 20_{(5)}, 25_{(6)}, 30_{(7)}, 36_{(8)} \dots).$$

To be more precise the GNLA structure of $\mathfrak{t}_{m,n} = \oplus g_i$ is given by the structure equations of Proposition 7 coupled with the following.

Proposition 9. *For $n = m$: $\mathfrak{t}_{m,m} = \mathfrak{f}_{m,m} \oplus g_0$, with the space g_0 Abelian and generated by e_0^{10}, e_0^{01} given by*

$$\begin{aligned} [e_0^{ac}, e_1] &= a e_1, \quad [e_0^{ac}, e'_1] = c e'_1, \\ [e_0^{ac}, e_k] &= ((k-1)a + c) e_k, \quad k \geq 2, \\ [e_0^{ac}, e'_k] &= ((k-2)a + 2c) e'_k, \quad k > 2. \end{aligned}$$

For $n > m > 1$ the space $g_0 \subset \mathfrak{t}_{m,n}$ is a Borel subalgebra of $\mathfrak{gl}(g_{-1})$ and is generated by $e_0^{100}, e_0^{010}, e_0^{001}$ given by

$$\begin{aligned} [e_0^{abc}, e_1] &= a e_1 + b e'_1, \quad [e_0^{abc}, e'_1] = c e'_1, \\ [e_0^{abc}, e_k] &= ((k-1)a + c) e_k + (k-2)b e'_k, \quad k \geq 2, \\ [e_0^{abc}, e'_k] &= ((k-2)a + 2c) e'_k, \quad k > 2. \end{aligned}$$

Further on, $\mathfrak{t}_{m,n} = \mathfrak{f}_{m,n} \oplus g_0 \oplus \cdots \oplus g_{n-m-1}$, where g_i for $i > 0$ are one-dimensional and generated by e_{-i} given by (only non-trivial relations are shown, and we omit commutators of non-negative gradations⁵):

$$\begin{aligned} [e_{-1}, e_1] &= e_0^{010}, \quad [e_{-1}, e_k] = \frac{(k-2)(k-3)}{2} e'_{k-1}, \\ [e_{-2}, e_1] &= e_{-1}, \quad [e_{-2}, e_k] = \frac{(k-2)(k-3)(k-4)}{6} e'_{k-2}, \quad \dots \\ \dots \quad [e_{m+1-n}, e_1] &= e_{m+2-n}, \quad [e_{m+1-n}, e_k] = \binom{k-2}{n-m} e'_{k+m+1-n} \end{aligned}$$

(in the above formulae for $[e_{-p}, e_k]$ we restrict $p+3 \leq k \leq m+p+2$).

When $n > 2$, $m = 1$, the above formulae hold true with the only difference that g_1 is two-dimensional and is generated by e_{-1}^{10}, e_{-1}^{01} , so that we change e_{-1} to e_{-1}^{10} in the above, and add the only nontrivial relations

$$\begin{aligned} [e_{-1}^{01}, e_1] &= e_0^{-2,0,1} = e_0^{001} - 2 e_0^{100}, \quad [e_{-1}^{01}, e'_1] = (1 - n^2) e_0^{010}, \\ [e_{-1}^{01}, e_k] &= (n^2 - k^2 + 2k - 4) e_{k-1} \end{aligned}$$

for the second generator of g_1 .

Proof. Consider at first the case $n = m$. Then g_{-1} canonically splits as $\mathbb{R}e_1 \oplus \mathbb{R}e'_1$, where e_1 is given (up to scaling) by the relation $[e_1, e_{m+1}] = 0$ (with $e_{m+1} = \text{ad}_{e_1}^{m-1}(e_2)$) and e'_1 by the relation $[e'_1, e'_3] = 0$ (with $e'_3 = \text{ad}_{e'_1}(e_2)$). Thus g_0 is diagonal, and in fact it equals the space $\{e_0^{ac}\}$ from the display formula of the Proposition.

To see that $g_1 = 0$, consider its element e_{-1} . It is uniquely given by 4 parameters:

$$[e_{-1}, e_1] = e_0^{ac}, \quad [e_{-1}, e'_1] = e_0^{a'c'}.$$

This implies the further relations:

$$\begin{aligned} [e_{-1}, e_2] &= c e'_1 - a' e_1, \quad [e_{-1}, e_3] = (a + 2c) e_2, \quad [e_{-1}, e'_3] = (2a' + c') e_2, \\ [e_{-1}, e_4] &= (3a + 3c) e_3, \quad [e_{-1}, e'_4] = (2a' + c') e_3 + (a + 2c) e'_3, \\ [e_{-1}, e_5] &= (6a + 4c) e_4, \quad \dots \end{aligned}$$

Relation $0 = [e_{-1}, [e'_1, e'_3]] = (3a' + 3c') e'_3$ yields $c' = -a'$.

⁵which restore uniquely by the derivation property.

There are two ways to evaluate $[e_{-1}, e'_5]$ since $e'_5 = [e_1, e'_4] = [e'_1, e_4]$ and they do not match, implying $a' = c' = 0$, $c = 0$. We continue calculation and get

$$[e_{-1}, e_k] = \frac{(k-1)(k-2)}{2} a e_{k-1}, \quad [e_{-1}, e'_k] = \frac{(k-2)(k-3)}{2} a e'_{k-1}.$$

For $k = m + 2$ (writing $[e_{-1}, e_{m+1}] = 0$ instead of e_{m+2}) we get $a = 0$.

Let now $n > m > 1$. Then only the line $\mathbb{R}e'_1 \subset g_{-1}$ is canonical, and it's easy to check that g_0 is upper triangular in the basis e_1, e'_1 , with general element e_0^{abc} as in the display formula in the Proposition.

Again to calculate g_1 we describe its element e_{-1} via the formulae

$$[e_{-1}, e_1] = e_0^{abc}, \quad [e_{-1}, e'_1] = e_0^{a'b'c'},$$

which imply

$$\begin{aligned} [e_{-1}, e_2] &= (c - b') e'_1 - a' e_1, \\ [e_{-1}, e_3] &= (a + 2c - b') e_2, \quad [e_{-1}, e'_3] = (2a' + c') e_2, \\ [e_{-1}, e_4] &= (3a + 3c - b') e_3 + b e'_3, \quad [e_{-1}, e'_4] = (2a' + c') e_3 + (a + 2c) e'_3, \\ [e_{-1}, e_5] &= (6a + 4c - b') e_4 + 3b e'_4, \quad \dots \end{aligned}$$

Next $0 = [e_{-1}, [e'_1, e'_3]] = (3a' + 3c') e'_3$ yields $c' = -a'$. Evaluation of $[e_{-1}, e'_5]$ in two ways using $e'_5 = [e_1, e'_4] = [e'_1, e_4]$ gives two different results, whence $a' = c' = 0$, $b' = c$.

The further calculations go without contradictions with the result

$$\begin{aligned} [e_{-1}, e_k] &= \left(\frac{(k-1)(k-2)}{2} a + (k-2)c\right) e_{k-1} + \frac{(k-2)(k-3)}{2} b e'_{k-1}, \\ [e_{-1}, e'_k] &= \left(\frac{(k-2)(k-3)}{2} a + 2(k-3)c\right) e'_{k-1}. \end{aligned}$$

Now we need to specify and consider first $n = m + 1$. Substituting the last expression into $[e_{-1}, [e_1, e_{m+2}]] = 0$ and $[e_{-1}, [e_1, e'_{m+2}]] = 0$ we get 3 independent constraints implying $a = b = c = 0$. Thus in this case $g_1 = 0$.

Let now $n > m + 1$. Then the relations are $[e_{-1}, [e_1, e_{n+1}]] = 0$ and $[e_{-1}, [e_1, e'_{m+2}]] = 0$ implying only $a = c = 0$, thus making e_{-1} with $b = 1$ the generator of g_1 .

Next we consider g_2 with element e_{-2} given by

$$[e_{-2}, e_1] = \alpha e_{-1}, \quad [e_{-2}, e'_1] = \alpha' e_{-1}.$$

This yields

$$\begin{aligned} [e_{-2}, e_2] &= -\alpha' e_0^{010}, \quad [e_{-2}, e_3] = \alpha' e'_1, \quad [e_{-2}, e'_3] = 0, \\ [e_{-2}, e_4] &= \alpha' e_2, \quad [e_{-2}, e'_4] = 0, \quad [e_{-2}, e_5] = \alpha' e_3 + \alpha e'_3, \quad \dots \end{aligned}$$

Evaluation of $[e_{-2}, e'_5]$ via $e'_5 = [e_1, e'_4] = [e'_1, e_4]$ gives $2\alpha' e'_3$ and 0, whence $\alpha' = 0$. If $n > m + 2$ there are no more contradictions and we get

$$[e_{-2}, e_k] = \frac{(k-2)(k-3)(k-4)}{6} \alpha e'_{k-2}, \quad [e_{-2}, e'_k] = 0,$$

i.e. g_2 is one-dimensional.

For $n = m + 2$ the last relation in $\mathfrak{f}_{m,n}$ forces $\alpha = 0$ and thus $g_2 = 0$ stops the prolongation. Continuing in the same way we uncover the result for $n > m > 1$.

The case $n > 2, m = 1$ is considered in the same manner giving us the required prolongation $\widehat{\mathfrak{f}}_{1,n}$. Indeed, the $g_0 = \{e_0^{abc}\}$ is the same with relations (δ_i^j is the Kronecker delta)

$$\begin{aligned} [e_0^{abc}, e_1] &= a e_1 + b e'_1, \quad [e_0^{abc}, e'_1] = c e'_1, \\ [e_0^{abc}, e_k] &= ((k-1)a + c) e_k + \delta_k^3 b e'_3 \quad (k \geq 2), \quad [e_0^{abc}, e'_3] = (a + 2c) e'_3. \end{aligned}$$

Next is g_1 the elements of which are specified by the relations

$$[e_{-1}, e_1] = e_0^{abc}, \quad [e_{-1}, e'_1] = e_0^{a'b'c'}.$$

Similar to the above we obtain $a' = c' = 0, a = -2c, b' = (1 - n^2)c$. Thus the general element is e_{-1}^{bc} given by the only nontrivial relations:

$$\begin{aligned} [e_{-1}^{bc}, e_1] &= e_0^{-2c,b,c}, \quad [e_{-1}^{bc}, e'_1] = e_0^{0,(1-n^2)c,0}, \\ [e_{-1}^{bc}, e_k] &= ((k-1)(3-k) + n^2 - 1)c e_{k-1} + \delta_k^4 b e'_3. \end{aligned}$$

Next comes g_2 and the same arguments show that it is one-dimensional. Continuing we get $g_p = \langle e_{-p} \rangle$ for $2 \leq p \leq n-2$ with the only non-trivial relations (between elements of positive and negative gradings):

$$[e_{-p}, e_1] = e_{1-p}, \quad [e_{-p}, e_{p+3}] = e'_3$$

(and we mean e_{-1}^{10} instead of e_{-1} in the very first formula for $p = 2$).

Prolongation of $\mathfrak{f}_{1,1}$ is rather standard, and the case of $\mathfrak{f}_{1,2}$ is due to Tanaka [T] (see also [Y]). \square

Corollary 3. $\dim \mathfrak{t}_{m,n} = 2n + 4$ for $n \geq m > 1$ and it equals $2n + 5$ for $n > 2, m = 1$. In exceptional cases $\dim \mathfrak{t}_{1,2} = 14$ and $\dim \mathfrak{t}_{1,1} = \infty$.

Thus in non-exceptional cases $\dim \widehat{\mathfrak{f}}_{m,n} \leq 2 \dim \mathfrak{f}_{m,n} - 1$ with the equality only for $m = 1$ (the deviation being $2m - 1$ for $m > 1$).

3.3. Symmetric models. In this section we shall calculate the internal infinitesimal symmetry algebra $\text{sym}(\mathcal{E}_{m,n})$ for the model equation (4). The upper bound on the number of independent symmetries is

given by Corollary 3, and we shall show that this upper bound is realized by an explicit construction. An isomorphism between $\widehat{\mathfrak{f}}_{m,n}$ and $\text{sym}(\mathcal{E}_{m,n})$ together with algebraic description of the latter will be given.

If $m > 1$, the symmetry algebra has dimension $\dim \text{sym}(\mathcal{E}_{m,n}) = 2n + 4$. These symmetries split into two types. The first type are the point symmetries with the generators: shifts

$$S_0 = \partial_x, \quad Y_i = \frac{x^i}{i!} \partial_y \quad (0 \leq i < m), \quad Z_j = \frac{x^j}{j!} \partial_z \quad (0 \leq j < n)$$

and scalings

$$S_1 = x \partial_x + (m-1)y \partial_y + (n - \frac{1}{2})z \partial_z, \quad R = y \partial_y + \frac{1}{2}z \partial_z.$$

These form a subalgebra of symmetries of dimension $m + n + 3$, which acts transitively on $\mathcal{E}_{m,n}$ (of dimension $m + n + 2$). As we shall see in Proposition 11, the subalgebra $\langle S_0, S_1, Y_i, Z_j \rangle \subset \text{sym}(\mathcal{E}_{m,n})$ coincides with the Carnot algebra $\widehat{\mathfrak{f}}_{m,n}$, while the vector field $S_1 + 3R$ corresponds to the Euler vector⁶ $\epsilon \in \mathfrak{g}_0$.

Then there are higher symmetries with generators ($0 \leq k \leq n - m$):

$$Z_{n+k} = 2 \sum_{p=0}^k (-1)^p \binom{m+p-1}{p} \frac{x^{k-p}}{(k-p)!} z_{n-m-p} \partial_y + \frac{x^{n+k}}{(n+k)!} \partial_z$$

To verify that these are symmetries we prolong them to

$$\hat{Z}_{n+k} = 2 \sum_{i=0}^m \sum_{p=0}^k (-1)^p \binom{m+p-1-i}{p} \frac{x^{k-p}}{(k-p)!} z_{n-m-p+i} \partial_{y_i} + \sum_{j=0}^n \frac{x^{n+k-j}}{(n+k-j)!} \partial_{z_j}$$

(with $\binom{p-1}{p} = \delta_p^0$) and check that

$$\hat{Z}_{n+k}(y_m - z_n^2) = 0.$$

When $n = m$, Z_n is also a point symmetry⁷; otherwise Z_{n+k} are genuine higher symmetries.

Thus the totality of symmetries is $2n + 4$ and since by Theorem 4 $\dim \text{sym}(\mathcal{E}_{m,n}) \leq \dim \widehat{\mathfrak{f}}_{m,n} = 2n + 4$, there are no more continuous symmetries.

For $m = 1$, $n > 2$ there is one additional symmetry⁸, namely

$$S_2 = x^2 \partial_x + (nz_{n-1})^2 \partial_y + (2n-1)xz \partial_z,$$

and then we have⁹ $\dim \text{sym}(\mathcal{E}_{1,n}) = 2(n+3) - 1$. Thus we have proved:

⁶this is the unique vector characterized by the condition $[\epsilon, v] = k \cdot v \quad \forall v \in \mathfrak{g}_k$.

⁷this explains why this case separates in Proposition 7.

⁸for $n = 2$ this symmetry persists as well, but there are 5 more symmetries.

⁹with $\dim \mathcal{E}_{1,n} = n + 3$, so that $2(n+3) - 1 = \dim ST^*M$, cf. [DZ].

Proposition 10. *The algebra $\text{sym}(\mathcal{E}_{m,n})$ of internal symmetries of equation (4) has dimension $2n + 4$ for $n \geq m > 1$ and dimension $2n + 5$ for $n > 2, m = 1$ with the basis S_l, R, Y_i, Z_l .*

Let us describe the Lie algebra structure for $\text{sym}(\mathcal{E}_{m,n})$. Assume at first $m > 1$ and let the index $l = 0, \dots, 2n - m$. The non-zero commutators are the following (with $Y_{-1} = Z_{-1} = 0$).

$$\begin{aligned} [S_0, S_1] &= S_0, [S_0, Y_i] = Y_{i-1}, [S_0, Z_l] = Z_{l-1}, \\ [S_1, Y_i] &= (i + 1 - m)Y_i, [S_1, Z_l] = (l + \frac{1}{2} - n)Z_l, \\ [Y_i, R] &= Y_i, [Z_l, R] = \frac{1}{2}Z_l. \end{aligned}$$

These are accompanied by splitting Z_l into Z_j ($0 \leq j < n$) and Z_{n+k} ($0 \leq k \leq n - m$) and the relations:

$$[Z_j, Z_{n+k}] = 2(-1)^k \binom{n-j-1}{k} Y_{j-n+m+k}, \quad n - m - k \leq j < n - k.$$

For $m = 1$ (when Y_i are reduced to solely Y_0) we add the following commutators for S_2

$$[S_0, S_2] = 2S_1, [S_1, S_2] = S_2, [S_2, Z_l] = ((l + 1 - n)^2 - n^2)Z_{l+1}.$$

By Theorem 4 and Proposition 10 the symmetry algebra of our model is isomorphic to the Tanaka algebra $\mathfrak{t}_{m,n} = \widehat{\mathfrak{f}_{m,n}}$. We make this explicit.

Proposition 11. *The isomorphism $\text{sym}(\mathcal{E}_{m,n}) \simeq \mathfrak{t}_{m,n}$ is given by the following identifications:*

$$\begin{aligned} S_0 &= e_1, S_1 = \frac{1}{2}e_0^{001} - e_0^{100}, S_2 = -e_{-1}^{01}, R = -\frac{1}{2}e_0^{001}, \\ Y_i &= e'_{m+2-i} \quad (0 \leq i < m), Z_j = e_{n+1-j} \quad (0 \leq j < n), Z_n = -2e'_1, \\ Z_{n+1} &= 2e_0^{010}, Z_{n+2} = -2e_{-1}^{10}, Z_{n+k+1} = 2(-1)^k e_{-k} \quad (1 < k < n - m). \end{aligned}$$

For $m > 1$ we shall remove S_2 and e_{-1}^{01} with change $e_{-1}^{10} \mapsto e_{-1}$. And for $m = n$ we identify $e_0^{abc} \mapsto e_0^{ac}$ and remove e_0^{010} (Z_{n+1} is non-existent in this case).

In particular, we obtain the weights giving gradation of $\text{sym}(\mathcal{E}_{m,n})$:

$$w(S_k) = k - 1, w(R) = 0, w(Y_i) = i - m - 2, w(Z_j) = j - n - 1.$$

For $m = 1$, the radical is $\mathfrak{r} = \langle Y_0, Z_l, R \rangle_{0 \leq l \leq 2n-1}$ and the vectors S_0, S_1, S_2 define an sl_2 complement. The length of the derived series for \mathfrak{r} equals 3. The nilradical is $\langle Y_0, Z_l \rangle_{0 \leq l < 2n}$ and its class (the length of lower central series) equals 2.

For $m > 1$ the symmetry algebra is solvable. The length of the derived series equals 3 for $n = m$ and 4 for $n > m$. The nilradical is $\langle S_0, Y_i, Z_l \rangle_{0 \leq i < m; 0 \leq l \leq 2n-m}$ and has class equal $2n - m + 1$.

3.4. End of the proof and a generalization. Now we can prove our main results.

Proof of Theorem 1. Let Δ be the distribution corresponding to equation (3) with non-degeneracy condition $F_{z_n z_n} \neq 0$ satisfied and \mathcal{C} the one for equation (4). We have demonstrated inequality $\dim \text{sym}(\Delta) \leq \dim \text{sym}(\mathcal{C})$.

Moreover by virtue of Theorem 4 the equality

$$\dim \text{sym}(\Delta) = \dim \text{sym}(\mathcal{C}) = \dim \mathfrak{t}_{m,n}$$

implies $\text{sym}(\Delta) \simeq \mathfrak{t}_{m,n}$ and the distribution Δ of the equation \mathcal{E} is flat.

Therefore it is locally isomorphic to the standard model, which coincides by the above with the distribution \mathcal{C} . Thus the equation \mathcal{E} is locally internally equivalent to the equation $\mathcal{E}_{m,n}$ (4). \square

Consider Monge ODEs (3) with non-degeneracy condition $F_{z_n z_n} \neq 0$ (sufficient for $\dim \text{sym}(\Delta) < \infty$). They correspond internally to rank 2 distributions on a manifold of $\dim(\mathcal{E}) = m + n + 2 = k$. Denote the class of all such distributions with $m \geq \mu$ and fixed k by $\mathfrak{D}(\mu, k)$. Our calculations imply:

Corollary 4. *Among distributions in the class $\mathfrak{D}(m, k)$ the most symmetric are locally equivalent to the distribution of equation (4).*

This implies in turn our Corollary 1. Thus if we denote $\mathfrak{p}_{n+3} = \mathfrak{f}_{1,n}$, then the largest algebra of symmetries of distributions in $\mathfrak{D}(1, k)$ is $\widehat{\mathfrak{p}}_k$.

Remarks: 1. Any rank 2 distribution can be realized as the Cartan distribution of an underdetermined ODE system. By the same trick as in D-modules this can be rewritten as a single higher order ODE. The equation manifold has however higher dimension and the former distribution is obtained from the latter by a sequence of de-prolongations. Thus without non-degeneracy condition Monge equations are internally indistinguishable from the general rank 2 distributions.

2. Dimension 5 admits only one Lie algebra \mathfrak{p}_5 of the type $(2, 1, 2)$. In dimension 6 there are already 3 GNLA of $(2, 1, 2, 1)$ -type: $\mathfrak{p}_6 = \mathfrak{f}_{1,3}$, $\mathfrak{h}_6 = \mathfrak{f}_{2,2}$, and one more $\mathfrak{e}\mathfrak{l}_6$ not realizable as a distribution of a nondegenerate single Monge ODE [AK]. There is a reason to call them parabolic, hyperbolic and elliptic, whence our choice of the letters.

3. In higher dimensions the number of GNLA's grows unboundedly, and it is impossible to trace the classification. Study of low-dimensional cases and some other types of underdetermined ODEs leads to the following claim, which will be discussed in a forthcoming paper [AK]:

Conjecture: Equation $\mathcal{E}_{1,k-3}$ (4) represents the most symmetric distribution \mathcal{C} among all rank 2 distributions Δ on a manifold \mathcal{E} of $\dim(\mathcal{E}) = k$ with $\dim \text{sym}(\Delta) < \infty$.

4. LIE-BÄCKLUND TYPE THEOREM

Consider a bundle $\pi : E \rightarrow M$. A Lie transformation of the jet space $J^k\pi$ is, by definition, a diffeomorphism preserving the canonical Cartan distribution $\mathcal{C} = \mathcal{C}^k$ (k is not the rank, but we write it as a superscript to avoid confusions). The classical Lie-Bäcklund theorem states that a Lie transformation F of the jet space $J^k\pi$ is the lift of a Lie transformation f of $J^\epsilon\pi$, where $\epsilon = 1$ and f is a contact transformation of $J^1\pi$ in the case $\text{rank } \pi = 1$, while $\epsilon = 0$ and f is a (local) diffeomorphism of $J^0\pi$ (point transformation) for $\text{rank } \pi > 1$.

Various generalizations of this to Lie transformations of equations $\mathcal{E} \subset J^k\pi$ are known [KLV, AKO, GK]. In this section we present a generalization to the case of Monge equations (3).

4.1. Lie transformations of the mixed jets. To prove our main result we first consider the case of mixed jets¹⁰

$$J^{k_1, k_2}(M, N_1 \times N_2) = J^{k_1}(M, N_1) \times_M J^{k_2}(M, N_2)$$

with points $(x, j_x^{k_1}(s_1), j_x^{k_2}(s_2))$, where $j_x^{k_i}(s_i)$ denotes k_i -jet of the map $s_i : M \rightarrow N_i$ at the point $x \in M$. There are natural projections

$$\begin{aligned} \pi_i &: J^{k_1, k_2}(M, N_1 \times N_2) \rightarrow J^{k_i}(M, N_i) \quad (i = 1, 2), \\ \pi^j &: J^{k_1, k_2}(M, N_1 \times N_2) \rightarrow J^{k_1 - j, k_2 - j}(M, N_1 \times N_2). \end{aligned}$$

The Cartan distributions on the space of pure jets induce the one on the mixed jets by the formula

$$\mathcal{C}^{k_1, k_2} = (d\pi_1)^{-1}(\mathcal{C}^{k_1}) \cap (d\pi_2)^{-1}(\mathcal{C}^{k_2}) \subset TJ^{k_1, k_2}.$$

Its (weak=strong) derived flag is

$$(\mathcal{C}^{k_1, k_2})_j = (d\pi^j)^{-1}(\mathcal{C}^{k_1, k_2}).$$

Note that in the case considered in the next section, namely $\dim M = \dim N_1 = \dim N_2 = 1$, the growth vector of \mathcal{C}^{k_1, k_2} is $(3, 2, \dots, 2, 1, \dots, 1)$.

To exclude the known case $J^k(M, N_1 \times N_2) = J^{k, k}(M, N_1 \times N_2)$ of pure jets, we assume in the next statement that $k_1 < k_2$.

¹⁰The results of this subsection have a straightforward generalization to the case of more general mixed jets J^{k_1, \dots, k_s} .

Theorem 12. 1. Every Lie transformation F of the mixed jet space $J^{k_1, k_2}(M, N_1 \times N_2)$ is the prolongation $F_0^{(k_1)}$ of a Lie transformation F_0 of the space $J^{0, k_2 - k_1}(M, N_1 \times N_2)$.

2. Under identification $J^{0, k}(M, N_1 \times N_2) = N_1 \times J^k(M, N_2)$ (for $k = k_2 - k_1$) every Lie transformation F_0 is given by the formula

$$F_0(y, j_x^k(s)) = (Y(y, j_x^k(s)), f^{(k-\epsilon)}(j_x^k(s))),$$

where $\epsilon = 1$ and $f : J^1(M, N_2) \rightarrow J^1(M, N_2)$ is a contact transformation in the case $\dim N_2 = 1$, while $\epsilon = 0$ and $f^{(k)}$ is the lift of a point transformation $f : M \times N_2 \rightarrow M \times N_2$ for $\dim N_2 > 1$.

Proof. For 1 we mimic the proof of the classical Lie-Bäcklund theorem: from the derived flag of \mathcal{C}^{k_1, k_2} we deduce that the Cauchy characteristic space of $(\mathcal{C}^{k_1, k_2})_{j+1}$ is $\text{Ker}(d\pi^j)$ and thus every Lie transformation of J^{k_1, k_2} fibers over a Lie transformation of $J^{k_1 - j, k_2 - j}$ for any $1 \leq j \leq k_1$. Standard arguments show that the former transformation is the prolongation of the latter.

To prove 2 let us notice that the Cauchy characteristics space of the Cartan distribution $\mathcal{C}^{0, k}$ equals $TN_1 = \text{Ker}(d\pi_2)$. Thus every Lie transformation of $J^{0, k}(M, N_1 \times N_2)$ fibers over a Lie transformation of the pure jet space $J^k(M, N_2)$ and an application of the classical Lie-Bäcklund theorem finishes the proof. \square

4.2. Lie transformations of Monge ODEs. Internal equivalence of two equations $\mathcal{E}, \mathcal{E}'$ of type (3) is a diffeomorphism $\phi : \mathcal{E} \rightarrow \mathcal{E}'$ which maps the induced Cartan distributions $\mathcal{C}_{\mathcal{E}}$ to $\mathcal{C}_{\mathcal{E}'}$. Proposition 7 implies that two nondegenerate Monge ODEs with different pairs (m, n) are never internally equivalent.

External equivalence of two Monge equations $\mathcal{E}, \mathcal{E}' \subset J^{m, n}(\mathbb{R}, \mathbb{R} \times \mathbb{R})$ is a Lie transformation of $J^{m, n}(\mathbb{R}, \mathbb{R} \times \mathbb{R})$ mapping \mathcal{E} to \mathcal{E}' . In particular, it induces an internal equivalence.

Theorem 13. Every internal equivalence of two non-degenerate Monge equations (3) with $n \geq m > 0$ and $(m, n) \neq (1, 1), (1, 2)$ is induced by a unique external Lie transformation.

Contrary to other Lie-Bäcklund type theorems [KLV, GK], it is not enough here to study Cauchy characteristics of the derived distributions or characteristic subspaces. We shall use the Tanaka theory.

Proof. Let \mathcal{C} be the Cartan distribution of (3). Its (weak=strong) derived flag \mathcal{C}_i was calculated in Section 3.1.

According to the proof of Proposition 9, the Cartan distribution \mathcal{C} of (3) contains the canonical (from internal viewpoint) line $\mathcal{L} = \langle e'_1 \rangle$.

Namely the Carnot algebra $\mathfrak{f}_{m,n}$ contains precisely one vector $v = e'_1 \in \mathfrak{g}_{-1}$ (up to scaling) satisfying $\text{ad}_v^3 = 0$. From external viewpoint this line \mathcal{L} is generated by the vertical vector (in internal coordinates ∂_{z_n} , in external $\partial_{z_n} - F_{z_n} \partial_{y_m}$). The quotient of \mathcal{E} by the corresponding foliation can be naturally identified with $J^{m-1,n-1}(\mathbb{R}, \mathbb{R}^2)$. Indeed, if we denote by $j : \mathcal{E} \rightarrow J^{m,n}$ the inclusion and by $\pi : \mathcal{E} \rightarrow \mathcal{E}/\mathcal{L}$ the projection, then for $\pi^1 : J^{m,n} \rightarrow J^{m-1,n-1}$ we have: $\pi = \pi^1 \circ j$.

Furthermore, the distribution $\mathcal{H} = \langle \mathcal{D}_x, \partial_{z_n}, \partial_{z_{n-1}}, \partial_{y_{m-1}} \rangle \subset \mathcal{C}_3$ is also canonical (intrinsically defined) as it equals $\mathcal{C}_2 \oplus [e'_1, \mathcal{C}_2]$ (in the Carnot algebra it corresponds to $\mathfrak{g}_{-1} \oplus \mathfrak{g}_{-2} \oplus \langle e'_3 \rangle$). In the quotient of \mathcal{E} by \mathcal{L} the distribution \mathcal{H} goes to the Cartan distribution of the jet space, $d\pi(\mathcal{H}) = \mathcal{C}^{m-1,n-1}$.

Consider two Monge ODEs $\mathcal{E}, \mathcal{E}' \subset J^{m,n}(\mathbb{R}, \mathbb{R}^2)$ and an internal equivalence $\Psi : \mathcal{E} \rightarrow \mathcal{E}'$. It satisfies $\Psi_*(\mathcal{L}) = \mathcal{L}'$ and $\Psi_*(\mathcal{H}) = \mathcal{H}'$ and thus induces a Lie transformation Φ of the jet-space, i.e. we have the following commutative diagram:

$$\begin{array}{ccc} \mathcal{E} & \xrightarrow{\Psi} & \mathcal{E}' \\ \pi \downarrow & & \downarrow \pi' \\ J^{m-1,n-1}(\mathbb{R}, \mathbb{R}^2) & \xrightarrow{\Phi} & J^{m-1,n-1}(\mathbb{R}, \mathbb{R}^2). \end{array}$$

By Theorem 12, Φ equals the prolongation of a Lie transformation ϕ of $J^{0,n-m}(\mathbb{R}, \mathbb{R}^2)$. To complete the proof we must show that $\Psi = \phi^{(n-m)}|_{\mathcal{E}}$.

First, we claim that $\phi^{(n-m)}$ maps \mathcal{E} to \mathcal{E}' . Indeed, a point $b \in \mathcal{E}$ is represented by the class of tangent spaces $T_{\pi(b)}(\pi(L^1))$, where L^1 runs through integral curves of $(\mathcal{E}, \mathcal{C})$ passing through b . Thus $\Psi(b)$ is determined by the collection of spaces $\Phi_* T_{\pi(b)}(\pi(L^1))$, which coincides with

$$T_{\Phi(\pi(b))}(\phi^{(n-m-1)}(\pi(L^1))) = T_{\pi'(\phi^{(n-m)}(b))}(\pi'(\phi^{(n-m)}(L^1)))$$

due to equality $\Phi = \phi^{(n-m-1)}$. This implies that $\phi^{(n-m)}(\mathcal{E}) = \mathcal{E}'$, and let us denote

$$I = \Psi^{-1} \circ \phi^{(n-m)} : \mathcal{E} \rightarrow \mathcal{E}.$$

This map is a Lie transformation fibered over the identity map of $J^{m-1,n-1}(\mathbb{R}, \mathbb{R}^2)$. Writing conservation of the Cartan form $dz_{n-1} - z_n dx$ we obtain that I is indeed the identity. \square

Remark 2. *The exact formulae from Section 3.3 show that symmetries of (4) are induced by exterior transformations of jets $J^{m,n}$ as the theorem claims. They have block form generalizing the feedback transformations in control theory (with variable z , control y and time x).*

Let us discuss the exceptional cases. When $m = n = 1$, the internal symmetry group of $\mathcal{E} : y_1 = f(x, y, z, z_1)$ is isomorphic to the contact group of $J^1(\mathbb{R}, \mathbb{R})$ and is parametrized by a single function in 3 variables. On the other hand, the external symmetries can be calculated directly since, by the classical Lie-Bäcklund theorem, they are prolongations of point symmetries of $J^0(\mathbb{R}, \mathbb{R}^2)$. In particular, for the model equation $\mathcal{E}_{1,1} : y' = (z')^2$ the group of point symmetries is 10-dimensional, and thus is much smaller than the infinite-dimensional internal group.

When $m = 1, n = 2$, the internal geometry of the equation $\mathcal{E} : y_1 = f(x, y, z, z_1, z_2)$ was described by E. Cartan [C1]. The internal symmetry group is at most 14-dimensional, and is realized by the equation $\mathcal{E}_{1,2} : y' = (z'')^2$ (see [AKO, AF] for the explicit generators). The symmetry group is the split real form of the exceptional Lie group G_2 . This group has a 9-dimensional subgroup of external symmetries coming from Lie transformations of $J^{1,2}(\mathbb{R}, \mathbb{R}^2)$. But the remaining 5 generators are not induced by the external symmetries. Thus, again in this case, the internal geometry is richer than the external geometry.

5. INTEGRABLE EXTENSIONS

In naive terms, a covering $\tilde{\mathcal{E}}$ or an integrable extension [KV, BG] of a general partial differential equation \mathcal{E} is given by introducing an additional dependent variable v and relations $D(v) = F[x, u, v]$, where F are vector differential operators on the vector-functions $u = u(x)$, $v = v(x)$, such that the expanded system is compatible by virtue of \mathcal{E} . In this case solutions of the new system $\tilde{\mathcal{E}}$ can be found from the solutions of \mathcal{E} by ODE methods.

In the case of (systems of) underdetermined ODEs with $u = (y, z, \dots)$ the additional equation has the form

$$v' = f(x, u, u', \dots, v)$$

and no integrability constraints arise.

In this section we prove that flat integrable extensions of flat model equations are in bijective correspondence with central extensions of the corresponding Carnot algebras \mathfrak{m} .

5.1. Central extensions of GNLA's. Any grading preserving automorphism of a fundamental GNLA is generated by its action on g_{-1} , and therefore can be identified with a subgroup $\mathfrak{H} \subset \mathrm{GL}(g_{-1})$. Motivated by our interest in central extensions, we call it the *gauge group* of \mathfrak{m} . The Lie algebra $\mathfrak{h} = \mathrm{Lie}(\mathfrak{H})$ equals the subspace $g_0 \subset \mathfrak{g} = \widehat{\mathfrak{m}}$ of degree 0 derivations of \mathfrak{m} .

A d -dimensional *central extension* $\tilde{\mathfrak{m}}$ of a Lie algebra \mathfrak{m} is defined by the exact sequence

$$0 \rightarrow \mathfrak{a} \hookrightarrow \tilde{\mathfrak{m}} \rightarrow \mathfrak{m} \rightarrow 0, \quad (5)$$

where \mathfrak{a} is a d -dimensional Abelian algebra belonging to the center of $\tilde{\mathfrak{m}}$. We are interested in the case, when both \mathfrak{m} and $\tilde{\mathfrak{m}}$ are fundamental GNLA, and \mathfrak{a} has a pure grading $-k$, i.e. $\mathfrak{a} \subset g_{-k}$.

For our applications the initial grading of GNLA is $(2, 1, 2, \dots)$, and then we have $3 < k \leq m + 1$, where m is the length of \mathfrak{m} . An extension of maximal grading $k = m + 1$ is

$$\tilde{\mathfrak{m}} = g_{-m-1} \oplus (g_{-m} \oplus \dots \oplus g_{-1}) = \mathfrak{a} \oplus \mathfrak{m}$$

(direct sum in the sense of vector spaces, but not Lie algebras).

It is well-known (see e.g. [SL, F]) that the 1-dimensional non-trivial central extensions of a Lie algebra \mathfrak{m} are enumerated by the elements of the Lie algebra cohomology $H^2(\mathfrak{m})$. Zero element of the cohomology group corresponds to the trivial extension $\tilde{\mathfrak{m}} = \mathfrak{m} \oplus \mathbb{R}$ and we will exclude such cases, which for exact sequence (5) means that it splits.

The same proof shows that for a GNLA \mathfrak{m} central extensions of grading $-k$ correspond to the graded part of the cohomology $H_k^2(\mathfrak{m})$. More generally, d -dimensional *nontrivial extensions* are given by d -dimensional subspaces in H_k^2 , that is, elements of the Grassmanian $\text{Gr}_d H_k^2$. Isomorphism classes of central extensions correspond to orbits of the gauge group action on $\text{Gr}_d H_k^2$, so that applying an automorphism to the cohomology class leads to an equivalent central extension.

Thus we have proved

Proposition 14. *Let \mathfrak{m} be a GNLA with fundamental space g_{-1} and the gauge group \mathfrak{H} . Its d -dimensional nontrivial central extensions of grading k are in bijective correspondence with $\text{Gr}_d H_k^2(\mathfrak{m})/\mathfrak{H}$.*

Let us notice that the space g_0 is always non-zero, since \mathfrak{H} contains the scaling corresponding to multiplication by ϵ^k on g_{-k} . However this scaling acts trivially on the Grassman space $\text{Gr}_d H_k^2(\mathfrak{m})$.

5.2. Integrable extensions of flat models. For general distributions Δ an *integrable extension* (or covering) is a submersion

$$\pi : (\tilde{M}, \tilde{\Delta}) \rightarrow (M, \Delta), \quad \text{such that}$$

- (i) $d_x \pi : \tilde{\Delta} \xrightarrow{\sim} \Delta$ is an isomorphism for any $x \in \tilde{M}$,
- (ii) for the derived distribution we have $\tilde{\Delta}_2 \cap \text{Ker } d\pi = 0$.

Then by the Frobenius theorem, any integral manifold $N \subset (M, \Delta)$ lifts to an integral manifold $\tilde{N} \subset (\tilde{M}, \tilde{\Delta})$ passing through a given point.

Indeed, the induced distribution $\tilde{\Delta} \cap T\Sigma_N$ on $\Sigma_N = \pi^{-1}N \subset \tilde{M}$ is Frobenius-integrable (and thus its integration is reduced to ODEs).

Remark 3. *To recast this into the terminology of [BG] let us choose a general connection on the bundle π , i.e. a distribution of horizontal spaces $H \subset T\tilde{M}$ such that $\tilde{\Delta}_2 \subset H$. This H defines a vertical Pfaffian system $J \subset \Omega^1(\tilde{M})$ and the integrable extension is given by $\tilde{I} = \text{Ann}(\tilde{\Delta})$, $I = \text{Ann}(\Delta)$, related by $\tilde{I} = \pi^*I + J$.*

In order to relate integrable extensions to central extensions, we restrict to integrable extensions for which tangents to the fiber V of π have pure grading $-k$ in the Carnot algebra $\tilde{\mathfrak{m}}$. The integrability condition $\tilde{\Delta}_2 \cap \text{Ker } d\pi = 0$ is then equivalent to the requirement $k > 2$.

If $\dim \Delta = 2$, then the integrability condition is vacuous, since we require the growth vector starts $(2, 1, 2, \dots)$. In this case $k > 3$.

As usual we assume that the distribution Δ is completely non-holonomic. An integrable extension π will be called *non-trivial* if $\tilde{\Delta}$ enjoys the same property, which requires additionally only that iterated brackets span TV (and surely only in this case the above grading $-k$ is defined).

Theorem 15. *1. Let $\pi : (\tilde{M}, \tilde{\Delta}) \rightarrow (M, \Delta)$ be a non-trivial integrable extension with d -dimensional fibers V . Let $\tilde{\mathfrak{m}}$ denote the Carnot algebra at $x \in \tilde{M}$ and \mathfrak{m} the Carnot algebra at $y = \pi(x) \in M$. Suppose that $\mathfrak{v} = T_x V$ correspond to pure grading $-k$ in $\tilde{\mathfrak{m}}$, $k > 2$. Then $\tilde{\mathfrak{m}}$ is a d -dimensional central extension of \mathfrak{m} (in grading $-k$).*

2. Conversely, let $\tilde{\mathfrak{m}}$ be a nontrivial central d -dimensional extension of a fundamental GNLA \mathfrak{m} in pure grading $-k$, $k > 2$, and let $(\tilde{M}, \tilde{\Delta})$ and (M, Δ) be the corresponding flat model distributions. Then $(\tilde{M}, \tilde{\Delta})$ is an integrable extension of (M, Δ) .

Remark 4. *In other words there is a bijective correspondence between classes (modulo diffeomorphisms) of integrable flat extensions of flat distributions (M, Δ) and classes (modulo automorphisms of GNLA) of central extensions of \mathfrak{m} (nontrivial and of pure grading in both cases).*

Before we start proving this theorem, let us establish the functoriality of the Tanaka map $\Delta \mapsto \mathfrak{m}$.

Proposition 16. *Any smooth map $\pi : (\tilde{M}, \tilde{\Delta}) \rightarrow (M, \Delta)$ induces a morphism of GNLA's $\tilde{\mathfrak{m}}_x \rightarrow \mathfrak{m}_{\pi(x)}$.*

Indeed, $d\pi$ maps the derived flags $\tilde{\Delta}_i \rightarrow \Delta_i$ and hence

$$d\pi : \tilde{g}_{-i} = \tilde{\Delta}_i / \tilde{\Delta}_{i-1} \rightarrow \Delta_i / \Delta_{i-1} = g_{-i}.$$

Since $\pi_* = d\pi$ respects commutators, the map is a homomorphism.

Now we can turn to the theorem about extensions.

Proof. 1. Due to Proposition 16 we have the following exact sequence of vector spaces

$$0 \rightarrow \mathfrak{v} \rightarrow \tilde{\mathfrak{m}} \xrightarrow{\pi_*} \mathfrak{m} \rightarrow 0.$$

Since π_* is an epimorphism of GNLA's it suffices to show that \mathfrak{v} belongs to the center of $\tilde{\mathfrak{m}}$.

Let $u \in \tilde{g}_{-l}$ and $v \in \mathfrak{v}$. The assumption $\mathfrak{v} \subset \tilde{g}_{-k}$ implies $[u, v] \in \tilde{g}_{-l-k}$. On the other hand, $[u, v]$ equals $[\tilde{\xi}, \eta]_x \bmod \tilde{\Delta}_{k+l-1}$, where $\tilde{\xi} \in \Gamma(\tilde{\Delta}_l)$ is a vector field with value u at x and $\eta \in \Gamma(V)$ is the vertical field with value v at x . Choosing a π -projectible vector field $\tilde{\xi}$ over some $\xi \in \Gamma(\Delta_l)$, we have $[\tilde{\xi}, \eta] \in \Gamma(V)$. This implies $[u, v] \in \mathfrak{v} \cap \tilde{g}_{-l-k} = 0$.

2. Consider a d -dimensional central extension

$$0 \rightarrow V \rightarrow \tilde{\mathfrak{m}} \xrightarrow{\varpi} \mathfrak{m} \rightarrow 0.$$

By the Tanaka theory the flat distribution Δ can be locally identified with the standard distribution on the Lie group M corresponding to \mathfrak{m} , see [T]. Let $(\tilde{M}, \tilde{\Delta})$ be the corresponding pair for $\tilde{\mathfrak{m}}$. We take both $\tilde{M} = M \times V$ and M simply-connected, and as nilpotent groups they are topological vector spaces.

The distributions Δ and $\tilde{\Delta}$ are spanned by the left-invariant vector fields from g_{-1} and \tilde{g}_{-1} respectively. Define the submersion $\pi : \tilde{M} \rightarrow M$ to be the group homomorphism, induced by the Lie algebras homomorphism ϖ . Then $d\pi$ maps $\tilde{\Delta}$ onto Δ and $\tilde{\Delta}_2$ onto Δ_2 . Since $k > 2$, the distribution $\tilde{\Delta}_2$ is horizontal and hence π is an integrable extension.

2'. Let us give an alternative more constructive proof.

Consider the integrable extension given by linearly independent cohomology classes $[\alpha_1], \dots, [\alpha_d]$, where the closed 2-forms $\alpha_i \in \Lambda^2 \mathfrak{m}^*$ have grading $k > 2$. These forms are not exact on \mathfrak{m} , but $\varpi^* \alpha_i$ are exact on the extension $\tilde{\mathfrak{m}}$, whence $\varpi^* \alpha_i = d\beta_i$ and $\beta_i|_V$ is a basis of V^* .

This construction can be made explicit as follows. Realize α_i as left-invariant 2-forms on M . Since M is contractible, $\alpha_i = d\sigma_i \in \Omega^2(M)$. We can choose σ_i and linear coordinates v_i on V in such a way that the forms $\beta_i = dv_i - \pi^* \sigma_i$ are left invariant on \tilde{M} .

Let $\tilde{\Delta} = \text{Ann}\langle \beta_1, \dots, \beta_d \rangle \cap \pi_*^{-1}(\Delta)$. We claim that $\tilde{\Delta}_2$ does not intersect the fiber V . Indeed, choose $\tilde{v}, \tilde{w} \in \Gamma(\tilde{\Delta})$. Without restriction of generality we can assume that \tilde{v}, \tilde{w} cover sections v, w of Δ . Then $\alpha_i(v, w) = 0$ thanks to the assumption $k > 2$, and so $d\beta_i(\tilde{w}, \tilde{v}) = \beta_i([\tilde{v}, \tilde{w}]) = 0$. Since $\tilde{\Delta}_2$ is generated by the commutators $[\tilde{v}, \tilde{w}]$, it is horizontal. Therefore π is an integrable extension.

Changing 2-forms within the cohomology class or applying an automorphism of \mathfrak{m} changes the distributions $\tilde{\Delta}$ to equivalent. \square

Theorem 17. *Consider a flat underdetermined ODE $\mathcal{E} : F[x, u] = 0$ ($u = u(x)$ is a vector function and F a vector differential operator) corresponding to a rank 2 distribution Δ with Carnot algebra \mathfrak{m} . Let $\tilde{\mathfrak{m}}$ be a nontrivial central extension of the GNLA \mathfrak{m} of grading $k > 3$. Then its integrable extension can be realized as an underdetermined ODE system (v has the same dimension as the extension)*

$$F[x, u] = 0, \quad v' = G[x, u]$$

In particular, if $\mathfrak{m} = \mathfrak{f}_{m,n}$ and $\mathcal{E} = \mathcal{E}_{m,n}$ is the model Monge equation, then the integrable extension can be realized as a pair of ODEs (which, in some cases, can be reduced to a higher order Monge equation).

Proof. To keep the notations simple we restrict to the case of a flat Monge equation. In Section 3.1 we constructed coordinates (x, y_i, z_j) on \mathcal{E} (which are not the canonical coordinates on $M = \mathcal{E}$ considered as the nilpotent Lie group corresponding to $\mathfrak{m} = \mathfrak{f}_{m,n}$). On the integrable extension \tilde{M} let us use the coordinates adapted for the projection π , that is the coordinates of M together with vertical coordinates v_l .

Let $\omega_1, \omega'_1, \omega_2, \dots$ be the left-invariant 1-forms dual to the basis e_1, e'_1, e_2, \dots on \mathfrak{m} chosen in Proposition 7. They satisfy the Maurer-Cartan equations $d\omega_1 = 0, d\omega'_1 = 0, d\omega_2 = \omega'_1 \wedge \omega_1, \dots$. Notice that $\omega_1 = -dx, \omega'_1 = dz_n$ and $\Delta = \text{Ann}\langle \omega_i, \omega'_j : i, j \geq 2 \rangle$.

Thus we can express the forms defining the extension (as in the previous proof) via the co-basis:

$$\beta_l = dv_l - c_{l1}\omega_1 - c_{l1'}\omega'_1 - c_{l2}\omega_2 - \dots,$$

where the coefficients $c_{l\sigma}$ are functions on M . By changing the 1-forms to cohomologous $\beta_l \mapsto \beta_l - d[\int c_{l1'} dz_n]$ we can assume that the coefficient $c_{l1'}$ is zero. Thus $\beta_l = dv_l - g_l dx \text{ mod } \langle \omega_i, \omega'_j : i, j \geq 2 \rangle$.

This implies that the integral extension $\tilde{\Delta} = \langle \omega_i, \omega'_j, \beta_l : i, j \geq 2 \rangle$ correspond to the underdetermined ODE system

$$y^{(m)} = (z^{(n)})^2, \quad v'_i = g_i(x, y, z, z', \dots).$$

For general ODE systems \mathcal{E} the proof is absolutely similar. \square

Returning to rank 2 distributions we observe that any fundamental GNLA \mathfrak{m} with gradings $(2, 1, 2, \dots)$ is a sequence of successive central extensions of the unique GNLA with gradings $(2, 1, 2)$ and this implies Theorem 3.

5.3. Examples. In this subsection we demonstrate how Proposition 14 and Theorem 17 work in practice.

5D→6D. Let us start with the unique GNLA $\mathfrak{f}_{1,2}$ with the grading $(2, 1, 2)$. Its multiplication table in the basis $e_1, e'_1, e_2, e_3, e'_3$ is (we list all non-zero commutators):

$$[e_1, e'_1] = e_2, [e_1, e_2] = e_3, [e'_1, e_2] = e'_3.$$

This is equivalent to the structure equations in the dual co-basis:

$$d\omega_1 = 0, d\omega'_1 = 0, d\omega_2 = \omega'_1 \wedge \omega_1, d\omega_3 = \omega_2 \wedge \omega_1, d\omega'_3 = \omega_2 \wedge \omega'_1. \quad (6)$$

The group $H^2(\mathfrak{m}) = H^2_4$ (of pure grading 4) is 3-dimensional, the gauge group is $\mathfrak{H} = \mathrm{GL}_2 = \mathrm{GL}(g_{-1})$ and the quotient $\mathbb{P}(H^2_4)/\mathfrak{H}$ consists of 3 points represented by the following 3 classes (called parabolic \mathfrak{p}_6 , hyperbolic \mathfrak{h}_6 and elliptic \mathfrak{ell}_6 6D algebras, because $H^2(\mathfrak{m})$ bears the natural conformal metric of Lorenzian signature $(2,1)$, see [AK]).

To calculate the integral extensions we need to realize the above abstract Maurer-Cartan forms. This is easy since they are dual to the vector fields $e_1 = -\mathcal{D}_x$, $e'_1 = \partial_{z_2}$, $e_2 = \partial_{z_1} + 2z_2\partial_y$, $e_3 = \partial_z$, $e'_3 = 2\partial_y$:

$$\begin{aligned} \omega_1 &= -dx, \omega'_1 = dz_2, \omega_2 = dz_1 - z_2dx, \\ \omega_3 &= dz - z_1dx, \omega'_3 = \frac{1}{2}(dy - z_2^2dx) - z_2(dz_1 - z_2dx). \end{aligned} \quad (7)$$

\mathfrak{p}_6 : The cohomology class of $\alpha = \omega_3 \wedge \omega_1$ corresponds to the extension

$$d\omega_4 = \omega_3 \wedge \omega_1 \quad (8)$$

or, in the dual (vector) form, to the additional commutator relation

$$[e_1, e_3] = e_4.$$

The flat model, calculated from (7) as $\omega_3 \wedge \omega_1 = -d(zdx)$, is the 1-dimensional integral extension obtained from Hilbert-Cartan equation by $v' = z$, whence the model Monge ODE is

$$\mathcal{E}_{1,3} : y' = (v''')^2.$$

$\mathfrak{h}_6 = \mathfrak{f}_{2,2}$: The cohomology class of $\alpha = \omega_3 \wedge \omega'_1 + \omega'_3 \wedge \omega_1$ corresponds to the extension $d\omega_4 = \alpha$ or, in the dual form, to

$$[e_1, e'_3] = e_4 = [e'_1, e_3].$$

The flat model, calculated from (7) as $\alpha = d\beta$ with

$$\beta = (z_1z_2 - \frac{1}{2}y)dx - z_2dx \equiv -\frac{1}{2}ydx \pmod{\langle \omega_2, \omega_3, \omega'_3 \rangle},$$

is the 1-dimensional integral extension obtained from Hilbert-Cartan equation by $v' = y$ (constant can be absorbed), whence the model Monge ODE

$$\mathcal{E}_{2,2} : v'' = (z'')^2.$$

\mathfrak{ell}_6 : The cohomology class of $\alpha = \omega_3 \wedge \omega_1 + \omega'_3 \wedge \omega'_1$ corresponds to the extension $d\omega_4 = \alpha$ or, in the dual form, to

$$[e_1, e_3] = e_4 = [e'_1, e'_3].$$

The flat model, calculated from (7) as $\alpha = d\beta$ with

$$\beta = -(z + \frac{1}{6}z_2^3)dx - \frac{1}{2}z_2dy + \frac{1}{2}z_2^2dz_1 \equiv -(z + \frac{1}{6}z_2^3)dx \pmod{\langle \omega_2, \omega_3, \omega'_3 \rangle},$$

is the 1-dimensional integral extension obtained from the Hilbert-Cartan equation by $v' = -(z + \frac{1}{6}z_2^3)$. The resulting systems is not representable as a single Monge and (after absorption of constants) equals

$$y' = (z'')^2, \quad v' = z - (z'')^3.$$

6D→**7D**. There are 8 different fundamental GNLA of dimension 7. We will consider only half of them – those which are extensions of \mathfrak{p}_6 . The structure equation of \mathfrak{p}_6 are (6)+(8).

The cohomology group consists of two pure gradings $H^2(\mathfrak{p}_6) = H_4^2 \oplus H_5^2$ and both summands are 2-dimensional. The gauge group \mathfrak{H} is the (3-dimensional) Borel subgroup of $GL(g_{-1})$. Both quotients $\mathbb{P}(H_4^2)/\mathfrak{H}$ and $\mathbb{P}(H_5^2)/\mathfrak{H}$ consist of two points and so there are 4 central extensions.

$(2, 1, 2, 1, 1)_1$. Consider at first central extension of maximal grading 5. The fixed point of the action is given by the cohomology class $[\omega_4 \wedge \omega_1]$. The corresponding central extension is the GNLA \mathfrak{p}_7 given by the additional commutation relation

$$[e_1, e_4] = e_5$$

and the integral extension is

$$y' = (z^{iv})^2.$$

$(2, 1, 2, 1, 1)_2$. Another central 1-dimensional extension is obtained with the cohomology class $[\omega_4 \wedge \omega'_1 - \omega_3 \wedge \omega_2]$ (\mathfrak{H} -orbit through it is open). The corresponding central extension is given by

$$[e'_1, e_4] = e_5 = [e_3, e_2]$$

and the integral extension is given by the system

$$y' = (z''')^2, \quad v' = (z'')^2.$$

$(2, 1, 2, 2)_1$. Now consider the central extensions of grading 4. The fixed point of the \mathfrak{H} -action is given here by the class $[\omega_3 \wedge \omega'_1 + \omega'_3 \wedge \omega_1]$. The corresponding central extension is the GNLA $\mathfrak{f}_{2,3}$ given by

$$[e_1, e'_3] = e'_4 = [e'_1, e_3]$$

and the integral extension is given by the Monge equation

$$y'' = (z''')^2.$$

$(2, 1, 2, 2)_2$. Finally, another central extensions of grading 4 corresponds to an open orbit of the \mathfrak{H} -action and is given by the class $[\omega'_3 \wedge \omega'_1]$. The corresponding central extension is given by

$$[e'_1, e'_3] = e'_4$$

and the integral extension is given by the system

$$y' = (z''')^2, \quad v' = (z''')^3.$$

5.4. Extensions of parabolic models \mathfrak{p}_n . The GNLA's $\mathfrak{p}_n = \mathfrak{f}_{1,n-3}$ are of special interest as the most symmetric cases. It is obvious that \mathfrak{p}_{n+1} is a central 1-dimensional extension of \mathfrak{p}_n . We would like to study other possible central extensions of \mathfrak{p}_n of pure grading.

Notice that any fundamental GNLA with $\dim g_{-1} = 2$ can be obtained from $(2, 1, 2)$ by successive d -dimensional central extensions of only maximal grading. In studying the latter for the sequence of parabolic models we observe the following \mathbb{Z}_2 -periodic pattern.

Theorem 18. *The Lie algebra \mathfrak{p}_{n+1} is a maximal grading central extension of \mathfrak{p}_n . For odd n such an extension is unique, while for even $n = 2l$ there are two maximal grading extensions \mathfrak{p}_{n+1} and \mathfrak{p}'_{n+1} .*

The second is however less symmetric (meaning $\dim \widehat{\mathfrak{p}'_{n+1}} < \dim \widehat{\mathfrak{p}_{n+1}}$) and has no maximal grading central extensions.

Proof. The Maurer-Cartan equations of \mathfrak{p}_n are (the index denotes the grading):

$$\begin{aligned} d\omega_1 &= 0, \quad d\omega'_1 = 0, \quad d\omega_2 = \omega'_1 \wedge \omega_1, \quad d\omega_3 = \omega_2 \wedge \omega_1, \quad d\omega'_3 = \omega_2 \wedge \omega'_1, \\ d\omega_4 &= \omega_3 \wedge \omega_1, \quad d\omega_5 = \omega_4 \wedge \omega_1, \quad \dots \quad d\omega_{n-2} = \omega_{n-3} \wedge \omega_1. \end{aligned}$$

The 2-cocycles in the maximal grading are

$$Z_{n-1}^2 = \langle \omega_{n-2} \wedge \omega_1, \omega_{n-2} \wedge \omega'_1 - \omega_{n-3} \wedge \omega_2 + \omega_{n-4} \wedge \omega_3 - \dots \pm \omega_l \wedge \omega_{l-1} \rangle$$

for even $n = 2l$ and $Z_{n-1}^2 = \langle \omega_{n-2} \wedge \omega_1 \rangle$ for odd $n = 2l + 1$

There are no coboundaries in this grading and so $\mathbb{P}H_{n-1}^2 = \mathbb{R}\mathbb{P}^\epsilon$, where $\epsilon = 0$ for odd n and $\epsilon = 1$ for even n . The action of 3-dimensional gauge group \mathfrak{H} has two orbits in the latter case: fixed point $[\omega_{n-2} \wedge \omega_1]$ and the complement, so that we can take the the second form of Z_{n-1}^2 for $n = 2l$ as a representative.

Extension through the first class is possible for any n and the result is \mathfrak{p}_{n+1} . Extension via the second class yields the GNLA \mathfrak{p}'_{2l+1} for $n = 2l$ with the following additional commutation relation:

$$[e'_1, e_{n-2}] = e_{n-1}, [e_2, e_{n-3}] = -e_{n-1}, [e_3, e_{n-4}] = e_{n-1} \dots$$

It's easy to check that there are no 2-cocycles $Z_{2l}^2(\mathfrak{p}'_{2l+1}) = 0$, and so $H_{2l}^2(\mathfrak{p}'_{2l+1}) = 0$. Thus no further maximal central extensions exist.

Finally to calculate the Tanaka prolongation of \mathfrak{p}'_{2l+1} notice that both directions $\mathbb{R}e_1$ and $\mathbb{R}e'_1$ are canonical (invariant under automorphisms) in g_{-1} because they can be uniquely characterized by the properties $\text{ad}_{e_1}|_{g_{2-2l}} = 0$, $\text{ad}_{e'_1}|_{g_{-3}} = 0$. Therefore the space g_0 is 2-dimensional Abelian (diagonal 2×2 matrices). Calculations similar to that of the proof of Proposition 9 show that $g_{-1} = 0$ and so $\dim \widehat{\mathfrak{p}'_{2l+1}} = 2l + 3$ is smaller than $\dim \widehat{\mathfrak{p}_{2l+1}} = 4l - 1$. \square

The flat ODE for \mathfrak{p}_n is given by the model equation $\mathcal{E}_{1,n-3}$. The flat ODE for \mathfrak{p}'_{2l+1} is not realized as a Monge equation, but only as a system. To find it one follows the scheme of Section 5.3 and therefore obtains the following (non-standard) integral extension of $\mathcal{E}_{1,2l-3}$:

$$y' = (z^{(2l-3)})^2, v' = (z^{(l-1)})^2.$$

6. APPLICATIONS TO INTEGRATION OF PDES

We conclude with some examples which illustrate how our study of rank 2 distributions for Monge equations can be applied to the integration of overdetermined and determined PDEs.

6.1. From underdetermined ODEs to overdetermined PDEs.

The examples here generalize those of Cartan [C1]. Specifically, Cartan considered compatible systems consisting of pairs of scalar 2nd order PDEs on the plane with one common characteristic (which we denote as $2E_2$). The equation $\mathcal{E} \subset J^2(\mathbb{R}^2, \mathbb{R})$ from the internal viewpoint is a 6-dimensional manifold N_6 equipped with a rank 3 distribution. This distribution always has a Cauchy characteristic and can therefore be reduced to a rank 2 distribution on a 5-dimensional manifold N_5 , which is necessarily equivalent to a single Monge equation (3).

Now we consider the case of mixed order equations \mathcal{E} consisting of pairs of compatible 2nd and 3rd order PDEs ($E_2 + E_3$). Again it can be seen that such systems can be reduced to rank 2 distributions, now on 6-dimensional manifolds, but we would like to emphasize that the reduction is slightly more complicated than in Cartan's case.

Indeed the system \mathcal{E} is represented internally as an 8-dimensional manifold M_8 equipped with a rank 3 distribution \mathcal{Q} . This distribution has one Cauchy characteristic, while its derived distribution \mathcal{Q}_2 has two. Thus the pair (M_8, \mathcal{Q}) has two reductions - one by Cauchy characteristic for \mathcal{Q} (quotient) and one by Cauchy characteristic for the derived distribution \mathcal{Q}_2 (de-prolongation).

These two reductions commute, so that we can do them in any order, and in Examples 1 and 2 we perform both at once. But for the purposes of Example 3 we shall need first to de-prolong M_8 to M_7 (with distribution $\bar{\mathcal{Q}}$ of rank 3), and then to quotient $(M_7, \bar{\mathcal{Q}})$ to (M_6, Δ) (rank of Δ is 2). These steps are summarized in the following sequence of maps:

$$(M_8, \mathcal{Q}) \longrightarrow (M_7, \bar{\mathcal{Q}}) \longrightarrow (M_6, \Delta).$$

In the first two examples (M_6, Δ) turns out to be one of the 6-dimensional symmetric models from Theorem 1 (this can be detected a priori via calculation of the symmetry algebra and concluding that the distribution is flat in the Tanaka sense). Then we use their integral curves to integrate the corresponding PDE systems.

Example 1: $\mathbf{E}_2 + \mathbf{E}_3 \mapsto \mathcal{E}_{1,3}$. Consider the PDE system, written parametrically as

$$u_{xx} = \tau^5 x^2, \quad u_{xy} = \tau^4 x^2, \quad u_{xyy} = \frac{5}{4}\tau u_{yyy} + \frac{4}{5}\tau^3 x. \quad (9)$$

It is compatible and has a common characteristic $\xi = \mathcal{D}_x - \frac{5}{4}\tau \mathcal{D}_y$.

Internally the equation is $M_8 = \mathbb{R}^8(x, y, u, u_x, u_y, \tau, u_{yy}, u_{yyy})$ and the derived distribution \mathcal{Q}_2 has a two-dimensional space of Cauchy characteristics

$$\begin{aligned} \Pi = \langle \hat{\xi} = & \partial_x - \frac{5}{4}\tau \partial_y + (u_x - \frac{5}{4}\tau u_y) \partial_u - \frac{1}{4}\tau^5 x^2 \partial_{u_x} \\ & + \frac{\tau}{4}(4\tau^3 x^2 - 5u_{yy}) \partial_{u_y} + \frac{4}{5}\tau^3 x \partial_{u_{yy}} - \frac{\tau}{2x} \partial_\tau, \partial_{u_{yyy}} \rangle. \end{aligned}$$

The distribution Π is integrable and the quotient by its leaves

$$M_6 = M_8/\Pi \simeq \mathbb{R}^6(t, v, w, w_1, w_2, w_3)$$

can be identified with the space of invariants

$$\begin{aligned} t &= y + \frac{5}{2}\tau x, \quad v = \frac{256}{625}u_x + \frac{128}{625}\tau^5 x^3, \\ w &= -\frac{7}{2}\tau^5 x^4 + \frac{25}{8}\tau^2 x^2 u_{yy} + \frac{5}{2}\tau x u_y - x u_x + u, \\ w_1 &= u_y + \frac{5}{2}\tau x u_{yy} - 3\tau^4 x^3, \quad w_2 = u_{yy} - \frac{8}{5}\tau^3 x^2, \quad w_3 = -\frac{16}{25}\tau^2 x. \end{aligned} \quad (10)$$

The quotient distribution on the manifold M_6 equals

$$\Delta = \langle \partial_t + w_3^2 \partial_v + w_1 \partial_w + w_2 \partial_{w_1} + w_3 \partial_{w_2}, \partial_{w_3} \rangle$$

and this corresponds to the model Monge equation $\mathcal{E}_{1,3}$:

$$v' = (w''')^2.$$

The inverse image of the general solution to this equation with respect to the projection map $\pi : M_8 \rightarrow M_6$ given by (10) (this introduces only 1 more parameter: π^{-1} decomposes into preimage and prolongation) yields the general solution of (9), written parametrically as

$$\begin{aligned} x &= -\left(\frac{8s}{25}\right)^2 f'''(t), \quad y = t - s f'''(t), \\ u &= f(t) - \frac{s^3}{20} f'''(t)^4 + \frac{s^2}{2} f''(t) f'''(t)^2 - s f'(t) f'''(t) - \frac{s^2}{4} f'''(t) g(t), \end{aligned}$$

where $g'(t) = f'''(t)^2$.

Example 2: $\mathbf{E}_2 + \mathbf{E}_3 \mapsto \mathcal{E}_{2,2}$. Now let's consider the system

$$2u_{xx} = y^2 - \tau^2, \quad 2u_{yy} = \tau^{-2}, \quad u_{xyy} - \tau^2 u_{yyy} = \tau^{-1}, \quad (11)$$

which is again compatible with common characteristic $\xi = \mathcal{D}_x - \tau^2 \mathcal{D}_y$.

Internally, the equation is $M_8 = \mathbb{R}^8(x, y, u, u_x, u_y, \tau, u_{xy}, u_{yyy})$ and the derived distribution \mathcal{Q}_2 has a two-dimensional space of Cauchy characteristics

$$\begin{aligned} \Pi = \langle \hat{\xi} &= \partial_x - \tau^2 \partial_y + (u_x - u_y \tau^2) \partial_u + \left(\frac{y^2 - \tau^2}{2} - u_{xy} \tau^2\right) \partial_{u_x} \\ &+ (u_{xy} - \frac{1}{2}) \partial_{u_y} + (y - \tau) \partial_{u_{xy}} - \tau^2 \partial_\tau, \partial_{u_{yyy}} \rangle. \end{aligned}$$

The distribution Π is integrable and the quotient by its leaves

$$M_6 = M_8 / \Pi \simeq \mathbb{R}^6(t, v, v_1, w, w_1, w_2)$$

is identified with the space of invariants

$$\begin{aligned} t &= y - \tau, \quad v = \ln \tau - 1 + 2u - 2\tau u_y + 2u_{xy} - 2xy - \tau x^2 y + 2\tau x + \\ &+ \frac{x^2(y^2 + \tau^2)}{2} - 2x u_x + 2\tau x u_{xy}, \quad v_1 = x^2 y + \frac{1}{\tau} - \tau x^2 - 2x u_{xy} + 2u_y, \\ w &= \frac{xy^2}{2} - y - \tau xy + \tau + \frac{\tau^2 x}{2} + \tau u_{xy} - u_x, \\ w_1 &= xy - \tau x - u_{xy}, \quad w_2 = -\frac{1}{\tau} + x. \end{aligned}$$

In these coordinates the quotient distribution Δ on M_6 coincides with the rank 2 distribution of the model Monge equation $\mathcal{E}_{2,2}$:

$$v'' = (w'')^2.$$

Again the general solution of this equation provides the general solution to system (11) as

$$x = f''(t) + \frac{1}{s}, \quad y = t + s, \quad u = \frac{(2s+t)t}{4}f''(t)^2 + \frac{s}{2}g'(t) + \frac{g(t)}{2} - \frac{2f'(t)s^2 + 2f(t)s - t^2}{2s}f''(t) - \frac{f(t)}{s} - \frac{t}{2s} + \frac{t^2}{4s^2} - \frac{1}{2}\ln(s),$$

where $g''(t) = f''(t)^2$.

We remark that the 2nd order PDE in (9) $x^2u_{xx}^4 = u_{xy}^5$ is a new example of a Darboux integrable equation. On the other hand, the 2nd order PDE in (11) $2u_{xx} + (2u_{yy})^{-1} = y^2$ is not Darboux integrable and can therefore only be solved by compatibility technique.

Example 3: $\mathbf{E}_2 + \mathbf{E}_3 \rightsquigarrow 2\mathbf{E}_2$. In this example we show that our theory of integrable extensions for Monge equations can be applied to construct integrable extensions of overdetermined PDE systems. Informally speaking, if \mathcal{E} and \mathcal{E}' are two PDE systems whose reduction by Cauchy characteristics and de-prolongations are rank 2 distributions corresponding to equations $\bar{\mathcal{E}}$ and $\bar{\mathcal{E}}'$, then an integrable extension $\varphi : \bar{\mathcal{E}} \rightarrow \bar{\mathcal{E}}'$ can be lifted to an integrable extension $\Phi : \mathcal{E} \rightarrow \mathcal{E}'$.

To be more precise, in the notations adopted at the beginning of this section we have the following commutative diagram (we mark by primes the canonical distributions of the manifolds N_i)

$$\begin{array}{ccc} (M_7, \bar{\mathcal{Q}}) & \xrightarrow{\Phi} & (N_6, \mathcal{Q}') \\ \pi \downarrow & & \downarrow \pi' \\ (M_6, \Delta) & \xrightarrow{\varphi} & (N_5, \Delta') \end{array}$$

where the horizontal arrows are the integrable extensions and the vertical arrows are the quotients of the rank 3 distributions $\bar{\mathcal{Q}}$ and \mathcal{Q}' by their Cauchy characteristics.

We illustrate this construction with the example of Cartan's involutive symmetric model [C1]

$$U_{XX} = \frac{1}{3}\tau^3, \quad U_{XY} = \frac{1}{2}\tau^2, \quad U_{YY} = \tau. \quad (12)$$

This system reduces to $\mathcal{E}_{1,2}$: $V' = (W'')^2$ with $V = V(T)$, $W = W(T)$ and the projection π' along the Cauchy characteristic defined by

$$\begin{aligned} T &= -U_{YY}, \quad V = -2U + 2XU_X + 2YU_Y - Y^2U_{YY} - XYU_{YY}^2 - \frac{1}{3}X^2U_{YY}^3, \\ W &= U_X - U_YU_{YY} + \frac{1}{2}YU_{YY}^2 + \frac{1}{6}XU_{YY}^3, \\ W_1 &= U_Y - YU_{YY} - \frac{1}{2}XU_{YY}^2, \quad W_2 = Y + XU_{YY} \end{aligned}$$

Basing on the fact that $\mathcal{E}_{1,3}$ is an integrable extension of $\mathcal{E}_{1,2}$ and on the reduction from Example 1 of this section, we will show that (9) is an integrable extension of (12).

Recall that $(M_7, \bar{\mathcal{Q}})$ is obtained from (M_8, \mathcal{Q}) by de-prolongation via a Cauchy characteristic for \mathcal{Q}_2 not belonging to \mathcal{Q} . Thus $\bar{\mathcal{Q}}$ still carries a Cauchy characteristic and we get $(M_7, \bar{\mathcal{Q}}) \simeq (\mathcal{E}_{1,3}, \mathcal{C}^{1,3}) \times (\mathbb{R}, \mathbb{R})$ and similarly $(N_6, \mathcal{Q}') \simeq (\mathcal{E}_{1,2}, \mathcal{C}^{1,2}) \times (\mathbb{R}, \mathbb{R})$ for the Cartan example, because $(M_6, \Delta) \simeq (\mathcal{E}_{1,3}, \mathcal{C}^{1,3})$ and $(N_5, \Delta') \simeq (\mathcal{E}_{1,2}, \mathcal{C}^{1,2})$.

The central extension $\varpi : \mathfrak{p}_6 \rightarrow \mathfrak{p}_5$ induces the integrable extension $\phi = \varphi \times 1 : (\mathcal{E}_{1,3}, \mathcal{C}) \times (\mathbb{R}, \mathbb{R}) \rightarrow (\mathcal{E}_{1,2}, \mathcal{C}') \times (\mathbb{R}, \mathbb{R})$, where

$$\varphi(t, v, w, w_1, w_2, w_3) = (t, v, w_1, w_2, w_3) = (T, V, W, W_1, W_2).$$

By this ϕ we construct the required map Φ in the diagram, and then we compute its inverse Φ^{-1} given by the formulae

$$\begin{aligned} x &= -\left(\frac{4}{5}X\right)^2(Y + XU_{YY})^{-1}, & y &= -U_{YY} - \frac{5}{2}X, \\ u_x &= \left(\frac{5}{4}\right)^4\left(-\frac{1}{3}X^2U_{YY}^3 - \frac{1}{2}X(2Y + X^2)U_{YY}^2 - (X^2 + Y)YU_{YY}\right. \\ &\quad \left.+ 2XU_X + 2YU_Y - 2U - \frac{1}{2}XY^2\right), \\ u_y &= \frac{1}{6}XU_{YY}^3 + \left(\frac{5}{4}X^2 + \frac{1}{2}Y\right)U_{YY}^2 + \left(\frac{25}{16}X^3 - U_Y + \frac{5}{2}XY\right)U_{YY} \\ &\quad + U_X - \frac{5}{2}XU_Y + \frac{25}{16}X^2Y, \end{aligned}$$

where $(X, Y, U, U_X, U_Y, \dots)$ are jet-coordinates on the equation N_6 (12) and $(x, y, u, u_x, u_y, \dots)$ are jet-coordinates on the equation M_7 . Prolongation of the latter yields the required integrable extension M_8 .

Notice that U is not expressed in the above formulae but is obtained via quadrature, as is always the case with flat integrable extensions. We leave it to the reader to check that the integrability condition $D_x(u_y) = D_y(u_x)$ holds by virtue of (12) and satisfies (9).

6.2. Relation to the method of Darboux. According to [AFV] any Darboux integrable single scalar PDE on the plane $\mathcal{E} \subset J^2(\mathbb{R}^2, \mathbb{R})$ is a group quotient of a pair of rank 2 distributions with common freely acting symmetry group G , that is

$$(\mathcal{E}, \mathcal{C}) = (M_1, \Delta_1) \times (M_2, \Delta_2)/G.$$

Notice that in this case $\dim \mathcal{E} = 7$, $\dim \mathcal{C} = 4$, but $\dim M_i$ can be sufficiently large, depending on the order of intermediate integrals of \mathcal{E} .

Our theory of rank 2 distributions of maximal symmetry restricts the possible types of the factors.

Theorem 19. *Suppose that (M_i, Δ_i) have Carnot algebras \mathfrak{f}_{m_i, n_i} . Then the pairs (m_1, n_1) , (m_2, n_2) with normalization $0 < m_i \leq n_i$, $n_2 - n_1 =$*

$k \geq 0$, and $m_1 \leq m_2$ for $k = 0$ satisfy the condition

$$m_1 + m_2 + k \leq 7 + \delta,$$

where $\delta = 1$ if $m_1 = m_2 = 1$ or $m_1 = 1, k > 0$, and $\delta = 0$ else. Thus there are no more than 53 different admissible triples (m_1, m_2, k) .

Proof. The equality

$$\dim M_1 + \dim M_2 - \dim G = \dim \mathcal{E} = 7$$

implies $\dim G = n_1 + m_1 + n_2 + m_2 - 3$. Since G acts as the symmetry group for both (M_i, Δ_i) and since the symmetry group of a distribution is bounded from above by Tanaka algebra $\widehat{\mathfrak{f}_{m,n}}$, i.e. $\dim G \leq \dim \mathfrak{t}_{m_i, n_i}$ we get that $\min\{\dim \mathfrak{t}_{m_i, n_i} : i = 1, 2\} \geq n_1 + m_1 + n_2 + m_2 - 3$. From Section 3.4 we know that $\dim \mathfrak{t}_{m_i, n_i} = 2n_i + 5$ for $m_i = 1$ and $2n_i + 4$ for $m_i > 1$. The claim follows. \square

Recall that distributions Δ_i corresponding to nondegenerate Monge equations (3) have Carnot algebras \mathfrak{f}_{m_i, n_i} . The estimate of the theorem is rough and can be sharpened by a detailed analysis of the admissible (M_i, Δ_i) which actually arise for various types of equations \mathcal{E} .

Example ([AF]). The Goursat equation $u_{xy} = a(x, y)\sqrt{u_x u_y}$ is Darboux integrable if and only if both factors (M_i, Δ_i) can be realized as rank 2 distributions of the Monge equation

$$y' = F(x, z, z', \dots, z^{(n)})$$

and the group G is either Abelian or 1-step solvable of dimension $2n+1$.

In particular, for the Darboux integrable equations

$$u_{xy} = \frac{2n}{x+y} \sqrt{u_x u_y}$$

both factors coincide with (the prolongation of) the model equations $\mathcal{E}_{1,n}$. In this case, the Lie algebra of G is equal to the nilradical \mathfrak{r} of the Lie algebra $\mathfrak{t}_{1,n}$ given in Section 3.3.

REFERENCES

- [AF] I. M. Anderson, M. Fels, *Transformations of Darboux integrable systems*, in: Differential equations: Geometry, Symmetry and Integrability. The Abel Symposium 2008, Ed: B. Kruglikov, V. Lychagin, E. Straume, 21–48 (2009).
- [AFV] I. M. Anderson, M. Fels, P. Vassiliou, *Superposition formulas for exterior differential systems*, Adv. Math. **221**, no. 6, 1910–1963 (2009).
- [AKO] I. M. Anderson, N. Kamran, P. Olver, *Internal, external and generalized symmetries*, Adv. Math. **100**, 53–100 (1993).
- [AK] I. M. Anderson, B. Kruglikov: in preparation.

- [BG] R. Bryant, Ph. Griffiths, *Characteristic cohomology of differential systems. II. Conservation laws for a class of parabolic equations*, Duke Math. J. **78**, no. 3, 531–676 (1995).
- [C1] E. Cartan, *Les systèmes de Pfaff, à cinq variables et les équations aux dérivées partielles du second ordre*, Ann. Sci. École Norm. Sup. (3) **27**, 109–192 (1910).
- [C2] E. Cartan, *Sur l'équivalence absolue de certains systèmes d'équations différentielles et sur certaines familles de courbes*, Bulletin de la S.M.F. **42**, 12–48 (1914).
- [DZ] B. Doubrov, I. Zelenko, *A canonical frame for nonholonomic rank two distributions of maximal class*, C.R. Math. Acad. Sci. Paris **342**, no. 8, 589–594 (2006).
- [F] D. B. Fuks, *Cohomology of infinite-dimensional Lie algebras*, Contemporary Soviet Mathematics. Consultants Bureau, New York (1986).
- [G] E. Goursat, *Leçons sur le problème de Pfaff*, Hermann, Paris (1923).
- [GK] R. Gardner, N. Kamran, *Characteristics and the geometry of hyperbolic equations in the plane*, J. Diff. Equations **104**, 60–116 (1993).
- [GKR] A. Giaro, A. Kumpera, C. Ruiz, *Sur la lecture correcte d'un résultat d'Élie Cartan*, C.R.Acad.Sci. Paris Sér. A-B **287**, no. 4, A241–A244 (1978).
- [H] D. Hilbert, *Über den begriff der klasse von differentialgleichungen*, Math. Ann. **73**, no. 1, 95–108 (1912).
- [KL] B. Kruglikov, V. V. Lychagin, *Geometry of Differential equations*, In: Handbook on Global Analysis, D.Krupka and D.Saunders Eds., 725–771, Elsevier Sci. (2008).
- [KLV] I. S. Krasilschik, V. V. Lychagin, A. M. Vinogradov, *Geometry of jet spaces and differential equations*, Gordon and Breach (1986).
- [KV] I. S. Krasilschik, A. M. Vinogradov, *Nonlocal trends in the geometry of differential equations: symmetries, conservation laws, and Bäcklund transformations. Symmetries of partial differential equations, Part I*, Acta Appl. Math. **15**, no. 1-2, 161–209 (1989).
- [L] S. Lie, *Zur allgemeinen theorie der partiellen differentialgleichungen beliebiger ordnung*, Gesammelte Abhandlungen, B.G. Teubner (Leipzig) - H.Aschehoung (Oslo), Bd.4 (1929), paper IX.
- [MT] T. Morimoto, N. Tanaka, *The classification of the real primitive infinite Lie algebras*, J. Math. Kyoto Univ. **10**, no.2, 207–243 (1970).
- [MZ] R. Montgomery, M. Zhitomirskii, *Geometric approach to Goursat flags*, Ann. Inst. H. Poincaré, Anal. Non Linéaire **18**, no. 4, 459–493 (2001).
- [SL] Séminaire “Sophus Lie” de l'École Normale Supérieure, 1954/1955. *Théorie des algèbres de Lie. Topologie des groupes de Lie*, Secrétariat mathématique, 11 rue Pierre Curie, Paris (1955).
- [T] N. Tanaka, *On differential systems, graded Lie algebras and pseudo-groups*, J. Math. Kyoto Univ. **10**, no.1, 1–82 (1970).
- [Y] K. Yamaguchi, *Differential systems associated with simple graded Lie algebras*, Adv. Studies in Pure Math. **22**, 413–494 (1993).
- [W] E. von Weber, *Zur Invariantentheorie der Systeme Pfaff'scher Gleichungen*, Berichte Ges. Leipzig, Math-Phys. Classe **L**, 207–229 (1898).