

1-1-2009

Geochemistry and potential playa sources of the January 7, 2008 Southwestern New Mexico "milky rain"

Joel Gilbert

Department of Geological Sciences, University of Texas, El Paso

Thomas E. Gill

Department of Geological Sciences, University of Texas, El Paso

David Borrok

Department of Geological Sciences, University of Texas, El Paso

Bonnie Frey

New Mexico Bureau of Geology and Mineral Resources, Socorro

Terry Hertel

New Mexico Environment Department, Air Quality Bureau, Santa Fe

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.usu.edu/nrei>

Recommended Citation

Gilbert, Joel; Gill, Thomas E.; Borrok, David; Frey, Bonnie; Hertel, Terry; Lane, Maria; Bleiweiss, Max; Lehmann, Chris; Gay, David; and Ramana, Chintalapalle V. (2009) "Geochemistry and potential playa sources of the January 7, 2008 Southwestern New Mexico "milky rain"," *Natural Resources and Environmental Issues*: Vol. 15, Article 21.

Available at: <http://digitalcommons.usu.edu/nrei/vol15/iss1/21>

This Article is brought to you for free and open access by the Quinney Natural Resources Research Library, S.J. and Jessie E. at DigitalCommons@USU. It has been accepted for inclusion in Natural Resources and Environmental Issues by an authorized administrator of DigitalCommons@USU. For more information, please contact becky.thoms@usu.edu.



Authors

Joel Gilbert, Thomas E. Gill, David Borrok, Bonnie Frey, Terry Hertel, Maria Lane, Max Bleiweiss, Chris Lehmann, David Gay, and Chintalapalle V. Ramana

Geochemistry and Potential Playa Sources of the January 7, 2008 Southwestern New Mexico "Milky Rain"

Joel Gilbert^{1,2}, Thomas E. Gill^{2,3}, David Borrok², Bonnie Frey⁴, Terry Hertel⁵, Maria Lane⁶, Max Bleiweiss⁷, Chris Lehmann⁸, David Gay⁸ & Chintalapalle V. Ramana⁹

¹Environmental Science Program, University of Texas-El Paso, TX 79968; ²Department of Geological Sciences, University of Texas-El Paso, TX 79968; ³Environmental Science and Engineering Program, University of Texas-El Paso, TX 79968; ⁴New Mexico Bureau of Geology & Mineral Resources, New Mexico Institute of Mining and Technology, 801 Leroy Place, Socorro, NM 87801; ⁵New Mexico Environment Department, New Mexico Air Quality Bureau, 1301 Siler Road, Building B, Santa Fe NM 87507; ⁶Department of Geography, University of New Mexico, 1 University Lane, Albuquerque, NM 87131; ⁷Department of Entomology, Plant Pathology and Weed Science, New Mexico State University, Las Cruces, NM 88003; ⁸National Atmospheric Deposition Program, Illinois State Water Survey, University of Illinois at Urbana-Champaign, 2204 Griffith Drive, Champaign, IL 61820; ⁹Metallurgical and Materials Engineering, University of Texas - El Paso, 500 West University Avenue, El Paso, TX 79968

Corresponding author:

Joel Gilbert

Environmental Science Program, University of Texas-El Paso, 500 West University Avenue, El Paso TX 79968, USA

E-mail: jhgilbert@utep.edu

On the afternoon of January 7, 2008, an approximately 1300 km² region of Grant County, New Mexico, USA (Figure 1A), was subject to an unusual rainfall of milky-white color that left a hard, solid residue on surfaces (Figure 1D). Concerned residents, who collected samples from cisterns, rain gauges, and roof runoff, contacted area universities and state agencies to arrange for chemical analyses. Initial speculation of the source targeted everything from local mine tailings to Pacific Rim volcanic ash. Scientific evaluation of wind back trajectories, MODIS satellite images, and photographic observations indicated a possible regional source mechanism: wind erosion and transport of whitish dust from Willcox Playa in southeastern Arizona, USA (Figure 1B), followed by downwind entrainment into clouds, nucleation and wet deposition in localized rain showers downwind. Other rare "milky rain" events worldwide were traced to rainout of dust from saline lake basins (Resane, T.H. et al. 2004, South African Journal of Science 100: 483-487), and plumes of Willcox Playa dust have been previously observed to have been advected above the region as aerosols. Milky rain pH was 7.20 compared with the

regional average of 5.66. Electrical conductivity ranged from 46.5 to 115 $\mu\text{S}/\text{cm}$ as opposed to an average of 4.3 $\mu\text{S}/\text{cm}$ for the region. Analysis of major ions indicated elevated levels of calcium, magnesium, sodium, potassium, sulfate and chloride, as would be expected in playa-type deposits. A National Atmospheric Deposition Program site at the Gila Cliff Dwellings determined the calcium concentration of the rain (5.8 mg/l) to be within the top 1.0% of all data gathered by the network in a typical 5 year period. Upon analysis, the New Mexico Environment Department concluded that the event samples had similar chemistry to surface water samples obtained from the Lordsburg Playas in the early 1990s and reassured the public regarding its possible (non-) toxicity; the Lordsburg Playas (a known dust source along the same trajectory between the affected region and Willcox Playa) may have also contributed to this event. SEM analysis of water filter residue (insolubles and precipitates) from the event showed particles $\leq 1 \mu\text{m}$ in diameter (Figure 1C) with an elemental composition (Si, Al, Ca, and Mg) consistent with aluminosilicate and evaporite minerals.




Figure 1—A: MODIS satellite image of the event showing plumes of dust rising off the Willcox Playa (1) and location of Lordsburg Playas (circle) and area affected by milky rain (square) (2). B: Dust plumes moving right to left (south to north) across Interstate Highway 10. The image was coincidentally taken on day of event. Willcox Playa is just off image to right. C: Scanning electron micrograph of filter residue. D: Image of dried residue left on car hood (Photo courtesy of Alyson Siwik).