

2002

# How Can I Make My House Safe For My Pets?

Clell Bagley

Follow this and additional works at: [http://digitalcommons.usu.edu/extension\\_histall](http://digitalcommons.usu.edu/extension_histall)

 Part of the [Education Commons](#)

**Warning:** The information in this series may be obsolete. It is presented here for historical purposes only. For the most up to date information please visit [The Utah State University Cooperative Extension Office](#)

---

## Recommended Citation

Bagley, Clell, "How Can I Make My House Safe For My Pets?" (2002). *All Archived Publications*. Paper 415.  
[http://digitalcommons.usu.edu/extension\\_histall/415](http://digitalcommons.usu.edu/extension_histall/415)

This Article is brought to you for free and open access by the Archived USU Extension Publications at DigitalCommons@USU. It has been accepted for inclusion in All Archived Publications by an authorized administrator of DigitalCommons@USU. For more information, please contact [dylan.burns@usu.edu](mailto:dylan.burns@usu.edu).





**A weekly question/answer column**

**How Can I Make My House Safe For My  
Pets?**

*Clell Bagley\* answers:*

Just as candy, pills and house plants attract children, they are just as great a temptation for pets. They can also be toxic to them.

- Chocolate is toxic to both cats and dogs, but dogs are much more likely to eat a toxic dose. As little as 2 1/2 ounces of baking chocolate could poison a 20 pound dog. Other forms of chocolate are less toxic, but they still can be potentially dangerous. The toxins, caffeine and theobromine, are rapidly absorbed but slowly excreted by cats and dogs.
- Ornamental plants such as mistletoe, holly and poinsettia are occasionally eaten by dogs and cats. Just one to three berries, especially mistletoe, may result in clinical signs and even death of pets.
- Keep pain medications put away where dogs don't have access to them. They seem to be attracted to ibuprofen, acetaminophen, aspirin and indomethacin. In one summary of poisoning cases, accidental exposures accounted for 75 percent of all the poisoning cases in dogs.
- Keep acetaminophen away from cats, as they lack the enzyme needed to detoxify one of its ingredients. Aspirin is also dangerous for cats since they can easily be overdosed.

---

\* Clell Bagley is Utah State University Extension Veterinarian