



## Blueberries in Utah? Difficult, but Maybe Not Impossible

*Dr. Brent Black*, USU Extension Fruit Specialist, *Dr. Grant Cardon*, USU Extension Soils Specialist  
*Dr. Mark Ehlenfeldt*, USDA-ARS Research Geneticist

U.S. consumers have become increasingly interested in blueberries. Part of this is due to the exceptional flavor and versatility of the berries, while some of the interest can be attributed to recent discoveries regarding the health benefits of this “super food.” In response to this increased demand, more chain stores are now selling blueberry plants to intrepid and/or gullible Utah gardeners. Here we describe the difficulties you will encounter in home blueberry production.

### Background

Commercially harvested blueberry fruit come from one of several different blueberry species, including lowbush, highbush, and rabbiteye. The lowbush or wild blueberry is native to the northeastern U.S. and eastern Canadian provinces. The highbush blueberry originated in the coastal forests of eastern North America and was first domesticated in the Pine Barrens of New Jersey. The rabbiteye blueberry is native to coastal regions of the southeastern U.S. including Florida, Georgia and the Carolinas. All of these forested coastal regions have sandy soils with high organic matter content, a low pH, and high annual rainfall. As one might expect, blueberries are uniquely adapted to this environment. Specifically, the root system is broad, shallow and fibrous, and requires soil that has an acidic pH, that is coarse textured, well drained, and has high organic matter content.

### Soil pH

Soil acidity/alkalinity is measured in units of pH, where the lower the pH the more acidic the soil, with a pH of 7 being neutral. Blueberries prefer a soil pH of 4.5 and will tolerate pH levels as high as 5.5. The pH of typical garden soils in Utah is above 7.5. The pH scale is an exponential measurement of the amount of acidic cations (positively charged ions) in solution, so a pH

difference of 1 unit equates to a 10-fold difference in acidic cation concentration. In other words, blueberries require an acidic cation concentration 1000 times higher than typically found in Utah soils. Soil pH determines the relative availability of essential elements. When the pH becomes either too high or too low, the plant usually shows nutrient deficiencies. In Utah’s high pH soils, this is commonly seen as iron chlorosis or zinc deficiency.

### Soil Acidification

In some cases, soil pH can be lowered by the addition of acids or materials such as sulfur that form acids in the soil. However, many Utah soils are “buffered” at higher pH by the presence of carbonates, bicarbonates, and cations such as calcium and magnesium. These soil buffers would need to be saturated by the addition of acid before the soil pH will begin to decrease. The observed consequence of this buffering capacity is that large amounts of acid must be added before the soil pH changes appreciably (Figure 1).




Figure 1. pH buffering effects in Utah soils.

Typical Utah soils contain lime (calcium carbonate) ranging from 1 percent to greater than 20 percent of the soil weight. One percent lime is equivalent to one pound of lime per square foot, and would require nearly a pound of sulfuric acid, or 4.7 ounces of elemental sulfur per square foot, just to neutralize the lime. In the case of soils with 20 percent lime content, 20 times that amount (5.9 pounds of elemental sulfur per square foot) would be required to neutralize the soil's buffering capacity. Once this buffering capacity has been satisfied, additional acid is required to lower the pH. The amount of acid required is dependant upon soil type and texture. A routine soil test, along with a test for percent lime, will provide the information needed to determine the potential for pH adjustment in your soil (see Additional Resources section below).

Acidifying soils may create additional problems. When soil pH is adjusted, some minerals in the soil become more soluble, and thus more available to the plant. In the case of aluminum and manganese, this increased solubility may result in plant toxicity.

## Soil Texture

The fine fibrous root system of blueberry plants requires a coarse-textured soil. Blueberry root growth in acidified soils is often restricted by fine-textured soils (clay). The pH, nutrient availability, and physical properties (texture) of the soil environment can be improved using organic amendments. Interestingly, in some commercial plantings, repeated side-dress applications of organic amendments, such as wood chips, results in a mound of decomposing organic matter that essentially replaces the existing soil as a substrate for root growth (Figure 2).



Figure 2. Highbush blueberries growing in a raised bed made up of decomposing wood chips, placed on the surface of a clay loam soil.

## Soil Replacement

With the difficulty of soil acidification, it might be easier and more cost effective to substitute the existing soil with a media that has the texture, organic matter content, and soil pH suitable for blueberry root growth. This can be accomplished by establishing windrows of organic matter, as shown in Figure 2, or by planting in pots containing a suitable potting medium. However, growing blueberries in a soil substitute is not without difficulties. First, the broad shallow root system is not well-suited to restricted root volumes such as in pots. Further, root zone temperatures in containers and raised beds will often exceed that which is optimal for blueberry roots. For growing in pots, use a 5 to 7 gallon pot that is wider than it is tall, and that is either light colored, or protected from direct sunlight. Finally, like the soils, Utah water is also alkaline and is buffered by the presence of dissolved carbonates. In order for the soil substitute to remain a favorable environment for root growth, irrigation water may also need to be acidified.

## Water Acidification

Alkaline irrigation water can be neutralized or acidified for the maintenance of pH when growing blueberries in acidic soil materials or growing media (peat, potting mixes, etc.). This is done by mixing acid (usually sulfuric acid) directly with water in appropriate amounts, or by oxidizing (burning) elemental sulfur in the presence of water to form sulfuric acid which is then mixed with the irrigation water (known as the sulfur burner process). Homeowner-scale sulfur burners are not available, so the process is generally restricted to commercial production.

Handling concentrated acid solutions is a difficult and potentially hazardous procedure and should only be undertaken with the utmost care and according to the safety precautions outlined by the manufacturer of the acid product. Always use rubber gloves and eye protection when handling acid. When mixing acid and water, ALWAYS add acid to water, NEVER add water to acid as this can be a violent, volatile reaction!

Typical water sources in Utah contain from 60 to 1200 mg/L (or ppm) dissolved carbonate and bicarbonate which buffers pH between 7 and 8. Neutralizing the alkalinity of the water and reducing the pH to about 5 (optimal for blueberries) requires from 160 to 3,175 pounds of sulfuric acid per acre foot of water. This is about 16 to 355 gallons of fertilizer-grade (62%) sulfuric acid (or 10 to 220 gallons of concentrated (98%) sulfuric acid) per acre foot of water treated. For the home garden, this would be equivalent to 0.5 to 8 teaspoons of fertilizer grade sulfuric acid per 100 gallons of water. The water will need to be treated each time the blueberry plants are irrigated. The amount added is critical, as too much acid will lower the pH below 4.0

and can damage the blueberry plants. It should also be noted that for secondary irrigation water sources, the carbonate and bicarbonate content varies over the course of the season. Therefore, the amount of acid required to treat your irrigation water will vary somewhat from spring to fall. For best results, you may need to invest in a test kit or pH meter to measure water pH, then monitor your treated irrigation water and periodically adjust the amount of acid added.

## Varieties and Plant Sources

In addition to the lowbush, northern highbush and rabbiteye types mentioned previously, additional blueberry types have been created through plant breeding. The southern highbush results from crosses between northern highbush and rabbiteye. The half-high varieties are the result of crosses between northern highbush and lowbush types. These general types differ in plant size, fruit ripening time, and most importantly in cold hardiness. For colder parts of the state, the half-high and northern highbush types would be best adapted. For the Dixie area, southern highbush and rabbiteye would be best adapted.

The following varieties are a good starting point. They have been included based upon vigor, toughness, and reliability. Many other varieties are available if these prove successful under your specific conditions. The different types are listed from most cold hardy (half-high) to least cold hardy and most heat tolerant (rabbiteye).

### Half-high varieties:

Friendship – a native selection from Wisconsin that is a taller (chest high), more vigorous bush than typical half-high types. Early ripening, with a medium to medium-small fruit size.

Northblue – a knee-high selection from Minnesota with a spreading bush habit. Blooms later than some other half-highs so it may offer better frost avoidance, but is still relatively early ripening. Medium-sized fruit with good fruit quality.

Northland – a mid-height (waist high) Michigan variety often grown by smaller growers in the northeastern U.S. Cold tolerant and productive, but it has relatively soft fruit that is not well-suited to shipping. Early ripening, with medium-sized fruit.

Northsky – a low, bushy, Minnesota variety with a growth habit somewhat like lowbush blueberry. Early ripening, with a medium to medium-small fruit size.

### Highbush varieties:

Bluecrop – one of the most widely grown varieties. It ripens in midseason, is a broadly-adapted, reliable producer year after year, with classic blueberry flavor.

Blueray – is a sibling of ‘Bluecrop’ with similar recommendations regarding ease of growing. ‘Blueray’ ripens slightly earlier, but is also slightly less productive. Herbert – a late ripening variety that has a vigorous bush and very good flavor. Productive, but fruit are less firm than many other varieties.

Lateblue – one of the latest ripening varieties. Plants are vigorous, upright, and productive year after year.

### Southern highbush:

Echota – an extremely vigorous North Carolina variety with a somewhat sprawling bush habit, but nice quality fruit. Ripens in late midseason.

Georgiagem – an upright bush with reliable production under many different conditions. Ripens in midseason and has medium-sized fruit.

Legacy – a New Jersey variety adapted to a wide range of conditions. Plants may be a bit brushy when young, but develop into an upright, vigorous plant. Very good flavor and fruit quality with a very productive mid- to late harvest season.

Star – a Florida variety that also has good cold adaptation. Large, attractive, early season fruit.

Rabbiteye (rabbiteye types ripen later than almost all highbush types under typical conditions, notes on ripening below refer to their relative order within the rabbiteye ripening season):

Baldwin – a very productive, late-ripening variety with a long harvest season and a sprawling bush habit.

Ira – a newer North Carolina midseason variety, ‘Ira’ is productive with very good quality fruit.

Montgomery – another new North Carolina midseason variety. ‘Montgomery’ has very good fruit quality and more volatile flavors than most typical rabbiteye.

Powderblue – a widely-grown, midseason, Georgia variety with upright habit and good fruit quality.

Tifblue – a classic Georgia variety that combines vigor, cold tolerance, and productivity. An upright type with considerable suckering, but reliable mid-late yields. Fruit is medium sized and light blue.

## General Culture

Healthy blueberry plants will form a medium-sized multi-stem shrub within 3 to 4 years, with a flush of new healthy shoots coming from the crown each year. Plants should be spaced 2.5 to 4 feet apart within the row, with rows 8 to 10 feet apart. More vigorous varieties such as the rabbiteye types may need more space.

Blueberries are sensitive to both drought and waterlogging. The porous soils that blueberries prefer will drain quickly, but also will dry out quickly, so the plants will need to be irrigated frequently. If amending an existing garden soil with organic matter, use raised beds to improve drainage and soil aeration. A raised bed 3 feet wide and 8 to 12 inches high will be sufficient.

Blueberries are also sensitive to over fertilization. Apply 2 tablespoons of 10-20-10 fertilizer per plant at planting. In subsequent years, apply one to four ounces of fertilizer per plant (depending on plant size) in both the spring and the fall. Fertilizer nitrogen can be in either the ammonium or nitrate form. Blueberries best utilize the ammonium form. Where possible, use balanced fertilizers that contain ammonium nitrogen. Acid fertilizers, such as those formulated for azaleas, generally contain ammonium nitrogen. Alternatively, use ammonium sulfate to supply fertilizer nitrogen needs.

During the first growing season, do not allow the blueberry plants to produce fruit as this will slow establishment. To prevent fruiting, pick off the flower clusters as they emerge in the spring.

Once established and fruiting, blueberry plants will need to be pruned to maintain productivity. Each spring as the buds begin to swell, prune out low growing shoots around the base, and any dead wood, or small non-vigorous shoots. Select for vigorous (larger) shoots with lateral branches and bright colored bark. After removing the small twigs and basal growth, additional renewal pruning may be required to thin out remaining canes and make room for the establishment of new more productive canes. The degree of renewal pruning required will depend on the health and vigor of the plant. Where plants are extremely vigorous, approximately 1/3 of the wood should be removed annually. Typically this would entail removing two or three of the largest oldest canes. With less vigorous varieties or on less vigorous sites approximately 1/5 to 1/6 of the wood (one or two of the oldest canes) will need to be removed annually.

## Additional Resources

Highbush Blueberry Production: Oregon State University Extension, Publication PNW215.  
<http://extension.oregonstate.edu>.

Additional online information available at:

[www.fallcreeknursery.com/Gardener/index.htm](http://www.fallcreeknursery.com/Gardener/index.htm), Fall Creek Nursery.

[www.extension.umn.edu](http://www.extension.umn.edu), The University of Minnesota.

The USU Analytical Laboratory (USUAL) maintains an information Web site at: [www.usual.usu.edu](http://www.usual.usu.edu). Prices and sample submission forms for soil and water testing services can be found at the Web site, or by visiting your local county Extension office.

**Nursery sources:** Following is an alphabetical list of blueberry nurseries, including location, phone number, and featured blueberry cultivars. Except where noted, these nurseries sell small quantities of plants directly to home gardeners.

Bottoms Nursery, Concord, Georgia. Phone 770-884-5661.  
Georgiagem, Star, Baldwin, Powderblue, Tifblue.

DeGrandchamps Farms, South Haven, Michigan.  
888-483-7431. Friendship, Northblue, Star.

Finch Blueberry Nursery, Bailey, North Carolina.  
252-235-4664. Bluecrop, Blueray, Echota, Georgiagem, Legacy, Baldwin, Ira, Montgomery, Powderblue, Tifblue.

Hartmann's Plant Company, Lacota, Michigan  
269-253-4281. Bluecrop, Blueray, Northblue, Northsky, Legacy.

Heartland Blueberries, Hodgenville, Kentucky.  
270-358-5837. Bluecrop, Blueray, Star, Georgiagem, Legacy.

Ison's Nursery, Brooks, Georgia. 800-733-0324.  
Powderblue, Tifblue.

Miller Nurseries, Canandaigua, New York 800-836-9630.  
Bluecrop, Blueray, Northblue, Herbert.

Morrison's Home and Garden, Plymouth Massachusetts.  
508-746-0970. Bluecrop, Blueray, Lateblue

Octoraro Farm and Gardens, Nottingham, Pennsylvania. 610-932-0225. Friendship, Herbert, Lateblue

Oregon Blueberry Farms, Silverton, Oregon. 503-873-1200.  
Bluecrop, Legacy, Powderblue.

Stark Brothers. Louisiana, Missouri. 800-325-4180. Bluecrop, Blueray, Legacy, Northblue, Tifblue.

### Commercial sales only:

Fall Creek Nursery, Lowell, Oregon. 800-538-3001.  
Bluecrop, Blueray, Legacy, Northblue.

Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran's status. USU's policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions.

Utah State University employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran's status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USU-sponsored events and activities.

This publication is issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Noelle E. Cockett, Vice President for Extension and Agriculture, Utah State University.