



Eriophyid Mites bud, blister, gall, and rust mites

Ryan S. Davis, Arthropod Diagnostician and Taun Beddes, Cache County Horticulture Agent

WHAT YOU SHOULD KNOW

- Eriophyid mites cannot be seen without a 20x hand lens or greater magnification.
- Eriophyid mites seldom cause serious injury or stress to plants; damage is normally aesthetic.
- Damage from eriophyid mites usually consists of leaf galls, bud or flower galls, blisters, scabbing, and deformities of leaves, stems, buds, and flowers.
- Control techniques include pruning, plant removal, insecticidal soap, horticultural oils, predatory mites, and broad-spectrum insecticides and miticides.

INTRODUCTION

Eriophyid mites are translucent, cigar-shaped microscopic mites that cause deformities on many plants species. These mites are noticed when their feeding causes abnormalities of plant tissues such as erineum, galls, brooms, leaf curling, blisters, rusts, silvering, fruit rusetting, and deformed buds, catkins, fruits, etc. Fortunately, these mites rarely cause serious harm to plants, and control is seldom needed. In fact, eriophyid mites can serve as an alternative food source for predatory mites when their primary food source is absent. In that way, eriophyid mites can actually help reduce outbreaks of spider mites and other insects on which predatory spider mites feed, by allowing predatory mites to survive in the absence of their primary food. There are 1,859 species of eriophyid mites




Fig. 2. Microscopic view of an eriophyid mite (*Aceria anthocoptes*).²

described from around the world with many species still undescribed. Needless to say, you are likely to encounter these mites some time in your gardening future.

GENERAL BIOLOGY

Family Name: Eriophyidae

Range: Worldwide

Hosts: Some mites are host specific, while others attack many plant, tree, and shrub species (Table 2).

Identification of Adult: Nearly invisible to the unaided eye (1/100" in length), cylindrical/cigar-shaped, tapering from head to rear, and translucent white (Fig. 1). Unlike most mites, eriophyids only have 4 legs located near the head.

Life History: Mites generally overwinter as fertilized adult females under bud scales, or protected sites on or near the host plant, and emerge at bud break in spring. Both males and females are present throughout the growing season. Reproduction is continuous, with generations completed every 2-3 weeks; overlapping generations are present.

Key Habits: Overwintering females emerge from under bud scales and other protected sites to lay eggs and feed on new foliage when buds begin to break in the spring. This time is most adventitious for insecticide applications. Some may transmit viruses that can deform plants or




Fig. 1. Citrus rust mite (*Phyllocoptura oleivora*).¹

© aggie-horticulture.tamu.edu

GENERAL BIOLOGY CONTINUED

cause economic losses to various crops.

Damage description:

Below are pictures of symptoms caused by eriophyid mites categorized by plant part.



Fig. 3. Galls caused by boxelder pouchgall mite (*Eriophyes negundi*).³



Fig. 4. Leaf blisters on pear caused by pearleaf blister mite (*Phytoptus pyri*).⁴



Fig. 5. Erieneum on Rocky Mountain Maple caused by *Eriophyes calaceris*.⁵



Fig. 6. Fingergalls on chokecherry caused by *Phytoptus emarginate*.⁴




Fig. 7. Bronzing of pear leaves caused by pear rust mite (*Eptrimerus pyri*).⁶



Fig. 8. Russetting of pear fruit by pear rust mite (*Eptrimerus pyri*).⁶



Fig. 9. Cottonwood catkin gall mite (*Eriophyes newesse-gi*).³



Fig. 10. Ash flower gall mite (*Eriophyes fraxiniflora*).¹



Fig. 11. Hackberry witches broom caused by *Eriophyes celtis*.³



Fig. 12. Example of a stem gall mite (*Aceria* sp.) on maple.⁸



Fig. 13. Poplar bud gall caused by *Eriophyes parapopuli*.³



Fig. 14. Damage caused by rose rosette virus transmitted by *Phyllocoptes fructiphilus*.⁴

PHOTO CREDITS

- ¹. Texas A&M University, <http://cookislands.bishopmuseum.org/species.asp?id=9294>.
- ². Eric Erbe, USDA Agricultural Research Service, Bugwood.org.
- ³. Whitney Cranshaw, Colorado State University, Bugwood.org.
- ⁴. Keifer, H.H., Baker, E.W., Kono, T., Delfinado, M., and Styer, W.E. An Illustrated Guide to Plant Abnormalities Caused by Eriophyid Mites.
- ⁵. Ciesla, W.M. Forest Health International. Bugwood.org.

- ⁶. Clark, J.C. University of California Statewide IPM Project.
- ⁷. Solomon, J. USDA Forest Service. Bugwood.org.
- ⁸. Csoka, G. Hungary Forest Research Institute. Bugwood.org.

CONTROL

eriphyid mite damage occurs on many plant species and often does not impact long term health. Because of this, chemical pesticides should not be considered a primary management option.

Non-Chemical Control

Cultural control practices that modify the growing environment can reduce the occurrence of unwanted pests.

- **Pesticide avoidance:** Overuse of insecticides and miticides can eliminate natural predators, and can lead to pest resistance, making long-term control difficult.
- **Monitor/scout:** Periodic examination of plant material reveal Eriophyid populations before they increase to damaging numbers.
- **Plant selection:** Use plants that are less susceptible to eriophyid mites, and that are adapted to Utah's arid climate; contact your local USU Extension office for recommendations. Make sure plants or planting material you select are free from eriophyid mites.
- **Prune:** Remove infested leaves or branches and remove from the property.
- **Plant removal:** Heavily infested plants such as tomatoes and wheat, may need to be removed to reduce the spread of mites. Some mites have secondary host plants, so removal of host weeds minimizes sources of re-infestation.

Chemical Control

If severe infestations are causing major aesthetic damage, tree stress, or crop economic loss, insecticides/miticides may be necessary. Consider reduced-risk pesticides to limit negative impacts on beneficial insects and the environment. Occasionally, broad-spectrum insecticides/miticides, or those that kill not only your target pest, but many other arthropods, may be necessary for quick reductions in mite numbers. The downside of broad-spectrum insecticides is that they can eliminate predators and lead to chemical resistance in eriophyid mites and other insect populations (if used improperly).

Reduced-Risk Chemicals

- **Horticultural soaps and oils:** Apply dormant weight horticultural oils 7-10 days before bud break and again at bud break. Proper timing targets eriophyid mites and preserves beneficial arthropods. During the summer soaps and summer weight oils offer mite control or suppression. Proper timing can also target other common plant pests such as aphids. Test for phytotoxicity before spraying the whole plant.
- **Kaolin clay:** A fine clay applied to plants in a liquid solution. Kaolin clay acts as a feeding deterrent, and

requires full coverage of foliage (top and bottom) to be effective. Frequent applications are needed, but may cause plant stress if used too often (inhibits photosynthesis).

- **Sulfur:** Sulfur has long been used for mite control. Full coverage is necessary for control. Do not apply when temperatures exceed 90°F, or during periods of high humidity.
- **Neem oil:** Azadirachtin (group unknown) is a botanical insect growth regulator acquired from neem trees.
- **Predatory mites:** Naturally occurring predatory mites often feed on Eriophyid mites and can keep populations at tolerable levels. Avoid the use of broad-spectrum insecticides to preserve predatory mites. Low or extinct populations of predatory mites may be supplemented with predatory mites available for order. One good option for hot arid regions is *Galendromus occidentalis* (western predatory mite). Other species of predatory mites are available, but work best in humid conditions, such as in a greenhouse.

Insecticides and Miticides

Table 1 contains an abbreviated list of active ingredients effective against Eriophyid mites. Make sure the host plant is listed on product label you choose.

- When using pesticides, rotate the chemical group on a yearly basis to avoid insect resistance. Chemical groups are listed in Table 1.
- For gall-forming mites, once visible damage such as galls are observed, it is often too late to treat during that growing season.

Active Ingredient	Chemical Group
abamectin	6
bifenthrin	3A
carbaryl	1A
deltamethrin	3A
imidacloprid	4A
malathion	1B
permethrin	3A
pyrethrin	3A

Table 1. Broad-spectrum insecticides for Eriophyid mite control, and their chemical group. Use a different chemical group every year to minimize pest resistance.¹

COMMON ERIOPHYID MITES AFFECTING PLANTS IN UTAH

Common Name	Scientific Name	Host Plants	Damage Description
tomato russet mite	<i>Aculops lycopersici</i>	tomato, nightshade plants (Solanaceae)	new infestations turn lower stems and leaves brown, eventually withering; russetting of tomato fruit; death
honeylocust rust mite	<i>Aculops spp.</i>	honeylocust	feeding on undersides of leaves causes bronzing that is visible by late summer
peach silver mite	<i>Aculus cornutus</i>	peach, nectarine, almond	silvering of leaves in late summer before leaf drop
willow pouchgall mite	<i>Aculops tetanothrix</i>	willow	2-3 mm diameter irregular and beadlike galls usually located near the leaf midrib or between veins; gall color is pink to red-purple to yellow and ranges from fuzzy, rough, to smooth; galls may cover whole leaf
apple rust mite	<i>Aculus schlechtendali</i>	apple	injured terminal growth; lengthwise leaf curl; fruit russetting
privet rust mite	<i>Asculus ligustri</i>	privet	scratched appearance on top of leaves turning brown; leaf curl on young leaves
grape erineum mite	<i>Colomerus vitis</i>	grape	deformation of bud clusters; felty erineum on lower leaf surface followed by blister-like swellings on upper leaf
Rocky Mountain Maple erineum mite	<i>Eriophyes calaceris</i>	Rocky Mountain Maple	greenish yellow to pink, crimson, or purple-red erineum on upper leaf surface, usually missing the major veins (Fig. 5)
hackberry mite	<i>Eriophyes celtis</i>	hackberry	thin, short, stunted, and tightly bunched twigs resulting in witches' brooms (Fig. 11)
ash flowergall mite	<i>Eriophyes fraxiniflora</i>	ash	inflorescence gall (Fig. 10)
boxelder pouchgall mite	<i>Eriophyes negundi</i>	boxelder	pouch-like galls on upper leaf surface created by indentations on the underside of leaves, filled with fine white hairs-galls are mostly solitary, located between veins
cottonwood catkin gall mite	<i>Eriophyes newesegi</i>	poplar (<i>Populus spp.</i>)	grape-like growths/distortions of catkins; affected catkins remain on tree until mid summer (Fig. 9)
poplar budgall mite	<i>Eriophyes parapopuli</i>	poplar	irregular, asymmetrical, bumpy, solid mass of fleshy swelling of the bud (Fig. 13)
wheat curl mite/bulb mite	<i>Eriophyes tulipae</i>	barley, oats, corn, wheat, <i>Poa spp.</i> , <i>Agropyron spp.</i> , foxtail barley, onion, garlic, tulip	leaf curling; red-streaked corn; leaf margin curling; brown blistering on bulbs; can transmit wheat streak mosaic virus
aspen gall mite	<i>Phyllocoptes didelphus</i>	quacking aspen	circular, shallow galls, 5-15 mm in diameter; pouch-like galls on upper side of leaf; pockets on undersides of leaves are filled with yellowish leaf growth
chokecherry fingergall mite	<i>Phytoptus emarginate</i>	chokecherry, various plum species (<i>Prunus spp.</i>)	elongate, finger-like galls scattered over upper surface of leaf; green to yellow turning brown (Fig. 6)
alder pouchgall mite	<i>Phytoptus laevis</i>	alder (<i>Alnus spp.</i>)	multi-sized, roundish, green-yellow to reddish-brown galls located on both sides of leaves
appleleaf blister mite pearleaf blister mite	<i>Phytoptus mali</i> <i>Phytoptus pyri</i>	apple pear	blister-like galls on upper surface of leaves; yellow-green turning brown; brown, blister-like lesions on fruit (Fig. 4)
mountain-ash pouchgall mite	<i>Phytoptus sorbi</i>	mountain-ash	blister-like galls on the upper surface of leaves; yellow-green turning brown
pine needle eriophyid mite	<i>Trisetacus spp.</i>	pine (<i>Pinus spp.</i>), Douglas-fir	clustering of aborted buds; stunted needles; needle yellowing or browning

Precautionary Statement: Utah State University Extension and its employees are not responsible for the use, misuse, or damage caused by application or misapplication of products or information mentioned in this document. All pesticides are labeled with active ingredients, directions for use, and hazards, and not all are registered for edible crops. "Restricted use" pesticides may only be applied by a licensed applicator. The pesticide applicator is legally responsible for proper use. USU makes no endorsement of the products listed herein.

Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran's status. USU's policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions. Utah State University employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran's status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USU-sponsored events and activities. This publication is issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Dept. of Ag., Noelle E. Cockett, Vice President for Extension and Agriculture, Utah State University.

Fact Sheet Series: Insects - Landscape and Ornamental; Tree Fruit; Small Fruit